Small-angle X-ray scattering from biomacromolecular solutions

Dmitri Svergun

EMBL

Biological SAXS @ EMBL-HH

Group leader: D. Svergun
Staff : M. Petoukhov, C.Blanchet, D.Franke, A.Kikhney, H.Mertens
Postdocs: A.Tuukkanen, M.Graewert , A.Spilotros, C.Jeffries, A.Panjkovich
Predocs: G.Tria, M.Kachala, E.Valentini, N.Hajizadeh

Major tasks:

- Development of data analysis methods
- Running and developing SAXS beamlines
- User support and collaborative projects
- Education and training

Modern synchrotron SAXS

Small-angle scattering: experiment



Monochromatic beam

Wave vector ***k***, $k=2\pi/\lambda$

Radiation sources:

X-ray generator (λ = 0.1 - 0.2 nm)
Synchrotron (λ = 0.03 - 0.35 nm)
Thermal neutrons (λ = 0.2 - 1 nm)

Scattering vector $\mathbf{s}=\mathbf{k}_1-\mathbf{k}$, $s=4\pi\sin\theta/\lambda$

Small-angle scattering: contrast



$I_{sample}(s)$ – $I_{matrix}(s)$ = $I_{particle}(s)$

- To obtain scattering from the particles, matrix scattering must be subtracted, which also permits to significantly reduce contribution from parasitic background (slits, sample holder etc)
- **Contrast** $\Delta\rho = \langle\rho(r) - \rho_s\rangle$, where ρ_s is the scattering density of the matrix, may be very small for biological samples

Crystal versus solution

Crystal versus solution

- In solution, no crystallographic packing forces are present

- For SAXS solution studies, one does not need to grow crystals
- SAXS is not limited by molecular mass and is applicable under nearly physiological conditions
- Using solution SAXS, one can more easily observe responses to changes in conditions
- SAXS permits for quantitative analysis of complex systems and processes


The scattering is related to the shape
(or low resolution structure)
Solid sphere
lg I(s), relative
0
-1
-2
-3
-4
-5
-6
0.0
0.1
0.2
0.3
0.4
0.5
s, nm-1
Hollow sphere
Dumbbell
Flat disc
Long rod


When biologists
go for SAXS
SAXSMAN © A.Kikhney
This is just trivial case:
SAS yields much more
Crystallization
Care for a
shape?

Some words of caution



- Sample preparation
- Experiment
- Data processing
- Unambiguous interpretation
- Changing conditions
- Relation to function

Methods development at EMBL-Hamburg



Data processing and manipulations
Rigid body refinement

Employed by over 9500 users worldwide



Ab initio modeling suite
Analysis of mixtures





Konarev, P.V., Petoukhov, M.V., Volkov, V.V. & Svergun, D.I. (2006). *J. Appl. Cryst.* **39**, 277

Scattering from dilute macromolecular solutions (monodisperse systems)

$$I(s) = 4\pi \int_0^D p(r) \frac{\sin sr}{sr} dr$$

The scattering is proportional to that of a single particle averaged over all orientations, which allows one to determine size, shape and internal structure of the particle at low (*1-10 nm*) resolution.

Overall parameters





"Simple" monodisperse systems



Shape and conformational changes of macromolecules and complexes

Validation of high resolution models and oligomeric organization



Rigid body models of complexes using high resolution structures



Addition of missing fragments to high resolution models

Ab initio methods





Bead (dummy atoms) model

A sphere of radius D_{max} is filled by densely packed beads of radius $r_0 << D_{max}$



Vector of model parameters:

$$\text{Position } (j) = X(j) = \begin{cases} 1 & \text{if particle} \\ 0 & \text{if solvent} \end{cases}$$

(phase assignments)

Number of model parameters $M \approx (D_{max} / r_0)^3 \approx 10^3$ is too big for conventional minimization methods – Monte-Carlo like approaches are to be used

But: This model is able to describe rather complex shapes

Chacón, P. *et al.* (1998) *Biophys. J.* 74, 2760-2775.

Svergun, D.I. (1999) *Biophys. J.* 76, 2879-2886

How to reconstruct 3D from 1D



The 3D model is required not only to fit the data but also to fulfill (often stringent) physical and/or biochemical constrains



Poly(ethylene) glycol-conjugated hemoglobin: implications for a new oxygen therapeutic



SAXS visualizes PEG tails and interactions in Hemospan®, an oxygen therapeutic with increased performance



Svergun DI, Ekström F, Vandegriff KD, Malavalli A, Baker DA, Nilsson C & Winslow RM (2008) *Biophys J.* **94**, 173-181.

Shapes from recent projects at EMBL-HH

Complexes and assemblies

S-layer proteins

Fagan *et al* Mol. Microbiol (2009)

α-synuclein oligomers



Giehm *et al* PNAS USA (2011)

Domain and quaternary structure

Dcp1/Dcp2 complex



She *et al*, Mol Cell (2008)

Toxin B

Albesa-Jové *et al* JMB (2010)

Flexible/transient systems

Complement factor H



Morgan *et al* NSMB (2011)

Structural transitions

Cytochrome/adrenodoxin



Xu et al JACS (2008)

Src kinase



Bernado et al JMB (2008)

Modern life sciences widely employ hybrid methods

The most known and popular tool is, of course, Photoshop

SAXS also allows for a very effective hybrid model building where high resolution portions are positioned to fit the low resolution scattering data

Catalytic core of E2 multienzyme complex is an irregular 42-mer assembly



The E2 cores of the dihydrolipoyl acyl-transferase (E2) enzyme family form either octahedral (24-mer) or icosahedral (60-mer) assemblies. The E2 core from Thermoplasma acidophilum assembles into a unique 42-meric oblate spheroid. SAXS proves that this catalytically active 1.08 MDa unusually irregular protein shell does exists in this form in solution.



Marrott NL, Marshall JJ, Svergun DI, Crennell SJ, Hough DW, Danson MJ & van den Elsen JM. (2012) *FEBS J.* **279**, 713-23

Principle of rigid body modelling



Using spherical harmonics, the amplitude(s) of arbitrarily rotated and displaced subunit(s) are analytically expressed *via* the initial amplitude and the six positional parameters: $C_{lm}(s) = C_{lm}(B_{lm}, \alpha, \beta, \gamma, x, y, z)$.

The scattering from the complex is then rapidly calculated as

$$I(s) = I_A(s) + I_B(s) + 4\pi^2 \sum_{0}^{\infty} \sum_{-l}^{l} \mathrm{Re}\left[A_{lm}(s) C_{lm}^*(s)\right]$$

Svergun, D.I. (1991). *J. Appl. Cryst.* **24**, 485-492

A global refinement run with distance constraints

A tyrosine kinase MET (118 kDa) consisting of five domains

Program SASREF



Single curve fitting with distance constraints: C to N termini contacts

Gherardi, E., Sandin, S., Petoukhov, M.V., Finch, J., Youles, M.E., Ofverstedt, L.G., Miguel, R.N., Blundell, T.L., Vande Woude, G.F., Skoglund, U. & Svergun, D.I. (2006) *PNAS USA,* **103**, 4046.

Architecture of nuclear receptor heterodimers on DNA direct repeat elements

Nuclear hormone receptors (NHRs) control numerous physiological processes through the regulation of gene expression. SAXS, SANS and FRET were employed to determine the solution structures of NHR complexes, RXR–RAR, PPAR–RXR and RXR–VDR, free and in complex with the target DNA



RXR–RAR–DR5

Ab initio and rigid body models of NHRs complexed with direct repeat elements

RXR–VDR–DR3



N.Rochel, F.Ciesielski, J.Godet, E.Moman, M.Roessle, C.Peluso-Iltis, M.Moulin, M. Haertlein, P.Callow, Y.Mely, D.Svergun & D.Moras (2011) *Nat Struct Mol Biol* 18, 564-70

Architecture of nuclear receptor heterodimers on DNA direct repeat elements

NHR-DNA complexes show extended asymmetric shape and reveal conserved position of the ligand-binding domains at the 5′ ends of the target DNAs. Further, the binding of only one coactivator molecule per heterodimer, to RXR's partner, is observed.



RAR–RXR-DR1
PPAR–RXR-DR1





The models and the polarity of RXR–RAR–DR5 and RAR–RXR–DR1 were validated using neutron scattering and FRET



N.Rochel, F.Ciesielski, J.Godet, E.Moman, M.Roessle, C.Peluso-Iltis, M.Moulin, M. Haertlein, P.Callow, Y.Mely, D.Svergun & D.Moras (2011) *Nat Struct Mol Biol* 18, 564-70

Addition of missing fragments

- Flexible loops or domains are often not resolved in high resolution models
- Their tentative configuration can be reconstructed by fixing the known portion and adding the missing parts to fit the scattering from the full-length macromolecule.

- Moreover, addition of missing fragments can be combined with rigid body refinement (programs BUNCH and CORAL)

Petoukhov, M. V. & Svergun, D. I. (2005). *Biophys. J.* **89**, 1237-1250



C-terminal flexibility of the influenza virus M1 protein

Influenza A virus matrix protein M1 is one of the most important and abundant proteins in the virus particles. The low resolution SAXS models reveal a polarized M1 molecule consisting of a compact NM-fragment, compatible with its crystal structure, and an extended and partially flexible C-terminal domain.







The M1 monomers co-exist in solution with a small fraction of large assemblies revealing a layered architecture similar to that observed in the authentic influenza virions. The flexibility of the C-terminus is an essential feature relevant for the multifunctionality of the entire protein.

E.V. Shtykova, L.A. Baratova, N.V. Fedorova, V.A. Radyukhin, A.L. Ksenofontov, V. V. Volkov, A.V. Shishkov, A.A. Dolgov, L.A. Shilova, O.V. Batishchev, C.M. Jeffries, D.I. Svergun (2013), PLoS One, 8(12):e82431.



Life is more complicated: mixtures and processes



Equilibrium oligomeric mixtures

Stoichiometry and complex formation



Natively unfolded proteins and multidomain proteins with flexible linkers



Protein folding/unfolding kinetics



Assembly/disassembly processes

Scattering from mixtures

$$I(s)=\sum_k v_k I_k(s)$$

The scattering is proportional to that of a single particle averaged over all orientations, which allows one to determine size, shape and internal structure of the particle at low (*1-10 nm*) resolution.

For equilibrium and non-equilibrium mixtures, solution scattering permits to determine the number of components and, given their scattering intensities $I_k(s)$, also the volume fractions

An encounter adrenodoxin/cytochrome C complex



Volume fractions

Mon:Dim:Tri:Tet

0 : 0 : 50 : 50

0 : 8 : 47 : 45

5 : 25 : 55 : 15

25 : 25 : 50 : 0

0 : 100 : 0 : 0 NMR data



A model of heterotetrameric native complex at high salt and high solute concentration



The cross-linked complex Ad/CC is always a heterodimer independent on conditions The native complex strongly depends on the sample concentration and on the amount of NaCl in the buffer.

X. Xu, W. Reinle, F. Hannemann, P. V. Konarev, D. I. Svergun, R. Bernhardt & M. Ubbink (2008) *JACS,* **130**, 6395-6403

Fibrillation of α-synuclein (αSN)

Aggregation of αSN leads to Parkinson decease. The fibril formation process characterized by SAXS reveals that there exists an intermediate oligomer formed by several partially unfolded αSN molecules, and the mature fibril is formed by association of these oligomers. Such mechanism was earlier found for insulin fibrillation.







The oligomer is shown to disrupt liposomes, i.e. it is potentially cytotoxic

Giehm, L., Svergun, D.I., Otzen, D.E. & Vestergaard, B. (2011) *PNAS USA,* **108,** 3246

SAXS from folded *vs* disordered protein





Folded: relatively small R_g and D_{max}, bell-shaped Kratky plot (e.g. for folded α-amylase (448 AAs) R_g=2.4 nm)

Disordered: large R_g and D_{max}, increasing Kratky plot (e.g. for IUP tau (441 AAs) Rg=6.5 nm)

Quantitative assessment of flexibility



- Automated classification (folded, partially or completely unfolded) is available

D.Franke DATCLASS

- More quantitative estimates are provided by ensemble methods
- One generates a large pool covering the conformational space and selects sub-ensemble(s) such that their mixture fits the available experimental data
- The structural properties of the selected ensemble(s) are compared to those of the pool

EOM, Bernadó et al. (2007)
J. Am. Chem. Soc. **129**, 5656.

34

Quantitative assessment of flexibility: EOM 2.0

- Extended possibilities for the pool generation, e.g. use of (partial) point symmetry
- Broader search capabilities
- Quantification of flexibility using enthropy and variation



$$R_{flex} = -\mathrm{H_b(S)} = \sum_{i=1}^{n} p(x_i)\log_b(p(x_i))$$

$$\mathrm{R}\sigma = \sigma_{ensemble} / \sigma_{pool}$$



G.Tria, M.Kachala

Crystal structures of substrate-bound chitinase from *Moritella marina* and its structure in solution

Chitinases break down glycosidic bonds in chitin and only few crystal structures are reported because of the flexibility of these enzymes.

The dimeric crystal structure (at BESSY) of chitinase 60 from M. marina (MmChi60) contains four domains: catalytic, two Ig-like, and chitin-binding (ChBD). SAXS (at EMBL) demonstrates that MmChi60 is monomeric and flexible in solution. The flexibly hinged Ig-like domains may thus allow the catalytic domain to probe the surface of chitin.





P. H. Malecki, C. E. Vorgias, M. V. Petoukhov, D. I. Svergun and W. Rypniewski. *Acta Cryst.* (2014) **D70**, 676-684

Not only proteins: magnetic nanoparticles





Highly monodisperse NPs are prepared by thermal decomposition of iron compounds including oxygen-containing ligands in boiling surfactants. The NPs are coated by phospholipids with PEG Tails to become soluble.

Rigid body analysis reveals equilibrium clusters of the NPs stabilized by magnetic interactions

Ab initio analysis: peculiarities of organization of different NPs





Shtykova, E.V, Huang, X., Remmes, N., Baxter, D., Dixit, S., Stein, B., Dragnea, B., Svergun, D. I. & Bronstein, L. M. (2007) J. Phys. Chem. C, **111**, 18078-18086

Mixed Monolayer Protected Gold Nanoparticles

Gold nanoparticles (NPs) are efficiently synthesized using self-assembled monolayers (SAMs) as stabilizing agents. SAMs of thiolated ligand molecules on gold substrates are ordered two-dimensional crystals. When mixtures of molecules are used, these may spontaneously form domains. The ligand shell provides the NP with important interfacial properties.

Scanning tunneling microscopy on dodecanethiol-hexanethiol (C12 : C6) 2:1 protected gold nanoparticles reveals stripe-like domains persistent across many images and retain their direction and overall morphology at different scan angles



M. Moglianetti, Q.K. Ong, J. Reguera, K.M. Harkness, M. Mameli, A. Radulescu, J. Kohlbrecher, C. Jud, D.I. Svergun, F. Stellacci (2014) Chem. Sci., **5**, 1232-1240

X-rays versus neutrons

- **X-rays:** scattering factor increases with atomic number, no difference between H and D
- **Neutrons:** scattering factor is irregular, may be negative, huge difference between H and D

Element	H	D	C	N	O	P	S	Au
At. Weight	1	2	12	14	16	30	32	197
N electrons	1	1	6	7	8	15	16	79
b_X, 10^{-12} cm	0.282	0.282	1.69	1.97	2.16	3.23	4.51	22.3
b_N, 10^{-12} cm	-0.374	0.667	0.665	0.940	0.580	0.510	0.280	0.760

Neutron contrast variation

Mixed Monolayer Protected Gold Nanoparticles

SANS was performed on the samples with specifically deuterated dodecanethiol (C6:d-C12; red circles) and hexanethiol (d-C6:C12; blue triangles).

Contrast, 10^{-6} Å^{-2}	C6: d-C12	d-C6: C12
Phase 1: gold	1.4	1.4
Phase 2: C12	2.5	-3.4
Phase 3: C6	-3.4	1.9





The scattering patterns and distance distribution functions from the two hybrid particles are distinctly different indicating that SANS sees the internal structure

M. Moglianetti, Q.K. Ong, J. Reguera, K.M. Harkness, M. Mameli, A. Radulescu, J. Kohlbrecher, C. Jud, D.I. Svergun, F. Stellacci (2014) Chem. Sci., **5**, 1232-1240

Mixed Monolayer Protected Gold Nanoparticles

The shape and internal structure of the particles was reconstructed using a multi-phase model depicting the three components (yellow: gold; magenta: C6; cyan: C12 and fitting both data sets from specifically deuterated thiols.





M. Moglianetti, Q.K. Ong, J. Reguera, K.M. Harkness, M. Mameli, A. Radulescu, J. Kohlbrecher, C. Jud, D.I. Svergun, F. Stellacci (2014) Chem. Sci., **5**, 1232-1240

Small-angle scatterers and biologists



Beamyear 2013: 335 user-visits, 291 unique users, 159 projects, 119 unique groups (of those, 10 mail-in/remote and 3 proprietary)

CRG access (HZG): 39 user-visits, 33 unique users, 23 projects, 19 unique groups (NOT counted in the above graph)

EMBL SAXS X33 beamline, 1979-2012



This is how it worked in 2005



High brilliance beamline P12 (Petra-3)

(P)reincarnation:

BioSAXS P12 beamline inherits not only the soul but also fully utilizes the automation developed for X33



Automation of SAS experiment and analysis





Processing with an automated pipeline and interpretation with an expert system

SEC + Malvern TDA online

BioStructX

Malvern



SEC elution profile of BSA on a Superdex 200 column:
- UV detector signal
- Light scattering
- Refractive index







2500 frames collected while protein was eluted from the column

Recent EMBL SAXS collaborative projects

Transcription factors

De *et al* PNAS (2014)

PDH complex E2 core

Marrott *et al* FEBS J (2012)

Human chromatin remodeler CHD4

Watson *et al* JMB (2012)

Flexible trimeric gephyrin

Sander *et al* Acta Cryst D (2013)

Arc1p-aminoacyl-tRNA synthetases



Koehler *et al* NAR (2013)

BARF1/hCSF-1 complex



Elegheert *et al* NSMB (2012)

Colonization factor GbpA



Wong *et al* Plos Pathog (2012)

Nanocomposites

Shtykova *et al* JPC (2012)

What does SAS tell about biological macromolecules



- Nothing known: *ab initio* low resolution structure
- Incomplete high resolution structure known: probable configuration of missing portions
- Complete high resolution structure known: validation in solution and biologically active oligomers
- High resolution structure of domains/subunits known: quaternary structure by rigid body refinement
- Mixtures/assemblies: volume fractions of components
- Flexible systems: quantitative analysis of configurational ensembles

What does SAS tell about biological macromolecules

- Nothing known: *ab initio* low resolution structure
- Incomplete high res… structure kn… configu…
- Com… in s…
- Hig… quat…
- Mixtu… …ents

…ghput SAXS

Not just high throughput: important present and future applications of SAXS are functional complexes and processes (flexible, dynamic, transient, evolving), where SAXS is among the few methods providing quantitative structural information

Syn… …ron SAXS… …terial, concentration from 0.5 mg/ml, exposure times: (sub)seconds

Recent reviews on solution SAS

Blanchet CE, Svergun DI (2013) Small-angle X-ray scattering on biological macromolecules and nanocomposites in solution. Annual Review of Physical Chemistry 64(1): 37–54.

Schneidman-Duhovny D, Kim S, Sali A. (2012) Integrative structural modeling with small angle X-ray scattering profiles. BMC Structural Biology 12(1):17.

Graewert MA, Svergun DI (2013) Impact and progress in small and wide angle X-ray scattering (SAXS and WAXS). Curr Opin Struct Biol 23: 748-754.

Rambo RP and Tainer JA (2013) Super-resolution in solution X-ray scattering and its applications to structural systems biology., Annu Rev Biophys. 42, 415-441

Books on SAXS

" The origins" (no recent edition) : Small Angle Scattering of X-rays. A. Guinier and A. Fournet, (1955), in English, ed. Wiley, NY

Small-Angle X-ray Scattering: O. Glatter and O. Kratky (1982), Academic Press. PDF available on the Internet at http://physchem.kfunigraz.ac.at/sm/Software.htm

Structure Analysis by Small Angle X-ray and Neutron Scattering. L.A. Feigin and D.I. Svergun (1987), Plenum Press. PDF available on the Internet at http://www.embl-hamburg.de/ExternalInfo/Research/Sax/reprints/feigin_svergun_1987.pdf

Small Angle X-Ray and Neutron Scattering from Solutions of Biological Macromolecules. D.I,Svergun, M.H.J. Koch, P.A.Timmins, R.P. May (2013) Oxford University Press, London.

Young scientists were always very interested in SAXS



… in preparation for EMBL PhD program … (courtesy of F.Gabel, IBS, Grenoble)

The brand new SAS book is not yet available for everyone (courtesy of M.Graewert, EMBL-HH)

Acknowledgments



EMBL-Hamburg: D.Franke, A. Kikhney, P. Konarev, M. Petoukhov, M. Roessle, C.Blanchet, H.Mertens, A.Spilotros, G.Tria, M. Graewert, C.Jeffries, A.Tuukkanen, M.Kachala, E.Valentini

http://www.embl-hamburg.de/biosaxs

Collaborative projects at EMBL beamlines:

- **Cytochrome/adrenodoxin:** M.Ubbink (Leiden Univresity)
- **BARF1, Flt3:** S. Savvides (Gent University)
- **Insulin, α-synuclein:** B.Vestergaard (Pharmaceutical Uni Copenhagen)
- **PEG-hemoglobin:** F.Ekström (FOM, Sweden), K.D.Vandegriff (Sangart, USA)
- **Myomesin:** N.Pinotsis (University Vienna), M.Wilmanns (EMBL, Hamburg)
- **Nuclear receptors:** D.Moras (CNRS Strasbourg)
- **S-layer proteins:** N. Fairweather, K. Brown (Imperial College, London)
- **Frataxin:** A.Pastore (NIMR MRC, London)
- **E2 core:** J. van den Elsen (University of Bath)
- **eRF1/3:** H.Song (Institute of Molecular and Cell Biology, Singapore)
- **Magnetic NPs:** L. Bronstein (Indiana University), E.Shtykova (ICRAS Moscow)
- **Chitinase:** W. Rypniewski (Poznan University),
- **Quantum dots:** L. Bronstein (Indiana University)
- **M1 protein:** E.Shtykova (ICRAS Moscow) L.Baratova (Moscow University)
- **Arcp1:** C.Koehler, D.Suck (EMBL, Heidelberg)
- **Gold NPs:** F.Stellacci (EPFL, Lausanne)