

Sneutrino or Stau as the Lightest Supersymmetric Particle in mSUGRA with R-parity Violation

Sebastian Grab


University of Bonn

VTI - Seminar
Bonn, May 20, 2008

Deutsche Telekom
Stiftung



Bonn-Cologne Graduate School
of Physics and Astronomy



in collaboration with

*B. C. Allanach, M. A. Bernhardt, S. P. Das, H. K. Dreiner,
C. H. Kom, P. Richardson and M. Trenkel*

- 1 Introduction
 - Supersymmetry
 - Minimal supergravity (mSUGRA)
- 2 Phenomenology of the stau LSP
 - Stau decays (2-body & 4-body)
 - Single slepton production
 - Sparticle pair production
- 3 Sneutrino as the LSP
 - Sneutrino LSP parameter space
 - Hadron collider phenomenology
- 4 Summary and Outlook

Supersymmetry (SUSY)

Why SUSY?

- Higgs mass is protected from quadratic divergencies.
- Unification of gauge couplings at $M_{GUT} = \mathcal{O}(10^{16})$ GeV.

What is SUSY?

$$\begin{aligned} Q |boson\rangle &= |fermion\rangle \\ \overline{Q} |fermion\rangle &= |boson\rangle \end{aligned}$$

- Q doesn't change gauge charges.
- Q doesn't change mass.

No SUSY partners observed so far.

$\Rightarrow$ SUSY must be broken.

Particle content of the MSSM

Minimal supersymmetric extension of the SM:




SM Particles	Superfields	spin 0	spin 1/2	spin 1
Quarks	Q_i $\bar{U}_i$ $\bar{D}_i$	$(\tilde{u}_{L_i}, \tilde{d}_{L_i})$ $\tilde{u}_{R_i}^c$ $\tilde{d}_{R_i}^c$	(u_{L_i}, d_{L_i}) $u_{R_i}^c$ $d_{R_i}^c$	
Leptons	L_i $\bar{E}_i$	$(\tilde{\nu}_i, \tilde{e}_{L_i})$ $\tilde{e}_{R_i}^c$	(ν_i, e_{L_i}) $e_{R_i}^c$	
Gauge Bosons	V_1 V_2 V_3		$\tilde{B}^0$ $\tilde{W}^\pm, \tilde{W}^0$ $\tilde{g}_a$	B^0 $W^\pm, W^0$ g_a
Higgs	H_u H_d	(H_u^+, H_u^0) (H_d^0, H_d^-)	$(\tilde{H}_u^+, \tilde{H}_u^0)$ $(\tilde{H}_d^0, \tilde{H}_d^-)$	

MSSM with R-parity violation (RPV)

General Superpotential of the Minimal Supersymmetric extension of the SM (MSSM):

$$W_{R_p} = (\mathbf{Y}_E)_{ij} L_i H_d \bar{E}_j + (\mathbf{Y}_D)_{ij} Q_i H_d \bar{D}_j + (\mathbf{Y}_U)_{ij} Q_i H_u \bar{U}_j + \mu H_d H_u ,$$

$$W_{\mathcal{R}_p} = \underbrace{\frac{1}{2} \lambda_{ijk} L_i L_j \bar{E}_k + \lambda'_{ijk} L_i Q_j \bar{D}_k}_{\Delta L \neq 0} + \underbrace{\frac{1}{2} \lambda''_{ijk} \bar{U}_i \bar{D}_j \bar{D}_k}_{\Delta B \neq 0} + \underbrace{\kappa_i L_i H_u}_{\Delta L \neq 0} .$$



The **lepton/baryon number violating** terms lead to **proton decay**.

It is sufficient to suppress $\Delta L \neq 0$ or $\Delta B \neq 0$ terms to keep proton stable.

[Dreiner, Luhn, Thormeier, Phys.Rev.D73:075007,2006]

Minimal supergravity (mSUGRA)

number of new parameters

- $\mathcal{O}(100)$ if R_p is conserved.
- $\mathcal{O}(200)$ if R_p is violated.


Assume simple boundary conditions at the scale $M_{GUT} = \mathcal{O}(10^{16})$ GeV.

mSUGRA parameter space

- M_0 : Universal soft breaking scalar mass.
- $M_{1/2}$: Universal gaugino soft breaking mass.
- A_0 : Universal trilinear scalar interaction.
- $\tan \beta$: Ratio of vevs. of the two Higgs doublets H_u, H_d .
- $\text{sgn } \mu$: Solution of EW symmetry breaking scalar potential.

Parameters at the scale $M_{EW} = \mathcal{O}(10^2)$ GeV are obtained by RGEs.
Programs: Softsusy, SPheno, Suspect, Isajet etc.


Running masses in mSUGRA



$\tilde{\chi}_1^0$ LSP versus $\tilde{\tau}$ LSP

[Allanach, Dedes, Dreiner, Phys.Rev.D69:115002,2004]

$$M_0 = A_0 = 0, \text{sgn}\mu = +1$$



- If R_p conserved:
Scenario is excluded.
(neutral LSP & $m_{h^0} > 114$ GeV).
- If R_p violated:
Most of the $\tilde{\tau}$ LSP region is allowed.

Add one parameter at M_{GUT} : $\mathbf{\Lambda} \in \{\lambda_{ijk}, \lambda'_{ijk}, \lambda''_{ijk}\}$.

$\Rightarrow$ **R-parity violating mSUGRA**

mSUGRA with R-parity violation (RPV)

We add one parameter at M_{GUT} : $\mathbf{\Lambda} \in \{\lambda_{ijk}, \lambda'_{ijk}, \lambda''_{ijk}\}$.

Differences to R_p conserving mSUGRA:

- Further R_p violating couplings are generated through RGEs at M_{EW} .
 $\Rightarrow$ 2-body and 4-body decays of $\tilde{\tau}$ LSP.
- Single sparticle production is possible.
 $\Rightarrow$ single slepton production.
- In principle, any SUSY particle could be the LSP.
 $\Rightarrow$ $\tilde{\tau}$ LSP region allowed.
- R_p violating RGEs change the SUSY mass spectrum at M_{EW} .
 $\Rightarrow$ $\tilde{\nu}$ LSP.
- Neutrino masses are generated.

What is the phenomenology of a $\tilde{\tau}$ LSP scenario at hadron colliders?

- $\tilde{\tau}$ LSP decays (2-body & 4-body).
[Dreiner, SG, Trenkel, work in progress]
- First example: Resonant single slepton production.
[Dreiner, SG, Trenkel, work in progress]
- Second example: Sparticle pair production.
[Allanach, Bernhardt, Dreiner, SG, Kom, Richardson, arXiv:0710.2034]

Typical mass ordering for $\tilde{\tau}$ LSP scenarios.

$$m_{\tilde{g}} > m_{\tilde{q}_2} > m_{\tilde{q}_1} > m_{\tilde{\chi}_2^+} > m_{\tilde{\chi}_1^+} \approx m_{\tilde{\ell}_2} > m_{\tilde{\chi}_1^0} \approx m_{\tilde{\mu}_1} \approx m_{\tilde{e}_1} > m_{\tilde{\tau}_1}$$

If $\Lambda \leq \mathcal{O}(10^{-3})$

- Sparticles are produced in pairs via gauge interactions, e.g. $\tilde{g}\tilde{g}$, $\tilde{q}\tilde{q}$.
- Sparticle undergo 2-body decays to the $\tilde{\tau}_1$ via gauge interactions.

$$\begin{aligned}\tilde{g} &\rightarrow \tilde{t}\bar{t} \\ &\hookrightarrow \tilde{\chi}_1^+ b \\ &\hookrightarrow \tilde{\nu}_\mu \mu^+ \\ &\hookrightarrow \tilde{\chi}_1^0 \nu_\mu \\ &\hookrightarrow \tilde{\tau}_1^- \tau^+\end{aligned}$$


If $\Lambda \geq \mathcal{O}(10^{-2})$

- Single sparticle production may dominate.
- RPV 2-body decays may alter the decay chains.




RPV decays of the $\tilde{\tau}$ LSP (naive picture)

- The dominant operator is: $L_3 L_j \bar{E}_k, L_i L_3 \bar{E}_k, L_i L_j \bar{E}_3$ or $L_3 Q_j \bar{D}_k$.
 $\Rightarrow$ 2-body decays.
- The dominant operator is: $L_{i \neq 3} L_{j \neq 3} \bar{E}_{k \neq 3}, L_{i \neq 3} Q_j \bar{D}_k$ or $\bar{U}_i \bar{U}_j \bar{D}_k$.
 $\Rightarrow$ 4-body decays.

For example $\lambda_{233} \neq 0$:




For example $\lambda'_{211} \neq 0$:



Dynamical generation of RPV couplings

Generation of λ_{i33} via λ'_{ijk}



$$16\pi^2 \frac{d\lambda_{i33}}{dt} = 3(\mathbf{Y}_E)_{33} \lambda'_{ijk} (\mathbf{Y}_D)_{jk} + \dots$$

Assume: $\mathbf{Y}_E = \text{diag} \Rightarrow$ e.g. if you break only L_e then $L_{\mu/\tau}$ will not be broken via RGEs.


Quark mixing: We know $\mathbf{V}_{CKM} = V_{uL}^+ V_{dL}$.

up-mixing: $\mathbf{Y}_U(M_Z) \times v_u = \mathbf{V}_{CKM}^+ \text{diag}(m_u, m_c, m_t) \mathbf{V}_{CKM},$
 $\mathbf{Y}_D(M_Z) \times v_d = \text{diag}(m_d, m_s, m_b)$

down-mixing: $\mathbf{Y}_U(M_Z) \times v_u = \text{diag}(m_u, m_c, m_t),$
 $\mathbf{Y}_D(M_Z) \times v_d = \mathbf{V}_{CKM} \text{diag}(m_d, m_s, m_b) \mathbf{V}_{CKM}^+.$


Running of RPV couplings: down-mixing

Dynamical generation of λ_{233}




For comparison:

Running of λ'_{2jk}




Running of RPV couplings: up-mixing

Dynamical generation of λ_{233}



For comparison:

Running of λ'_{2jk}



Decays of the $\tilde{\tau}$ LSP

Naive picture


- The dominant operator is: $L_3 L_j \bar{E}_k, L_i L_3 \bar{E}_k, L_i L_j \bar{E}_3$ or $L_3 Q_j \bar{D}_k$,
e.g. $\lambda_{233} \neq 0 \Rightarrow$ 2-body decays.
- The dominant operator is: $L_{i \neq 3} L_{j \neq 3} \bar{E}_{k \neq 3}, L_{i \neq 3} Q_j \bar{D}_k$ or $\bar{U}_i \bar{U}_j \bar{D}_k$,
e.g. $\lambda'_{2jk} \neq 0 \Rightarrow$ 4-body decays.

But: λ'_{2jk} will generate λ_{233} .

Question: 2-body or 4-body decay dominant?

2-body versus 4-body decays

BR(2-body decay), down-mixing



$$BR_2 = \frac{1}{1 + \Gamma_4/\Gamma_2}$$

with

$$\Gamma_2 \propto \lambda_{233}^2 m_{\tilde{\tau}_1}$$

$$\Gamma_4 \propto \lambda_{2jk}^{\prime 2} \frac{m_{\tilde{\tau}_1}^7}{m_{\tilde{\chi}}^2 m_f^4}$$

$$\Rightarrow \Gamma_4/\Gamma_2 \propto m_{\tilde{\tau}_1}^6$$

$m_{\tilde{\tau}_1}$ dependence on $\tan\beta$

$$\text{Recall: } \Gamma_4/\Gamma_2 \propto m_{\tilde{\tau}_1}^6$$

$$M_{\tilde{\tau}}^2 = \begin{pmatrix} m_{\tilde{\tau}_R}^2 & m_{\tilde{\tau}_{RL}} m_{\tau} \\ m_{\tilde{\tau}_{RL}} m_{\tau} & m_{\tilde{\tau}_L}^2 \end{pmatrix}$$

with

$$m_{\tilde{\tau}_R}^2 = m_{\tau}^2 + M_0^2 + 0.15M_{1/2}^2 - 0.23M_Z^2 \cos 2\beta - 2/3 X_{\tau}$$

$$m_{\tilde{\tau}_L}^2 = m_{\tau}^2 + M_0^2 + 0.52M_{1/2}^2 - 0.27M_Z^2 \cos 2\beta - 1/3 X_{\tau}$$

$$m_{\tilde{\tau}_{RL}} = A_{\tau} - \mu \tan\beta$$

where

$$X_{\tau} = 10^{-4} (1 + \tan^2\beta) (M_0^2 + 0.15M_{1/2}^2 + 0.33A_0^2)$$

[Drees, Martin, 1995]

$\tan\beta$ increase $\Rightarrow m_{\tilde{\tau}_1}$ decrease $\Rightarrow BR_2 = 1/(1 + \Gamma_4/\Gamma_2)$ increase.

$\tilde{\tau}$ LSP phenomenology


So far: $\tilde{\tau}$ LSP decays.

First Example: Resonant single slepton production.


Resonant single slepton production was only investigated for $\tilde{\chi}_1^0$ LSP!
e.g. [Allanach et al., Searching for R parity violation at Run II of the Tevatron, 1999]

Single slepton production via λ'_{ijk}

$\tilde{\mu}_L^-$ production via λ'_{211} :




$\tilde{\mu}_L^-$ decay via λ'_{211} :



suppressed if $\lambda' \leq \mathcal{O}(10^{-2})$

RPC $\tilde{\mu}_L^-$ decay:



$\tilde{\chi}_1^0$ LSP

- $\tilde{\chi}_1^0$ decays via λ'_{211} .
(3-body decay)

$\tilde{\tau}$ LSP

- $\tilde{\chi}_1^0 \rightarrow \tilde{\tau}_1 \tau$.
- $\tilde{\tau}_1$ decays via λ'_{211} or λ_{233} .

Promising signatures at hadron colliders

$\tilde{\chi}_1^0$ LSP:

$$\tilde{\mu}_L^- \rightarrow \mu^- \mu^- u \bar{d}.$$

$\tilde{\tau}$ LSP:

$$\tilde{\mu}_L^- \rightarrow \mu^- \mu^- \tau^+ \tau^- u \bar{d} \quad (4\text{-body decay}).$$

$$\tilde{\mu}_L^- \rightarrow \mu^- \mu^- \tau^+ \nu_\tau \quad (2\text{-body decay}).$$

⇒ Promising signature: Like-sign muon final states!


⇒ Low SM background: 5 events at LHC for 10 fb^{-1} after cuts!

[Dreiner, Richardson, Seymour, Phys.Rev.D63:055008,2001]

Cross sections at hadron colliders.

$$PP(\bar{P}) \rightarrow \tilde{\mu}_L^+ + X \text{ via } \lambda'_{211} = 0.01 \text{ at } M_{EW}.$$

[Dreiner,SG,Krämer,Trenkel, Phys.Rev.D75:035003]



Note: $\lambda'_{211} = 0.01$ at $M_{GUT} \Rightarrow \lambda'_{211} \approx 0.03$ at M_{EW} .

Numerical example for LHC

$M_0 = 0$ GeV, $M_{1/2} = 700$ GeV, $A_0 = 1150$ GeV, $\tan\beta = 26$, $\text{sgn}\mu = +1$.

- σ_{prod} : Cross section for $\tilde{\mu}_L$ production.
- $\sigma_{\lambda'}$: $\sigma_{prod} \times BR(\tilde{\mu}_L \rightarrow \mu^\pm \mu^\pm + X)$ & $\tilde{\tau}_1$ decay via λ' .
- σ_λ : $\sigma_{prod} \times BR(\tilde{\mu}_L \rightarrow \mu^\pm \mu^\pm + X)$ & $\tilde{\tau}_1$ decay via λ .

$m_{\tilde{\mu}_L} = 470$ GeV		σ_{prod} [fb]	up mixing		down mixing	
			$\sigma_{\lambda'}$ [fb]	σ_λ [fb]	$\sigma_{\lambda'}$ [fb]	σ_λ [fb]
$\lambda'_{211} = 1 \times 10^{-2}$	$\mu^- \mu^-$	476	1.02	99.2	—	100
	$\mu^+ \mu^+$	885	1.90	184	—	186
$\lambda'_{221} = 1 \times 10^{-2}$	$\mu^- \mu^-$	309	61.8	—	—	65.1
	$\mu^+ \mu^+$	105	21.1	—	—	22.2

- Final state might reveal quark mixing and $\tan\beta$.
- Ratio $(\#\mu^+\mu^+)/(\#\mu^-\mu^-)$ can reveal the indices j, k of λ'_{ijk} .

$\tilde{\tau}$ LSP phenomenology

First Example:
Resonant single slepton production.


Second example:
Sparticle pair production at the LHC.

Benchmark scenario BC1

BC1

- $M_0 = A_0 = 0$
- $\lambda_{121}(M_{GUT}) = 0.032$
- $\tan \beta = 13$
- $M_{1/2} = 400 \text{ GeV}$
- $\text{sgn}(\mu) = +1$.

[Allanach et. al., Phys.Rev.D75:035002,2007]



Branching ratios in benchmark scenario BC1

	mass [GeV]	channel	BR	channel	BR
$\tilde{\tau}_1$	148	$\mu^+ \bar{\nu}_e e^- \tau^-$	32 %	$e^+ \bar{\nu}_\mu e^- \tau^-$	32 %
		$\mu^- \nu_e e^+ \tau^-$	18 %	$e^- \nu_\mu e^+ \tau^-$	18 %
$\tilde{e}_R$	161	$e^- \nu_\mu$	50 %	$\mu^- \nu_e$	50 %
$\tilde{\mu}_R$	161	$\tilde{\tau}^+ \mu^- \tau^-$	51 %	$\tilde{\tau}^- \mu^- \tau^+$	49 %
$\tilde{\chi}_1^0$	162	$\tilde{\tau}_1^+ \tau^-$	50 %	$\tilde{\tau}_1^- \tau^+$	50 %
$\tilde{\nu}_\tau$	265	$\tilde{\chi}_1^0 \nu_\tau$	67 %	$W^+ \tilde{\tau}_1$	33 %
$\tilde{\nu}_e(\tilde{\nu}_\mu)$	266	$\tilde{\chi}_1^0 \nu_e(\nu_\mu)$	92 %	$\mu^+(e^+)e^-$	7.5 %
$\tilde{e}_L^-(\tilde{\mu}_L^-)$	280	$\tilde{\chi}_1^0 e^-(\mu^-)$	92 %	$e^- \bar{\nu}_\mu(\bar{\nu}_e)$	8.1 %
$\tilde{\tau}_2$	283	$\tilde{\chi}_1^0 \tau^-$	63 %	$Z^0 \tilde{\tau}_1^-$	18 %
		$h^0 \tilde{\tau}_1^-$	19 %		

Signal rates of benchmark scenario BC1

$$\sigma(\text{total sparticle pair production}) = 4.8 \text{ pb}$$

e^+ or μ^+	e^- or μ^-	τ^+	τ^-	$\cancel{p}_T$	event fraction
2	2	2	2	yes	35 %
3	2	2	2	yes	12 %
2	3	2	2	yes	8.3 %
3	3	2	2	yes	7.3 %
2	2	2	1	yes	4.7 %
2	2	3	2	yes	4.3 %
2	2	3	3	yes	1.4 %
4	3	2	2	yes	1.1 %


- Multi-lepton final states (≈ 8 leptons).
- Multi-tau final states (≈ 4 taus).
- 2-4 jets
- Missing p_T due to neutrinos from $\tilde{\tau}_1$ decay.

Benchmark scenario BC2

BC2

- $M_0 = A_0 = 0$
- $\lambda'_{311}(M_{GUT}) = 3.5 \cdot 10^{-7}$
- $\tan \beta = 13$
- $M_{1/2} = 400 \text{ GeV}$
- $\text{sgn}(\mu) = +1$.

[Allanach et. al., Phys.Rev.D75:035002,2007]



Branching ratios in benchmark scenario BC2

	mass [GeV]	channel	BR	channel	BR
$\tilde{\tau}_1$	148	$\bar{u}d$	100 %		
$\tilde{e}_R(\tilde{\mu}_R)$	161	$\tilde{\tau}_1^+ e^-(\mu^-)\tau^-$	51 %	$\tilde{\tau}_1^- e^-(\mu^-)\tau^+$	49 %
$\tilde{\chi}_1^0$	162	$\tilde{\tau}_1^+ \tau^-$	50 %	$\tilde{\tau}_1^- \tau^+$	50 %
$\tilde{\nu}_\tau$	265	$\tilde{\chi}_1^0 \nu_\tau$	67 %	$W^+ \tilde{\tau}_1$	33 %
$\tilde{\nu}_e(\tilde{\nu}_\mu)$	266	$\tilde{\chi}_1^0 \nu_e(\nu_\mu)$	100 %		
$\tilde{e}_L^-(\tilde{\mu}_L^-)$	280	$\tilde{\chi}_1^0 e^-(\mu^-)$	100 %		
$\tilde{\tau}_2$	283	$\tilde{\chi}_1^0 \tau^-$	63 %	$Z^0 \tilde{\tau}_1^-$	18 %
		$h^0 \tilde{\tau}_1^-$	15 %		

Signal rates of benchmark scenario BC2

$$\sigma(\text{sparticle pair production}) = 4.8 \text{ pb}$$

e^+ or μ^+	e^- or μ^-	τ^+	τ^-	$\cancel{p}_T$	event fraction
0	0	1	1	no	14 %
0	0	2	0	no	7.1 %
0	0	0	2	no	6.8 %
1	0	1	1	yes	6.5 %
0	0	1	1	yes	4.5 %
1	0	0	2	yes	3.3 %
1	0	2	0	yes	3.2 %
1	1	1	1	yes	2.4 %

- Like-sign τ events.
- 6-8 jets
- Not necessarily missing p_T signature.
- Detached vertex, i.e. $c \cdot \tau_{\tilde{\tau}_1} = 0.3 \text{ mm}$.

So far:
 $\tilde{\tau}$ LSP in mSUGRA.

Now:
Sneutrino as the lightest supersymmetric particle
in mSUGRA.

[Bernhardt, Dreiner, SG, Das, work in progress]

Effects of RPV

What will change due to **one additional RPV coupling** at the GUT scale?

The RGEs get additional contributions.

⇒ Additional RPV couplings at M_{EW} .

⇒ Sparticle masses can change at M_{EW} .

running sneutrino mass

$$16\pi^2 \frac{d(m_{\tilde{\nu}_i}^2)}{dt} = - \left(\frac{6}{5} g_1^2 |M_1|^2 + 6 g_2^2 |M_2|^2 + \frac{3}{5} g_1^2 S \right) \\ + 6 \lambda_{ijk}'^2 \left[m_{\tilde{\nu}_i}^2 + (\mathbf{m}_{\tilde{\mathbf{Q}}}^2)_{jj} + (\mathbf{m}_{\tilde{\mathbf{D}}}^2)_{kk} \right] + 6 (\mathbf{h}_{\mathbf{D}^k})_{ij}^2$$


$$\text{with } (\mathbf{h}_{\mathbf{D}^k})_{ij} = \lambda_{ijk}' \cdot A_0 \text{ at } M_{GUT},$$

$$S = f(\tilde{m}^2).$$

Note: Contribution of $(\mathbf{h}_{\mathbf{D}^k})_{ij}$ can dominate for negative A_0 .

What is the LSP?


A non-vanishing coupling $\lambda'(M_{GUT})$ leads to a new LSP candidate.
For SPS1a:



$\Rightarrow \tilde{\nu}_\mu$ LSP; also possible: $\tilde{\nu}_e$ & $\tilde{\nu}_\tau$ LSP.

$\tilde{\nu}_\mu$ LSP parameter space: M_0 - $M_{1/2}$ plane

$$\lambda'_{221}(M_{GUT}) = 0.1, A_0 = -500 \text{ GeV}, \tan \beta = 10, \mu > 0.$$



$\tilde{\nu}_\mu$ LSP parameter space: M_0 - $M_{1/2}$ plane


Different LSP regions because:

- $m_{\tilde{\tau}_R}^2 = M_0^2 + 0.15M_{1/2}^2 + \dots$
(right-handed stau couples only via U(1) charges.)
- $m_{\tilde{\nu}_\mu}^2 = M_0^2 + 0.52M_{1/2}^2 + \dots$
(left-handed sneutrino couples via U(1) & SU(2) charges.)
- $m_{\tilde{\chi}_1^0}^2 \simeq M_1^2 = 0.17M_{1/2}^2$.
($\tilde{\chi}_1^0$ is bino-like.)

[Ibanez, Lopez, Munoz, Nucl.Phys.B256,1985]

$\tilde{\nu}_\mu$ LSP parameter space: A_0 - $\tan\beta$ plane

$$\lambda'_{221}(M_{GUT}) = 0.1, M_0 = 50 \text{ GeV}, M_{1/2} = 500 \text{ GeV}, \mu > 0.$$



Different LSP regions because:

- $m_{\tilde{\tau}_R}^2 = m_\tau^2 + M_0^2 + 0.15M_{1/2}^2 - 0.23M_Z^2 \cos 2\beta - 2/3 X_\tau$
 with
 $X_\tau = 10^{-4}(1 + \tan^2 \beta)(M_0^2 + 0.15M_{1/2}^2 + 0.33A_0^2)$
- $m_{\tilde{\tau}_{RL}} = A_\tau - \mu \tan \beta$.
- $16\pi^2 \frac{dm_{\tilde{\nu}_i}^2}{dt} = 6(\mathbf{h}_{\mathbf{D}^k})_{ij}^2 + \dots$
 with
 $(\mathbf{h}_{\mathbf{D}^k})_{ij} = \lambda'_{ijk} \cdot A_0$ at M_{GUT} .

So far:

$\tilde{\nu}$ LSP in extended regions
of RPV mSUGRA parameter space.

Now: Phenomenology of a $\tilde{\nu}$ LSP at hadron colliders.

- Sparticle pair production.
- Single slepton production.

Sparticle pair production at LHC

Example: $\lambda'_{221}(M_{GUT}) = 0.1$, $\tan \beta = 10$, $\mu > 0$, $M_0 = 110$ GeV, $M_{1/2} = 440$ GeV, $A_0 = -500$ GeV.

$$\sigma_{LHC}(PP \rightarrow 2 \text{ Sparticles}) = 3.2 \text{ pb}.$$


Characteristic signatures

- Not necessarily missing p_T .
(22% of events).
- 4-7 non b-jets and 0-2 b-jets.
- High p_T muon plus high p_T jet.
(10% of events)

	mass	channel	BR
$\tilde{\nu}_\mu$	135	$\bar{s}d$	100 %
$\tilde{\mu}_L^-$	157	$\bar{c}d$	100 %
$\tilde{\chi}_1^0$	179	$\tilde{\nu}_\mu \bar{\nu}_\mu$	39 %
		$\tilde{\mu}_L^- \mu^+$	11 %
$\tilde{\tau}_1^-$	192	$\tilde{\chi}_1^0 \tau^-$	100 %
$\tilde{\nu}_e$	316	$\tilde{\chi}_1^0 \nu_e$	100 %
$\tilde{d}_R$	881	$\mu^- c$	44 %
		$\nu_\mu s$	44 %
		$\tilde{\chi}_1^0 d$	12 %
$\tilde{c}_L$	931	$\tilde{\chi}_1^+ s$	55 %
		$\tilde{\chi}_2^0 c$	27 %
		$\mu^+ d$	17 %

High- p_T muons

Muon p_T distributions from the decays $\tilde{d}_R \rightarrow \mu^- c$ and $\tilde{c}_L \rightarrow \mu^+ d$:



$$m_{\tilde{d}_R} = 881 \text{ GeV}, m_{\tilde{c}_L} = 931 \text{ GeV}.$$

- Squark mass can be reconstructed via high p_T muon and high p_T jet.
- $\tilde{\nu}_\tau$ LSP $\Rightarrow$ high p_T taus.


$\tilde{\nu}$ LSP phenomenology

So far: Sparticle pair production

Now: Resonant single slepton production.

Resonant single slepton production was only investigated for $\tilde{\chi}_1^0$ LSP!
e.g. [Allanach et al., Searching for R parity violation at Run II of the Tevatron, 1999]

Single $\tilde{\mu}_L$ and $\tilde{\nu}_\mu$ production via λ'_{221}



$$PP(\bar{P}) \rightarrow \tilde{\nu}_\mu + X$$

$$\hookrightarrow \bar{s}d$$



$$PP(\bar{P}) \rightarrow \tilde{\mu}_L^- + X$$

$$\hookrightarrow \bar{c}d$$

$$\hookrightarrow \tilde{\chi}_1^0 \mu^-$$

$$\hookrightarrow \tilde{\nu}_\mu \nu_\mu$$

$$\hookrightarrow \bar{s}d$$



Single $\tilde{\mu}_L$ and $\tilde{\nu}_\mu$ production via λ'_{221}

$$\lambda'_{221}(M_{GUT}) = 0.1, \quad M_0 = 170 \text{ GeV}, \quad M_{1/2} = 300 \text{ GeV}, \quad A_0 = -500 \text{ GeV}, \quad \tan \beta = 10, \quad \mu > 0.$$

$$\Rightarrow M_{\tilde{\mu}_L} = 140 \text{ GeV}, \quad M_{\tilde{\chi}_1^0} = 120 \text{ GeV}$$

Tevatron


$$\sigma(P\bar{P} \rightarrow \tilde{\mu}_L \rightarrow \tilde{\chi}_1^0 \mu) = 1.2 \text{ pb.}$$

LHC

$$\sigma(P\bar{P} \rightarrow \tilde{\mu}_L \rightarrow \tilde{\chi}_1^0 \mu) = 29 \text{ pb.}$$

$\Rightarrow \tilde{\nu}$ LSP scenarios might be found at the Tevatron!

Bottleneck: Small p_T of Muons.



Muon p_T from $P\bar{P} \rightarrow \tilde{\mu}_L \rightarrow \tilde{\chi}_1^0 \mu$
at the Tevatron

Summary and Outlook

Summary

- Including R-parity violation allows $\tilde{\tau}$ LSP in mSUGRA.
- Including R-parity violation changes RGEs in mSUGRA.
 $\Rightarrow$ 2-body versus 4-body $\tilde{\tau}$ decays.
 $\Rightarrow \tilde{\nu}$ LSP possible.
- Promising hadron collider signatures for $\tilde{\tau}$ LSP:
 detached vertices, multi-lepton final states, like-sign leptons.
- Promising hadron collider signatures for $\tilde{\nu}$ LSP:
 high- p_T muons, muons from single slepton production.
 $\Rightarrow$ Tevatron might find $\tilde{\nu}$ LSP scenarios.

Outlook


- Detailed analysis including background, detector simulations and data.
- Additional LSP candidates: $\tilde{e}_R$ with λ , $\tilde{t}_1$ with λ'' .

backup slides

2-body versus 4-body decay: A_0 -dependence

BR(2-body decay), down-mixing

$\lambda'_{211} = 0.01$, $M_0 = 0$ GeV, $M_{1/2} = 500$ GeV, $\text{sgn}\mu = +1$



$$BR_2 = \frac{1}{1 + \Gamma_4/\Gamma_2}$$

with

$$\Gamma_2 \propto \lambda_{233}^2 m_{\tilde{\tau}_1}$$


$$\Gamma_4 \propto \lambda_{2jk}'^2 \frac{m_{\tilde{\tau}_1}^7}{m_{\tilde{\chi}}^2 m_f^4}$$

$$\Rightarrow \Gamma_4/\Gamma_2 \propto m_{\tilde{\tau}_1}^6$$

2-body versus 4-body decay: $M_{1/2}$ -dependence

BR(2-body decay), down-mixing

$$\lambda'_{211} = 0.01, M_0 = 0 \text{ GeV}, A_0 = 600 \text{ GeV}, \text{sgn}\mu = +1$$



$$BR_2 = \frac{1}{1 + \Gamma_4/\Gamma_2}$$

with

$$\Gamma_2 \propto \lambda_{233}^2 m_{\tilde{\tau}_1}$$


$$\Gamma_4 \propto \lambda_{2jk}^{\prime 2} \frac{m_{\tilde{\tau}_1}^7}{m_{\tilde{\chi}}^2 m_f^4}$$

$$\Rightarrow \Gamma_4/\Gamma_2 \propto m_{\tilde{\tau}_1}^6$$

2-body versus 4-body decay: M_0 -dependence

BR(2-body decay), down-mixing

$\lambda'_{211} = 0.01$, $A_{600} = 0$ GeV, $M_{1/2} = 1400$ GeV, $\text{sgn}\mu = +1$



$$BR_2 = \frac{1}{1 + \Gamma_4/\Gamma_2}$$

with

$$\Gamma_2 \propto \lambda_{233}^2 m_{\tilde{\tau}_1}$$

$$\Gamma_4 \propto \lambda_{2jk}'^2 \frac{m_{\tilde{\tau}_1}^7}{m_{\tilde{\chi}}^2 m_{\tilde{f}}^4}$$

$$\Rightarrow \Gamma_4/\Gamma_2 \propto m_{\tilde{\tau}_1}^6$$

Possible Signatures

$\tilde{\tau}_1$ decay	$\tilde{\mu}_L$ production	$\tilde{\nu}_\mu$ production
via λ'_{2jk}	$\tau^+ \tau^- \quad \mu^- \mu^\pm \quad [\ell^+ \ell^-] \quad jj$	$\tau^+ \tau^- \quad \mu^\pm \quad [\ell^+ \ell^-] \quad \cancel{E}_T \quad jj$
	$\tau^+ \tau^- \quad \mu^- \quad [\ell^+ \ell^-] \quad \cancel{E}_T \quad jj$	$\tau^+ \tau^- \quad [\ell^+ \ell^-] \quad \cancel{E}_T \quad jj$
via λ_{233}	$\tau^+ \tau^- \quad \mu^- \quad [\ell^+ \ell^-] \quad \cancel{E}_T$	$\tau^+ \tau^- \quad [\ell^+ \ell^-] \quad \cancel{E}_T$
	$\tau^\pm \quad \mu^- \mu^\mp \quad [\ell^+ \ell^-] \quad \cancel{E}_T$	$\tau^\pm \quad \mu^\mp \quad [\ell^+ \ell^-] \quad \cancel{E}_T$

with $\ell = e, \mu$ if decays $\tilde{\chi}_1^0 \rightarrow \tilde{\ell}_R^\pm \ell^\mp$ and $\tilde{\ell}_R^- \rightarrow \ell^- \tau^\pm \tilde{\tau}_1^\mp$ allowed.

$$\begin{aligned}
 \bar{u}_j d_k &\xrightarrow{\lambda'} \tilde{\mu}_L^- \rightarrow \mu^- \tilde{\chi}_1^0, \\
 &\hookrightarrow \tau^+ \tilde{\tau}_1^- \\
 &\xrightarrow{\lambda'} \tau^- \mu^- u_j \bar{d}_k \\
 &\xrightarrow{\lambda} \nu_\tau \mu^-, \\
 &\hookrightarrow \tau^- \tilde{\tau}_1^+ \\
 &\xrightarrow{\lambda'} \tau^+ \mu^- u_j \bar{d}_k
 \end{aligned}$$

$\Rightarrow$ Multi-lepton final states,
e.g. four μ in final state.
 $\Rightarrow$ Like sign-muon events.

SM background for $\mu^\pm \mu^\pm$ events

4.9 ± 1.6 like-sign μ events after cuts
at the LHC for $10 fb^{-1}$. [Dreiner,
Richardson, Seymour, Phys.Rev.D63:055008]

RPV couplings leading to a sneutrino LSP

couplings λ'_{ijk} with upper bounds of $\mathcal{O}(0.1 - 1)$ at M_{EW}



coupling	LSP
λ'_{112}	$\tilde{\nu}_e$
λ'_{121}	$\tilde{\nu}_e$
λ'_{131}	$\tilde{\nu}_e$
λ'_{212}	$\tilde{\nu}_\mu$
λ'_{221}	$\tilde{\nu}_\mu$
λ'_{231}	$\tilde{\nu}_\mu$
λ'_{312}	$\tilde{\nu}_\tau$
λ'_{321}	$\tilde{\nu}_\tau$
λ'_{331}	$\tilde{\nu}_\tau$

and up-mixing.

Charm physics, e.g. $D_0 - \bar{D}_0$ mixing, will test couplings λ'_{i21} & λ'_{i12} .

Running of $(h_{D^k})_{ij}$


$$\lambda'_{ijk}(M_{GUT}) = 0.1, M_{1/2} = 500 \text{ GeV}$$



$$16\pi^2 \frac{d(\mathbf{h}_{D^k})_{ij}}{dt} = -(\mathbf{h}_{D^k})_{ij} \left(\frac{7}{15} g_1^2 + 3g_2^2 + \frac{16}{3} g_3^2 \right) + \lambda'_{ijk} \left(\frac{14}{15} M_1^2 + 6M_2^2 + \frac{32}{3} M_3^2 \right).$$


A_0 dependence

$$\lambda'_{221}(M_{GUT}) = 0.149, M_0 = 50 \text{ GeV}, \tan \beta = 10.$$




$\tilde{\nu}_\tau$ LSP parameter space

$$\lambda'_{331}(M_{GUT}) = 0.12, A_0 = -550 \text{ GeV}, \\ \tan \beta = 14, \mu > 0.$$




$$\lambda'_{331}(M_{GUT}) = 0.12, M_0 = 200 \text{ GeV}, \\ M_{1/2} = 270 \text{ GeV}, \mu > 0.$$



muon anomalous magnetic moment: $\delta a_\mu = a_\mu|_{exp} - a_\mu|_{SM} = 2.95 \times 10^{-9}$.
 $\Leftrightarrow 3.4\sigma$ deviation to SM prediction!


$\delta a_\mu|_{SUSY} = 2.95 \times 10^{-9}$ (red line), $\pm 1\sigma$, $\pm 2\sigma$.

Single $\tilde{\mu}_L$ and $\tilde{\nu}_\mu$ production via λ'_{221}




Problem: Large QCD background.


$W + \geq 2$ jets at the Tevatron




Dijet production at the Tevatron



p_T distributions in benchmark scenario BC1




p_T distribution of the τ from $\tilde{\tau}_1$ decays.




p_T distribution of the neutrinos.

- Taus with $p_T > 30$ GeV might be useful to identify the scenario.
- Missing p_T is less than in the R_p conserving MSSM.

p_T distributions in benchmark scenario BC2



p_T distribution of the τ from $\tilde{\chi}_1^0$ decays.




p_T distribution of the d-jets from $\tilde{\tau}_1$ decays.

- Tau identification is difficult but possible.
- Reconstruction of the $\tilde{\tau}_1$ mass is possible via the two jets.

Benchmark scenario BC3

BC3

- $M_0 = 100$ GeV
- $A_0 = -100$ GeV
- $\lambda'_{331}(M_{GUT}) = 0.122$
- $\tan \beta = 10$
- $M_{1/2} = 250$ GeV
- $\text{sgn}(\mu) = +1$.



Branching ratios in benchmark scenario BC3

	mass [GeV]	channel	BR	channel	BR
$\tilde{\nu}_\tau$	93	$\bar{b}d$	100 %		
$\tilde{\chi}_1^0$	97	$\tilde{\nu}_\tau \nu_\tau$	50 %	$\tilde{\nu}_\tau \bar{\nu}_\tau$	50 %
$\tilde{\tau}_1^-$	105	$\nu_\tau \bar{b}d\tau^-$	37 %	$\bar{\nu}_\tau \bar{b}d\tau^-$	37 %
		$\tilde{\chi}_1^0 \tau^-$	26 %		
$\tilde{e}_R^-(\tilde{\mu}_R^-)$	146	$\tilde{\chi}_1^0 e^-(\mu^-)$	100 %		
$\tilde{\tau}_2^-$	159	$\tilde{\chi}_1^0 \tau^-$	100 %		
$\tilde{\chi}_2^0$	181	$\tilde{\nu}_\tau \nu_\tau$	27 %	$\tilde{\nu}_\tau \bar{\nu}_\tau$	27 %
		$\tilde{\tau}_1^+ \tau^-$	22 %	$\tilde{\tau}_1^- \tau^+$	22 %
$\tilde{\chi}_1^-$	181	$\tilde{\nu}_\tau \tau^-$	63 %	$\tilde{\tau}_1^- \nu_\tau$	35 %
$\tilde{\nu}_e(\tilde{\nu}_\mu)$	189	$\tilde{\chi}_1^0 \nu_e(\nu_\mu)$	85 %	$\tilde{\chi}_1^+ e^-(\mu^-)$	11 %
$\tilde{e}_L^-(\tilde{\mu}_L^-)$	206	$\tilde{\chi}_1^0 e^-(\mu^-)$	48 %	$\tilde{\chi}_1^- \bar{\nu}_e(\bar{\nu}_\mu)$	33 %
		$\tilde{\chi}_2^0 e^-(\mu^-)$	19 %		


Signal rates of benchmark scenario BC3

$$\sigma(\text{sparticle pair production}) = 4.7 \cdot 10^4 \text{ fb}$$


e^+ or μ^+	e^- or μ^-	τ^+	τ^-	$\cancel{p}_T$	event fraction
0	0	0	0	yes	27 %
0	0	1	0	yes	19 %
0	0	0	1	yes	16 %
0	0	1	1	yes	14 %
0	0	1	1	no	4.4 %
0	0	2	1	yes	4.0 %
0	0	1	2	yes	3.0 %
1	0	0	1	yes	1.9 %

- Most difficult scenario to trigger, although light spectrum.
- 4.7 million sparticle events at the LHC with $\int \mathcal{L} = 100 \text{ fb}^{-1}$.
- b-tagging should be possible.

p_T distributions in benchmark scenario BC3



p_T distribution of the b-jets from $\tilde{\nu}_\tau$ decays.




p_T distribution of the τ from $\tilde{\tau}_1$ decays.

- b-tagging should be possible.
- Most of the taus from $\tilde{\tau}_1$ decays are invisible ($p_T \leq 30$ GeV).

Benchmark scenario BC4

BC4

- no-scale mSUGRA
- $\lambda''_{212}(M_{GUT}) = 0.5$
- $\tan \beta = 30$
- $M_{1/2} = 600$ GeV
- $\text{sgn}(\mu) = +1$.



Branching ratios in benchmark scenario BC4

	mass [GeV]	channel	BR	channel	BR
$\tilde{\tau}_1$	169	$c d s \tau^-$	79 %	$\bar{c} \bar{d} \bar{s} \tau^-$	21 %
$\tilde{e}_R(\tilde{\mu}_R)$	236	$\tilde{\tau}_1^+ e^- (\mu^-) \tau^-$	58 %	$\tilde{\tau}_1^- e^- (\mu^-) \tau^+$	42 %
$\tilde{\chi}_1^0$	249	$\tilde{\tau}_1^+ \tau^-$	47 %	$\tilde{\tau}_1^- \tau^+$	47 %
$\tilde{\nu}_\tau$	393	$W^+ \tilde{\tau}_1$	89 %	$\tilde{\chi}_1^0 \nu_\tau$	12 %
$\tilde{\nu}_e(\tilde{\nu}_\mu)$	402	$\tilde{\chi}_1^0 \nu_e (\nu_\mu)$	100 %		
$\tilde{e}_L^-(\tilde{\mu}_L^-)$	413	$\tilde{\chi}_1^0 e^- (\mu^-)$	100 %		
$\tilde{\tau}_2$	417	$Z^0 \tilde{\tau}_1^-$	48 %	$h^0 \tilde{\tau}_1^-$	38 %
		$\tilde{\chi}_1^0 \tau^-$	15 %		
$\tilde{d}_R(\tilde{s}_R)$	897	$\bar{c} \bar{s} (\bar{d})$	99 %	$\tilde{\chi}_1^0 d (s)$	1.2 %
$\tilde{c}_R$	906	$\bar{s} \bar{d}$	95 %	$\tilde{\chi}_1^0 c$	4.7 %


Signal rates of benchmark scenario BC4

$$\sigma(\text{sparticle pair production}) = 7.1 \cdot 10^2 \text{ fb}$$


e^+ or μ^+	e^- or μ^-	τ^+	τ^-	$\cancel{p}_T$	event fraction
0	0	1	1	no	23 %
0	0	0	0	no	18 %
0	0	2	2	no	8.0 %
1	0	2	2	yes	5.6 %
0	0	2	1	yes	4.1 %
1	1	2	2	no	3.7 %
1	0	1	1	yes	3.6 %
0	1	2	2	yes	3.2 %

- Many jets in final state (6-8 jets).
- Very little missing p_T .
- Heavy spectrum.
- First two generations of $\tilde{q}_R$ undergo RPV decays.

p_T distributions in benchmark scenario BC4



p_T distribution of the τ from $\tilde{t}_1$ decay.




p_T distribution of the d-jets from $\tilde{t}_1$ decay.


- Triggering to taus should be possible.

Dynamical generation of RPV couplings

Dynamical generation of λ_{ikk}



Dynamical generation of κ_i




$$16\pi^2 \frac{d}{dt} \lambda_{ikk} = (Y_E)_{kk} [3\lambda'_{iaq} (Y_D)_{aq}^* + \lambda_{ill} (Y_E)_{ll}^*]$$

$$16\pi^2 \frac{d}{dt} \lambda'_{ijk} = \lambda'_{ijl} 2(Y_D^\dagger Y_D)_{kl} + \lambda'_{ilk} [(Y_D Y_D^\dagger)_{lj} + (Y_U Y_U^\dagger)_{lj}] \\ + 3\lambda'_{iaq} (Y_D)_{aq}^* (Y_D)_{jk} + \lambda_{iaa} (Y_E)_{aa}^* (Y_D)_{jk}$$


$$16\pi^2 \frac{d}{dt} \kappa_i = \mu [3\lambda'_{iaq} (Y_D)_{aq}^* + \lambda_{ill} (Y_E)_{ll}^*].$$

Breaking of one lepton number does not break the two other lepton numbers.

More p_T distributions in benchmark scenario BC1




p_T distribution of the $\tilde{\tau}_1$.




p_T distribution of τ coming from $\tilde{\chi}_m^0$ decays.

More p_T distributions in benchmark scenario BC1




p_T distribution of the ℓ^+ coming from $\tilde{\tau}_1$ decays.




p_T distribution of the ℓ^- coming from $\tilde{\tau}_1$ decays.

More p_T distributions in benchmark scenario BC2




p_T distribution of the $\tilde{t}_1$.




p_T distribution of the u-jets from $\tilde{t}_1$ decays.

More p_T distributions in benchmark scenario BC3




p_T distribution of the d-jets from $\tilde{\nu}_\tau$ decays.




p_T distribution of the neutrinos.

More p_T distributions in benchmark scenario BC3




p_T distribution of the d-jets from $\tilde{\tau}_1$ decays.




p_T distribution of the s-jets from $\tilde{\tau}_1$ decays.

More p_T distributions in benchmark scenario BC4




p_T distribution of the c-jets from $\tilde{\tau}_1$ decay.




p_T distribution of the s-jets from $\tilde{\tau}_1$ decay.

More p_T distributions in benchmark scenario BC4




p_T distribution of the $\tilde{\tau}_1$.




p_T distribution of the neutrinos.

More p_T distributions in benchmark scenario BC4



p_T distribution of τ from $\tilde{\chi}_m^0$ decays.



p_T distribution of the d-jets from $\tilde{\chi}_R$ decay.