

Belle and Belle II (Schwerpunkt deutsche Gruppen)

Jens Sören Lange (Justus-Liebig-Universität Gießen)

KET Meeting, Bad Honnef, 15.11.2014





Foto 04.11.2014



Bundesministerium
für Bildung
und Forschung




Nano-Beam Scheme

Belle → Belle II


Luminosity x 40

$$\mathcal{L} \leq 0.8 \times 10^{36} \text{ cm}^{-2} \text{ s}^{-1}$$

KEKB (*without crab*)



SuperKEKB




originally proposed for SuperB
by P. Raimondi (INFN)

graphics E. Paoloni (Pisa)

Belle II Physik (Beispiele)

- Präzisionsvermessung des Unitaritätsdreiecks (CKM)




Ziel:
 $\gamma (1^\circ)$
 $V_{ub} (5\%)$
 $V_{cb} (2\%)$

- Seltene B Zerfälle

- $B^+ \rightarrow \tau^+ \underline{\nu}$
- $b \rightarrow s \underline{\nu} \underline{\nu}$

Vorteil: BB Paar-Rekonstruktion
 (zweites B Meson, andere Seite)

- XYZ Zustände (Charmonium, Bottomonium)
- A_{FB} in $e^+e^- \rightarrow \mu^+\mu^-$
- Leptonenzahlverletzung in τ Zerfällen
- Suche nach dunklem Photon
- ...



Sensitivität
 CKM Kopplungen

$$\Lambda_{NP} \leq 10^4 \text{ TeV}$$

rechtshändige Kopplungen

$$\Lambda_{NP} \leq 10^6 \text{ TeV}$$

Sensitivität
 Kontaktwechselwirkungen

$$\Lambda_{NP} \leq 33 \text{ TeV}$$


Belle II Kollaboration (Status 09.11.2014)

679 Mitglieder, 99 Institute, 23 Länder



Belle II Mitglieder

(inkl Studenten und Techniker)



Status Belle and Belle II (deutsche Gruppen)


- Belle 62 Mitglieder, 7 Institute (Status 09.11.2014)
Belle II 96 Mitglieder, 11 Institute
mit neuen deutschen Mitgliedern $\rightarrow \geq 100$
- BMBF Anträge
neue Institute: Univ. Mainz (Sfienti), Univ. Hamburg (Hagner)
neue Gruppen: KIT Karlsruhe IPE (Peric)
- DPG 2014 Mainz
29 Beiträge (Belle I und II)
- Vorträge auf Konferenzen/Workshops in 2014
19 (Belle I und II)
- erfolgreicher EU RISE Antrag – JENNIFER (Belle II und T2K)
Japan and Europe Network for Neutrino and Intensity Frontier Experimental Research
5 deutsche Institute
Reisekosten: 93 Personenmonate am KEK (über 4 Jahre)


Deutsche Beiträge zu Belle II

- PXD **alle**
- spare-PXD
nächste Förderperiode (2015–18)
- L1 z-Vertex-Neurotrigger **TUM, MPI**
- Computing
 - Softwareentwicklung
 - PXD **alle**
 - non-PXD **KIT**
 - Tracking Algorithmen
DESY, KIT, LMU, TUM, MPI (Vertexing), **Bonn, Giessen** (Online)
 - GRID Ressourcen **DESY, KIT, MPI**
 - Alignment **DESY**

Pixel Vertex Detector for Belle II @ SuperKEKB

Precise Measurement
of particle origin









Bundesministerium
für Bildung
und Forschung


Belle II DEPFET Pixel Detector
z Vertexauflösung
 $50 \mu\text{m}$ (Belle) $\rightarrow 25 \mu\text{m}$ (Belle II)

PXD+SVD DESY team test (01/2014)

$\geq 2 \times 10^7$ events recorded




IR Mechanics, Remote Vacuum Connection
(DESY, MPI)



CO₂ Cooling PXD+SVD (MPI, DESY)





Thermal Mockup (DESY, Hamburg)



Belle und Belle II Daten in Deutschland

- Alle Belle I Daten ($\simeq 1 \text{ ab}^{-1}$, μDST) nach DESY kopiert
Zugang aller Gruppen zur DESY National Analysis Facility
- Belle II MC Produktion 10/2014
 - 4.7M Jobs
 - entspricht 2.9 ab^{-1}
 - Peak vergleichbar mit LHCb (26k Jobs)
 - DESY **28.5%**, KIT 3.7%, MPI 2.8%



Belle Publikationen mit starker deutscher Beteiligung (Status 09.11.2014) (Belle I Analysen nicht BMBF gefördert)

- Measurement of branching fractions and CP violation parameters in $B \rightarrow \omega K$ decays with first evidence of CP violation in $B^0 \rightarrow \omega K S^0$,

V. Chobanova, J. Dalseno, C. Kiesling, et al. (Belle Collaboration),

[arXiv:1311.6666\[hep-ex\]](https://arxiv.org/abs/1311.6666), publiziert in Phys. Rev D90(2014)012002,

Suche nach CP Verletzung in einem Pinguin-dominierten Zerfall, sensitiv auf den Winkel ϕ_1 im Unitaritätsdreieck

$$B \rightarrow \omega K$$

- Study of $B^0 \rightarrow \rho^0 \rho^0$ decays, implications for the CKM angle ϕ_2 and search for other B^0 decay modes with a four-pion final state,

P. Vanhoefer, J. Dalseno, C. Kiesling, et al. (Belle Collaboration),

[arXiv:1212.4015\[hep-ex\]](https://arxiv.org/abs/1212.4015), veröffentlicht in Phys. Rev. D89(2013)072008,

Constraint für den Winkel $\phi_2 = 84.9 \pm 12.9$ grad im Unitaritätsdreieck

$$B \rightarrow \rho^0 \rho^0$$

- Angular analysis of $B^0 \rightarrow \phi K^*$ decays and search for CP violation at Belle

M. Prim, et al. (Belle Collaboration)

[arXiv:1308.1830\[hep-ex\]](https://arxiv.org/abs/1308.1830), veröffentlicht in Phys. Rev. D 88(2013)072004

$b \rightarrow s \bar{s} s$ Pinguin Zerfall, in welchem eine überraschend geringe longitudinale Polarisation gemessen wurde (Hinweis auf Verletzung der Faktorisierung).

$$B \rightarrow J/\psi K^*$$

- Search for $B \rightarrow h^{(*)} \nu \bar{\nu}$ with the full Belle data sample,

O. Lutz, S. Neubauer, M. Heck, T. Kuhr, et al. (Belle collaboration),

[arXiv:1303.3719\[hep-ex\]](https://arxiv.org/abs/1303.3719), veröffentlicht in Phys. Rev. D87(2013)111103,

Seltener Zerfall, welcher sensitiv auf rechtshändige Kopplungen jenseits des Standardmodells ist.

$$B \rightarrow h^{(*)} \nu \bar{\nu}$$

- Measurement of the inclusive semileptonic branching fraction $B(B_s^0 \rightarrow X l \nu)$ at Belle,

C. Oswald, P. Urquijo, J. Dingfelder, et al. (Belle collaboration),

[arXiv:1212.6400\[hep-ex\]](https://arxiv.org/abs/1212.6400), veröffentlicht in Phys. Rev. D87(2013)072008,

Test von SU(3) Symmetrie

$$B_s \rightarrow X l \nu$$

- First observation of CP violation and improved measurement of the branching fraction and polarization of $B^0 \rightarrow D^{*+} D^{*-}$ decays,

B. Kronenbitter, et al. (Belle collaboration),

[arXiv:1207.5611\[hep-ex\]](https://arxiv.org/abs/1207.5611), veröffentlicht in Phys. Rev. D86(2012)071103(R),

Messung eines CP-verletzenden Zerfalls mit Charm-Mesonen

$$B \rightarrow D^{*+} D^{*-}$$

- Measurements of Branching Fractions and Time-dependent CP Violating Asymmetries in $B^0 \rightarrow D^{(*)+} D^-$ Decays,

M. Rohrken, et al. (Belle collaboration),

[arXiv:1203.6647\[hep-ex\]](https://arxiv.org/abs/1203.6647), veröffentlicht in Phys. Rev. D85(2012)091106(R),

Messung eines CP-verletzenden Zerfalls mit Charm-Mesonen

$$B \rightarrow D^{(*)+} D^-$$

Belle II Organization

Executive Board

Chair : H. Aihara

aihara@phys.s.u-tokyo.ac.jp

D.M.Asner, T.Aziz, A.Bozek, P.Chang,
F.Forti, T.Iijima, P.Krizan, S.Lange,
P.Podesta, M.Roney, C.Schwanda,
M.Sevior, E.Won, C.Z.Yuan, K.Akai

Spokesperson : Thomas E. Browder

teb@phys.hawaii.edu

Project Manager : Yoshihide Sakai

Yoshihide.Sakai@kek.jp

Financial Board

Chair : Y.Sakai

Yoshihide.Sakai@kek.jp

W.Abdullah, H.Aihara, D.Asner, R.Ayad, T.Aziz,
A.Bozek, T.Browder, P.Chang, Z.Dolezal,
G. Finocchiaro, P.Krizan, C.Lacasta, M.I.Martinez,
H.G.Moser, C.Niebuhr, P.Pakhlov, A.Rekalo,
M.Konie, C.Schwanda, M.Sevior, C.P. Shen,
U.Tippawan, T.Tran, N.Wermes, E.Won, M.Zeyrek

Institutional Board

Chair : Z.Dolezal

dolezal@ipnp.troja.mff.cuni.cz

Speakers Committee

Chair : A.Schwartz

alan.j.schwartz@uc.edu

T.Iijima, I. Peruzzi, Y.Sakai, C.Schwanda

Physics

Coordinator

: P.Urquijo

purquijo@unimelb.edu.au

Semileptonic & Missing Energy
: A. Zupanc, G. De Nardo

Radiative & Electroweak Penguin
: A. Ishikawa, J. Yamaoka

T-Dep. CP Violation
: T. Higuchi, K. Sumisawa

Hadronic B Decay & DCPV
: J. Libby, P. Goldenzweig

Y(nS) : R. Mizuk, T. Pedlar

Charm & Carmonium
: R. Briere, G. Casarosa

Tau & Low Multiplicity : T. Ferber,
: K. Hayasaka, H. Nakazawa

Technical

Coordinator

: Y.Ushiroda

ushiroda@post.kek.jp

Integration Leaders : I. Adachi (Outer)
S. Tanaka (Inner)

PXD : H.G. Moser
C. Kiesling

SVD : K. Hara
C. Imler

CDC : S. Uno
TOP : J. Fast

ARICH : S. Nishida
S. Korpar

ECL : A. Kuzmin

EKLM : P. Pakhlov

BKLM : L. Piiionen

TRG : Y. Iwasaki
DAQ : R. Itoh
IR : H. Nakayama
STR : J. Haba
BKG : S. Vahsen

Liaisons :

S. Tanaka (PXD)
I. Adachi (BPID)
I. Nakamura (ECL)
K. Sumisawa (BKLM/EKLM)

Software

Coordinator

: T.Kuhr

Thomas.Kuhr@kit.edu

Simulation : D. Kim
Tracking : M. Heck, E. Paoloni
Alignment : S.Yashchenko
Database : M. Bracko
: L. Wood

Computing

Coordinator

: T.Hara


takanori.hara@kek.jp

Distributed Computing : M. Sevier
Data Transfer / Network : M. Schram
MC prod. manager : H. Miyake
Monitor : K. Hayasaka

Installation of 100 new LER Dipole Magnets



field measurement




move into tunnel



carry on an air-pallet

Installation of 100 new LER bending magnets done



install done

~3 magnets per day

Install over HER magnets



SuperKEKB Status, 7th BPAC, Mar. 11, 2013, K. Akai

B-KLM Installation



2013

Completed on November 16th

~2 months delay




During installation, modules have been checked, found to be healthy.

The 1st New Detector Installed !!

CDC wire stringing finished (01/2014)

51456 wires



FWD side: ground cabling finished (11/2014)

white: axial super layers

green: stereo super layers

TOP Assembly of 1st Module






Scene at the Fuji Hall clean room just before Halloween 2014

ECL cosmic data taking w/ ~ 4000 crystals

31.10.2014, w/ new electronics:

- shaping time 1 μs (before) $\rightarrow$ 0.5 μs (Belle II)
- waveform digitization 2 MSamples/s




Kürzungen des KEK Operations-Budget

betrifft z.B. Commissioning des Beschleunigers

Jahr	Betrag benötigt und beantragt: 50.4 MEuro
2014	10.9 MEuro
2014	13.7 MEuro (Forderung Rückzahlung von 2013)
2015	16.9 MEuro


→ Verzögerung des Schedules

SuperKEKB Luminositätsextrapolation (post-KEK-Budgetkürzung FY2014)



Start der Physikdatennahme verzögert sich 10/2016 → **10/2017**
Impliziert Planänderung: alle TOP Module werden verfügbar sein
Aus- und Wiedereinbau von PXD, SVD, CDC nicht mehr notwendig
(keine Datennahme mehr mit $\frac{1}{2}$ TOP)

Belle II Schedule



Zusammenfassung

- Deutschland ist zweitstärkste Gruppe in Belle II (≥ 100 Mitglieder)
- signifikante Beiträge (PXD, Computing)
- KEK Budgetkürzungen erfordern Verschiebung des Schedules
→ erste Physikdatennahme geplant für 10/2017

Vielen Dank.

CLICK → [\[Belle deutsche Sektion WWW Homepage\]](#) ← CLICK

Backup

Spokesman: C. Kiesling (MPI), Technical coordinator: H.-G. Moser (MPI)

Group

IHEP Beijing, China (Z.A. Liu)
 Charles University, Prague, Czech Rep. (Z. Dolezal)
 DESY Hamburg (C. Niebuhr)
 University of Bonn (N. Wermes, H. Krüger)
 University of Hamburg (C. Hagner)
 University of Heidelberg (P. Fischer)
 University of Giessen (W. Kühn, S. Lange)
 University of Göttingen (A. Frey)
 University of Karlsruhe (T. Müller, M. Feindt, I. Peric)
 University of Mainz (C. Sfienti)
 Ludw.-Max.-University, Munich (Th. Kuhr)
 MPI for Physics, Munich (C. Kiesling, H.G.Moser)
 Semiconductor Lab (HLL) (J. Ninkovic)
 Technical University, Munich (S. Paul, A.Knoll)
 Struct. Biol.Research Center, KEK (S. Wakatsuki)
 IFJ PAN, Krakow, Poland (A. Bozek)
 University of Barcelona, Spain (A. Dieguez)
 CNM, Barcelona, Spain (E. Cabruja)
 IFCA Santander, Spain (I. Vila)
 IFIC, Valencia, Spain (C. Lacasta)
 University of Tabuk (R. Ayad)


Pers. Responsibilities

10	ATCA Boards
7	Beam test analysis
14	RVC, CO2, th. mock-up
15	DHP ASIC, DatCon ROIs
3	mech. mock-up
2	Slow Control
5	DAQ hardware
7	Beam tests, analysis
16	S/W framew., DCD, SW
4	Slow Control
3	P/S, Kapton, tracking
17	Sensors, mech., CO2
5	Sensors
7	DHH, Slow Control
4	Sensor testing
11	FTB for DatCon
3	ASIC design
2	(flip chipping)
3	environm. sensors
4	SMD + sensor rework
3	Comic ray test setup

Belle II VXD

4-layer SVD

$R = 3.8, 8.0, 11.5, 14 \text{ cm}$




2-layer PXD

$R = 1.4, 2.2 \text{ cm}$


beampipe $R = 1.0 \text{ cm}$

DEPFET Production




Phase I – before metal

- Implantations
- Polysilicon
- Dielectric depositions



Phase II - Alu

- Metal 1
- isolation
- Metal 2



Phase III – thinning and Cu

- Handle wafer removal
- Dielectric deposition
- Metal 3
- Passivation




Phase II:EMCM production(s) for Alu/Cu

z-Vertex-Neurotrigger

CDC Tracking:

■ 15000 wires $\Rightarrow$ 2336 track-segments (TS)




Which information is available for triggering?


- a) Identification numbers of active track-segments (TS-IDs)
 $\Rightarrow$ Position of priority wire
- b) Drift times of priority wires

z-Vertex-Neurotrigger

Algorithmus: Multi-Layer Perzeptron



Resolution (MC)



S. Skambraks et al., arXiv:1406.3319[physics.ins-det]
S. Neuhaus et al., arXiv:1410.1395[physics.ins-det]

Hardware platform
UT3 board (Xilinx Virtex-6 HXT)






2 Phase Accumulator Controlled Loop




“2PACL”, proposed by B. Verlaat (NIKHEF)



(C. Kiesling)

Two Dry Volumes: “cold” and “warm”



“warm” volumes between QCS and CDC inner wall, up to CDC end wall, both in FWD and BWD directions.

flushed by nitrogen mandatory (potential hazard by oil from dry air supplier)



VXD Cooling Requirements & Specs




- Minimize material in acceptance region
- SVD requirements:
 - coolant temperature for APV25 @ -20°C for SNR improvement
 - total dissipated power $\sim 700\text{ W}$
- PXD requirements:
 - sensor temp. $< 25^{\circ}\text{C}$, ASICs $< 50^{\circ}\text{C}$: cooling block @ $< -20^{\circ}\text{C}$
 - dissipated power $\sim 360\text{ W}$
- Cooling power $\sim 1.6\text{ kW}$ at VXD (incl. $\sim 540\text{ W}$ parasitic heat influx)
 - have to add heat loss through 15 m of transfer lines $\rightarrow \sim 2\text{-}3\text{ kW}$
- Dry Volume (preferably nitrogen):
 - dew point inside VXD volume: $\sim -30^{\circ}\text{C}$
- Outside of VXD: room temperature ($\sim 23^{\circ}\text{C}$) at the CDC
 - may need a thermal enclosure towards CDC, well isolated

— Cold CO₂ line
 — Cold R404a line
 — Warm service line
 (Cold lines require 32mm insulation)

System A

System B



IBBelle: The “Twin” at CERN

Installation at CERN (ATLAS Experiment)



Accum.
rack



SuperKEKB accelerator Japan

Mt. Tsukuba

SuperKEKB ring (HER+LER)

Belle II detector

Linac

KEK - Tsukuba

en@fnas@uni-bonn.de





Replace short dipoles with longer ones (LER)




Redesign the lattices of HER & LER to squeeze the emittance

TiN-coated beam pipe with antechambers



Colliding bunches

New superconducting / permanent final focusing quads near the IP



To get x40 higher interaction rate

Luminosity increase

1. beam current

1.64/1.19 A $\rightarrow$ 3.6/2.6 A for e⁺ (e⁻) beam
 $\rightarrow$ factor 2 increase in luminosity

2. beta function

β_y^*

5.9 mm $\rightarrow$ 0.27 mm

$\sigma_y \rightarrow$ 59 nm

"nanobeam"

$\rightarrow$ factor 20 increase in luminosity

$\rightarrow$ total factor 40 increase in luminosity


$$\sigma_y(z) \propto \sqrt{\varepsilon_{inj} \cdot \beta_y(z)}$$

$$\beta_y(z) = \beta_y^* \left(1 + \frac{(z - Z_0)^2}{\beta_y^{*2}} \right)$$

Z_0 is position
of minimum
beta function
(„waiste“)


IP Profile for Belle and Belle II („nanobeam“)

$100\mu\text{m}(H) \times 2\mu\text{m}(V) \rightarrow 10\mu\text{m}(H) \times 59\text{nm}(V)$





Belle Rotation 03/2013


for larger crossing angle $22 \text{ mrad} \rightarrow 83 \text{ mrad}$ (nanobeam)



18.6 cm @ end of platform (accuracy 0.5 mm)

Background increase x factor 10–20











Hardware Resources for Belle II

Yearly integrated luminosity




Site or Tier 1 fraction	Germany	This can be a site name as given on the Sites sheet or a number
Data fraction		If this is empty the data fraction will be equal to the number given on the Sites sheet or 0 for numerical Tier 1
Used Tier 1 fraction	0.12	The fraction of MC production and user analysis done at the site
Used data fraction	0.40	The fraction of mDST data stored at the site

Regional Center resources:


Year	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
Disk (data) [PB]	0.00	0.00	0.00	0.00	0.02	0.08	0.66	2.03	3.73	5.46	7.28
Disk (MC) [PB]	0.00	0.00	0.00	0.00	0.04	0.15	0.67	1.28	2.05	2.83	3.64
Disk (analysis) [PB]	0.00	0.00	0.00	0.00	0.00	0.01	0.10	0.30	0.56	0.82	1.09
Disk (challenges) [PB]	0.00	0.12	0.24	0.48	0.60	0.30	0.00	0.00	0.00	0.00	0.00
Disk [PB]	0.00	0.12	0.24	0.48	0.67	0.54	1.42	3.61	6.34	9.11	12.01
CPU (MC) [kHepSPEC]	0.00	0.00	0.00	0.00	4.33	14.36	32.02	38.28	47.55	56.71	68.23
CPU (an.) [kHepSPEC]	0.00	0.00	0.00	0.00	0.15	0.55	4.34	13.42	24.69	36.15	48.16
CPU (ch.) [kHepSPEC]	0.00	7.55	15.99	15.99	13.32	24.27	0.00	0.00	0.00	0.00	0.00
CPU [kHepSPEC]	0.00	7.55	15.99	15.99	17.81	39.18	36.37	51.70	72.23	92.86	116.39
CPU [# of cores 14.72HS/core]	0.00	512.95	1,086.25	1,086.25	1,210.17	2,661.56	2,470.63	3,512.18	4,907.23	6,308.22	7,906.99
WAN (data) [Gbit/s] (inward)	0.00	0.00	0.00	0.00	0.03	0.08	0.43	0.61	0.88	1.06	1.27
WAN (MC) [Gbit/s] (outward)	0.00	0.00	0.00	0.00	0.04	0.11	0.34	0.28	0.38	0.44	0.50
WAN (MC) [Gbit/s] (inward)	0.00	0.00	0.00	0.00	0.07	0.23	0.68	0.56	0.76	0.87	1.01
WAN (ch.) [Gbit/s] (in/out)	0.00	0.19	0.38	0.77	0.77	0.00	0.00	0.00	0.00	0.00	0.00
WAN [Gbit/s] (outward)	0.00	0.19	0.38	0.77	0.80	0.11	0.34	0.28	0.38	0.44	0.50
WAN [Gbit/s] (inward)	0.00	0.19	0.38	0.77	0.86	0.31	1.12	1.18	1.64	1.94	2.27

Belle II Physik


Präzisionsvermessung des Unitaritätsdreiecks (CKM, Cabbibo-Kobayashi-Maskawa)




2012 ($\sim 1000 \text{ fb}^{-1}$ at Belle and BaBar)



Expected constraint at 50 ab^{-1}



Tau lepton flavour violation




SM predictions are small

$$\text{Br}(\tau \rightarrow \text{lepton } \gamma) \sim 10^{-54}$$

$$\text{Br}(\tau \rightarrow 3 \text{ leptons}) \sim 10^{-14}$$

Lepton Flavour Violating Tau Decays



Future Prospect at Belle II

- Sensitivity will be...

- $\tau \rightarrow \mu \gamma$: BG non-free :

$$\propto 1/\sqrt{N_\tau}$$

- $\tau \rightarrow \mu \mu \mu$: BG free


$$\propto 1/N_\tau$$











Expected limit:

$$B(\mu \gamma) \sim O(10^{-9})$$

$$B(\mu \mu \mu) \sim O(10^{-9}) \rightarrow O(10^{-10})$$



LHCb can also push down $\tau \rightarrow \mu \mu \mu$ to $O(10^{-9})$ with $O(10 \text{ fb}^{-1})$ data



Observable	Belle 2006 ($\sim 0.5 \text{ ab}^{-1}$)	SuperKEKB (5 ab^{-1})	(50 ab^{-1})	¹ LHCb (2 fb^{-1})	(10 fb^{-1})	Observable	Belle (25 fb^{-1})	Belle/SuperKEKB (5 ab^{-1})	(50 ab^{-1})	LHCb ¹ (2 fb^{-1})	(10 fb^{-1})
Hadronic $b \rightarrow s$ transitions						B_s physics	(25 fb^{-1})	(5 ab^{-1})			
$\Delta S_{\phi K^0}$	0.22	0.073	0.029	0.14		$\mathcal{B}(B_s \rightarrow \gamma\gamma)$	$< 8.7 \times 10^{-6}$	0.25×10^{-6}	-	-	-
$\Delta S_{\eta' K^0}$	0.11	0.038	0.020			$\Delta\Gamma_s^{CP}/\Gamma_s$ ($Br(B_s \rightarrow D_s^{(*)} D_s^{(*)})$)	3%	1% (model dependency)	-	-	-
$\Delta S_{K_S^0 K_S^0 K_S^0}$	0.33	0.105	0.037	-	-	$\Delta\Gamma_s/\Gamma_s$ ($B_s \rightarrow f_{CP}$ t-dependent)	-	1.2%	-	-	-
$\Delta A_{K_S^0 K_S^0}$	0.15	0.072	0.042	-	-	ϕ_s (with $B_s \rightarrow J/\psi\phi$ etc.)	-	-	-	0.02	0.01
$\mathcal{A}_{\phi\phi K^+}$	0.17	0.05	0.014			$\mathcal{B}(B_s \rightarrow \mu^+\mu^-)$	-			6 fb^{-1} for 5σ discovery	
$\phi_1^{eff}(\phi K_S)$ Dalitz		3.3°	1.5°			ϕ_3 ($B_s \rightarrow KK$)	-			$7\text{--}10^\circ$	
Radiative/electroweak $b \rightarrow s$ transitions						ϕ_3 ($B_s \rightarrow D_s K$)	-			13°	
$S_{K_S^0 \pi^0}$	0.32	0.10	0.03	-	-	Υ decays	(3 fb^{-1})	(500 fb^{-1})			
$\mathcal{B}(B \rightarrow X_s \gamma)$	13%	7%	6%	-	-	$\mathcal{B}(\Upsilon(1S) \rightarrow \text{invisible})$	$< 2.5 \times 10^{-3}$	$< 2 \times 10^{-4}$			
$A_{CP}(B \rightarrow X_s \gamma)$	0.058	0.01	0.005	-	-		($\sim 0.5 \text{ ab}^{-1}$) ¹	(5 ab^{-1})	(50 ab^{-1})		
C_9 from $\overline{A}_{FB}(B \rightarrow K^* \ell^+ \ell^-)$	-	11%	4%			Charm physics					
C_{10} from $\overline{A}_{FB}(B \rightarrow K^* \ell^+ \ell^-)$	-	13%	4%			D mixing parameters					
C_7/C_9 from $\overline{A}_{FB}(B \rightarrow K^* \ell^+ \ell^-)$	-		5%	7%		x	0.25%	0.12%	0.09%	0.25% ^{††}	
R_K		0.07	0.02	0.043		y	0.16%	0.10%	0.06%	0.05% ^{††}	
$\mathcal{B}(B^+ \rightarrow K^+ \mu \nu)$	^{††} $< 3 \mathcal{B}_{SM}$		30%	-	-	$\delta_{K\pi}$	10°	6°	4°		
$\mathcal{B}(B^0 \rightarrow K^0 \mu \nu)$	^{††} $< 40 \mathcal{B}_{SM}$		35%	-	-	$ q/p $	0.16	0.1	0.05		
Radiative/electroweak $b \rightarrow d$ transitions						ϕ	0.13 rad	0.08 rad	0.05 rad		
S_{π^0}	-	0.3	0.15			A_D	2.4%	1%	0.3%		
$\mathcal{B}(B \rightarrow X_d \gamma)$	-	24% (syst.)		-	-	New particles ^x					
Leptonic/semileptonic B decays						$\gamma\gamma \rightarrow Z(3930) \rightarrow DD^*$		$> 3\sigma$			
$\mathcal{B}(B^+ \rightarrow \tau^+ \nu)$	3.5σ	10%	3%	-	-	$B \rightarrow KX(3872)(\rightarrow D^0 D^{*0})$		400 events			
$\mathcal{B}(B^+ \rightarrow \mu^+ \nu)$	^{††} $< 2.4 \mathcal{B}_{SM}$	4.3 ab^{-1} for 5σ discovery		-	-	$B \rightarrow KX(3872)(\rightarrow J/\psi \pi^+ \pi^-)$		1250 events			
$\mathcal{B}(B^+ \rightarrow D \tau \nu)$	-	8%	3%	-	-	$B \rightarrow KZ^+(4430)(\rightarrow \psi' \pi^+)$		1000 events			
$\mathcal{B}(B^0 \rightarrow D \tau \nu)$	-	30%	10%	-	-	$e^+ e^- \rightarrow \gamma_{ISR} Y(4260)(\rightarrow J/\psi \pi^+ \pi^-)$		3000 events			
LFV in τ decays (U.L. at 90% C.L.)						Electroweak parameters					
$\mathcal{B}(\tau \rightarrow \mu \gamma)$ [10^{-9}]	45	10	5	-	-	$\sin^2 \Theta_W$	-	3×10^{-4}			
$\mathcal{B}(\tau \rightarrow \mu \eta)$ [10^{-9}]	65	5	2	-	-						
$\mathcal{B}(\tau \rightarrow \mu \mu \mu)$ [10^{-9}]	21	3	1	-	-						
Unitarity triangle parameters											
$\sin 2\phi_1$	0.026	0.016	0.012	~ 0.02	~ 0.01						
ϕ_2 ($\pi\pi$)	11°	10°	3°	-	-						
ϕ_2 ($\rho\pi$)	$68^\circ < \phi_2 < 95^\circ$	3°	1.5°	10°	4.5°						
ϕ_2 ($\rho\rho$)	$62^\circ < \phi_2 < 107^\circ$	3°	1.5°	-	-						
ϕ_2 (combined)		2°	$\lesssim 1^\circ$	10°	4.5°						
ϕ_3 ($D^{(*)} K^{(*)}$) (Dalitz mod. ind.)	20°	7°	2°	8°							
ϕ_3 ($DK^{(*)}$) (ADS+GLW)	-	16°	5°	$5\text{--}15^\circ$							
ϕ_3 ($D^{(*)} \pi$)	-	18°	6°								
ϕ_3 (combined)		6°	1.5°	4.2°	2.4°						
$ V_{us} $ (inclusive)	6%	5%	3%	-	-						
$ V_{us} $ (exclusive)	15%	12% (LQCD)	5% (LQCD)	-	-						
$^{†††}\bar{\rho}$	20.0%		3.4%								
$^{†††}\bar{\eta}$	15.7%		1.7%								

Very broad physics program within Belle II!

See arXiv:1002.5012: “Physics at Super B Factory”

Very broad physics program within Belle II!




See arXiv:1002.5012: "Physics at Super B Factory"

$B \rightarrow \tau \nu$ [PRL110, 131801 (2013)]

- Data sample 711 fb⁻¹
- Hadronic tagging
- 2D fit in excess calorimeter energy (E_{ECL}) and M_{miss}^2


$$\mathcal{B}(B \rightarrow \tau \nu) = (0.72^{+0.27}_{-0.25} \pm 0.11) \times 10^{-4}$$

- Significance: 3σ



Tension in CKM fit between $\sin(2\phi_1)$, $\mathcal{B}(B \rightarrow \tau \nu) \sim 1.6\sigma$


Measurement of A_{FB}



$$A_{FB}(s) \approx \rho \frac{3G_F}{4\sqrt{2}\pi\alpha} \frac{sM_Z^2}{s - M_Z^2} g_A^e g_A^\mu$$

Stat. errors are x1000!
(otherwise not visible)

Belle 2: $\sigma_{\text{stat}}(A_{FB})/A_{FB} \approx 0.1\%$



$$\Delta\rho^{\text{new}} = \frac{\hat{\Pi}_{WW}^{\text{new}}(0) - \hat{c}_Z^2 \hat{\Pi}_{ZZ}^{\text{new}}(0)}{M_W^2(1 - \Delta\hat{r}_W)} \equiv \frac{\alpha T}{1 - \Delta\hat{r}_W} \approx \hat{\alpha}_Z T$$

mit Belle II Sensitivität auf Compositeness $\Lambda \leq 33 \text{ TeV}$
bei Interpretation im Rahmen von Contact Interactions)

Is there $c\bar{c}$ admixture inside the $X(3872)$?

$$|X(3872)\rangle = c_1 |c\bar{c}\rangle + c_2 |\bar{D}^0 D^{0*}\rangle$$

Belle, 711 fb⁻¹

E. Panzenboeck



(Göttingen/Nara COTUTELLE program)

Hadron 2013. LLWI 2014

Search for $X(3872) \rightarrow \chi_{c1} \pi^+ \pi^-$


No signal observed
(no indication of admixture)

MC simulation, assuming
 $\mathcal{B}(\chi_{c1} \pi^+ \pi^-) = \mathcal{B}(J/\psi \pi^+ \pi^-)$




Belle II Search for New Physics

$$\Delta_{NP} = (\text{exp.} - \text{SM}) \pm \sqrt{(\sigma_{\text{exp}})^2 + (\sigma_{\text{SM}})^2}$$
$$= c / (M_{NP})^{n=2}$$



$$\Delta_{NP} = (\text{exp.} - \text{SM}) \pm \sqrt{(\sigma_{\text{exp}})^2 + (\sigma_{\text{SM}})^2}$$


→ improve stat. and syst. errors



new detectors (PXD, TOP),
reduce hadronic uncertainties
(lattice QCD), ...

$$\Delta_{NP} = c / (M_{NP})^n$$

→ find decays with smaller n



e.g. penguin decays
n=1 for $b \rightarrow s\gamma$
 $m > \sim 10^6$ TeV in reach

Bounds on M_{NP} (or scale Λ) based upon existing data

Open Symposium on the European Strategy for Particle Physics

Isidori, Teubert, arXiv:1402.2844[hep-ph]

$M_{NP} \geq 140 \text{ TeV}$ (standard left-handed CKM couplings)

Operator	Bounds on Λ in TeV ($c_{NP} = 1$)		Bounds on c_{NP} ($\Lambda = 1 \text{ TeV}$)		Observables
	Re	Im	Re	Im	
$(\bar{s}_L \gamma^\mu d_L)^2$	9.8×10^2	1.6×10^4	9.0×10^{-7}	3.4×10^{-9}	$\Delta m_K; \epsilon_K$
$(\bar{s}_R d_L)(\bar{s}_L d_R)$	1.8×10^4	3.2×10^5	6.9×10^{-9}	2.6×10^{-11}	
$(\bar{c}_L \gamma^\mu u_L)^2$	1.2×10^3	2.9×10^3	5.6×10^{-7}	1.0×10^{-7}	$\Delta m_D; q/p _D, \phi_D$
$(\bar{c}_R u_L)(\bar{c}_L u_R)$	6.2×10^3	1.5×10^4	5.7×10^{-8}	1.1×10^{-8}	
$(\bar{b}_L \gamma^\mu d_L)^2$	6.6×10^2	9.3×10^2	2.3×10^{-6}	1.1×10^{-6}	$\Delta m_{B_d}; \sin(2\beta)$ from $B_d \rightarrow \psi K$
$(\bar{b}_R d_L)(\bar{b}_L d_R)$	2.5×10^3	3.6×10^3	3.9×10^{-7}	1.9×10^{-7}	
$(\bar{b}_L \gamma^\mu s_L)^2$	1.4×10^2	2.5×10^2	5.0×10^{-5}	1.7×10^{-5}	$\Delta m_{B_s}; \sin(\phi_s)$ from $B_s \rightarrow \psi \phi$
$(\bar{b}_R s_L)(\bar{b}_L s_R)$	4.8×10^2	8.3×10^2	8.8×10^{-6}	2.9×10^{-6}	

Table 1. Bounds on representative dimension-six $\Delta F = 2$ operators [96, 97]. The bounds on Λ are evaluated assuming an effective coupling $1/\Lambda^2$ (i.e. setting $c_{NP} = 1$). Alternatively, the bounds on the respective c_{NP} are obtained assuming $\Lambda = 1 \text{ TeV}$. In the last column we list the observables used to set such bounds; the observables related to CPV are separated from the CP conserving ones with semicolons.

Isidori, Teubert, arXiv:1402.2844[hep-ph]

→ Λ_{NP} scales with at most $N^{1/4}$,

where N is relative increase of number of events

$1 \text{ ab}^{-1} \rightarrow 50 \text{ ab}^{-1}$ ($50^{0.25} = 2.65$)

e.g. $\Lambda_{\text{NP}}=100 \text{ TeV}$ (Belle) → 265 TeV (Belle II)

If physics beyond the SM respects the SM gauge symmetry, as we expect from general arguments, the low-energy amplitudes describing the transition of a fermion ψ_i to a fermion ψ_j (of different flavour) can be decomposed in the following general form

$$\mathcal{A}(\psi_i \rightarrow \psi_j + X) = \mathcal{A}_0 \left[\frac{c_{\text{SM}}}{M_W^2} + \frac{c_{\text{NP}}}{\Lambda^2} \right] , \quad (16)$$

where Λ is the energy scale of the new degrees of freedom and the SM result is recovered in the limit $c_{\text{NP}} \rightarrow 0$. This structure is completely general: the coefficients $c_{\text{SM(NP)}}$ may include appropriate CKM coefficient factors and eventually a $\sim 1/(16\pi^2)$ suppression if the amplitude is loop-mediated. Given our ignorance about the c_{NP} coefficients, the values of the scale Λ probed by present experiments vary over a wide range. However, the general result in Eq. (16) allows us to predict how these bounds will improve with future experiments: increasing the statistic on a given observable, the corresponding bound on Λ scales at most as $N^{1/4}$, where N is the relative increase in the number of events used to measure the observable.² From Eq. (16) it is also clear that indirect searches can probe NP scales well above the TeV for models where ($c_{\text{SM}} \ll c_{\text{NP}}$), namely models which do not respect the symmetries and the symmetry-breaking pattern of the SM.