



Deutsches Elektronen-Synchrotron
(DESY), Hamburg




Double Parton Scattering experimental results

Paolo Gunnellini




DESY Tuesday Seminar
Hamburg - Zeuthen
19th-20th May 2015



- ① Introduction
- ② Choice of sensitive observables
- ③ Choice of physics channels
- ④ Summary of recent DPS measurements
- ⑤ Extraction of the DPS contribution
- ⑥ Other DPS-sensitive measurements
- ⑦ Summary and conclusion



Introduction: the Underlying Event



Hard scattering
Initial and Final State Radiation
Multiple Parton Interaction (MPI)
Beam-beam remnants



$$P_A = \frac{\sigma_A}{\sigma_{tot}^{pp}}$$

$$P_B = \frac{\sigma_B}{\sigma_{tot}^{pp}}$$

$$\sigma_{AB}^{DPS} \propto \frac{m}{2} P_A P_B \sigma_{tot}^{pp}$$




$$\sigma_{AB}^{DPS} = \frac{m \sigma_A \sigma_B}{2 \sigma_{eff}}$$

$$\sigma_{eff} \ll \sigma_{tot}^{pp}$$

Need for correlations!

Introduction: why do we care about DPS?

- Increasing contribution at the LHC when going to higher energy
 - Sizeable background for LHC processes (SM and searches), e.g. Higgsstrahlung
 - Information about the structure of the proton, i.e. parton correlations




**And...increasing interest and
number of entries in Spires!**

Choice of sensitive observables (I): a four-jet scenario

A four-jet final state may arise from one or two chains:



- the two additional jets may be produced via PS or a 2nd hard scattering



Various kinematical observables can discriminate the two processes:

$$\Delta_{soft}^{rel} p_T = \frac{|p_T(j_i, j_k)|}{|p_T(j_i)| + |p_T(j_k)|}$$

$$\Delta S = \arccos \left(\frac{\vec{p}_T(j^i, j^k) \cdot \vec{p}_T(j^l, j^m)}{|\vec{p}_T(j^i, j^k)| \cdot |\vec{p}_T(j^l, j^m)|} \right)$$



! Selection of jet pairs at different scales helps the jet association !

Choice of sensitive observables (II): a four-jet scenario



Which regions of the phase space are interesting for DPS detection?

Studies of SPS and DPS contributions performed with PYTHIA8:

Selection of a four-jet final state in $|\eta| < 4.7$ at two different p_T thresholds (20 and 50 GeV)

A SIMPLE scenario:


- SPS: MPI contribution switched off
- DPS: Two hard scatterings at the parton level forced to happen w/o parton shower



Different regions of the phase space are filled by the two processes

Discriminating power



Choice of physics channels



Measurement of a four-jet final state


Event selection

Exactly four jets in the final state in $|\eta| < 4.7$:
 2 jets: $p_T > 50$ GeV (hard), 2 jets: $p_T > 20$ GeV (soft)



$$\Delta_{\text{soft} p_T}^{\text{rel}} = \frac{|\vec{p}_T(j_i, j_k)|}{|\vec{p}_T(j_i)| + |\vec{p}_T(j_k)|}$$

$$\Delta S = \arccos \left(\frac{\vec{p}_T(j^i, j^k) \cdot \vec{p}_T(j^l, j^m)}{|\vec{p}_T(j^i, j^k)| \cdot |\vec{p}_T(j^l, j^m)|} \right)$$



Soft jets are expected to be produced also by a 2nd scattering


PRD 89 (2014) 092010

ΔS and $\Delta_{\text{soft} p_T}^{\text{rel}}$ sensitive to MPI contribution: → ROOM for DPS!


Measurement of a four-jet final state with b-jets

Event selection

Selection of at least four jets with $p_T > 20$ GeV:
2 b-jets: $|\eta| < 2.4$, 2 other jets: $|\eta| < 4.7$



$$\Delta S = \arccos \left(\frac{\vec{p}_T(j^i, j^k) \cdot \vec{p}_T(j^l, j^m)}{|\vec{p}_T(j^i, j^k)| \cdot |\vec{p}_T(j^l, j^m)|} \right)$$



Additional jets
may be
produced also
by DPS

CMS-FSQ-13-010




Sensitivity to higher orders..

..but also to MPI!

Measurement of a W+dijet final state

Event selection

Presence of a muon with $p_T > 35$ GeV in $|\eta| < 2.1$ and $E_T^{miss} > 50$ GeV
+ at least 2 jets: $p_T > 20$ GeV in $|\eta| < 2.0$




The jets are expected to be produced also by a 2nd scattering

JHEP 03 (2014) 032


Sensitivity to DPS!

How can one extract the DPS contribution from the measured observables?



How to extract σ_{eff} : the template method

- Measurement of DPS-sensitive observables
- Definition of signal and background
- Fit the relative fraction of signal and background
- The signal fraction translates into a value for σ_{eff}



From Ramandeep Kumar,
Talk at MPI@LHC 2012

W + jets channel

$$\sigma_{\text{eff}} = \frac{\sigma_A \cdot \sigma_B}{\sigma_{\text{DPS}}}$$

$$\sigma_{\text{eff}} = \frac{N_A^{\text{ev}}}{N_{A+B(\text{DPS})}^{\text{ev}}} \cdot \sigma_B$$

$$\sigma_{\text{eff}} = \frac{N_A^{\text{ev}}}{f_{\text{DPS}} \cdot N_{A+B}^{\text{ev}}} \cdot \sigma_B$$

Extraction of σ_{eff} from W+dijet final state (ATLAS)

First measurement of DPS signal at 7 TeV

New J. Phys. 15 (2013) 033038


SELECTION: 2j with $p_T > 20$ GeV in $|y| < 2.8$, standard W selection

CONSIDERED OBSERVABLES: normalized $\Delta_{\text{jets}}^n = \frac{|\vec{p}_T^{1j} + \vec{p}_T^{2j}|}{|\vec{p}_T^{1j}| + |\vec{p}_T^{2j}|}$

BACKGROUND: ALPGEN+HERWIG+JIMMY with hard MPI excluded

SIGNAL: selection of two independent collisions from data

DRIVING UNCERTAINTY: model dependence



$$\sigma_{\text{eff}} = \frac{N_{W+0j}}{f_{DPS} \cdot N_{W+2j}} \cdot \sigma_{2j}$$

with $f_{DPS} = 8.0\%$ and

$$\frac{N_{W+0j}}{N_{W+2j}} = 23$$

$$\sigma_{\text{eff}} = 15.0 \pm 3 \text{ (st.) } {}^{+5}_{-3} \text{ (sys.) mb}$$



Extraction of σ_{eff} from W+dijet final state (CMS)

CONSIDERED OBSERVABLES: normalized ΔS and $\Delta^{rel} p_T$

BACKGROUND: MADGRAPH+P8 with hard MPI above 15 GeV excluded

SIGNAL: Two mixed independent scatterings generated with P8 and MG+P8

DRIVING UNCERTAINTY: model dependence



$$\sigma_{eff} = \frac{N_{W+0j}}{f_{DPS} \cdot N_{W+2j}} \cdot \sigma_{2j}$$

$$f_{DPS} = 5.5\%,$$

$$\frac{N_{W+0j}}{N_{W+2j}} = 27.8$$

JHEP 03 (2014) 032

$$\sigma_{eff} = 20.7 \pm 0.8 \text{ (stat.)} \pm 6.6 \text{ (syst.)} \text{ mb}$$

The inclusive fit method

Experimental difficulties of the template method

→ **How to define the background?**

- Good to exclude hard MPI..but no such possibility in some generators

→ **How to define exclusive and inclusive events?**

- N_{W+0j} and N_{W+2j} are sensitive to the jet scales

→ These issues have an impact on the systematic uncertainty!

Is there a way out?

The inclusive fit method


- Run predictions for different choices of UE parameters
- Fit the MC predictions to the considered observables
- Improve the data description with the examined model
- (...look at the corresponding σ_{eff} ...)




Extraction of σ_{eff} in four-jet final states

$$\text{Minimization of the binned } \chi^2 = \sum_O \sum_{b \in O} \frac{(MC^b - DATA^b)^2}{\Delta_b^2}$$

Normalized ΔS in $pp \rightarrow 4j$ in $|\eta| < 4.7$ at $\sqrt{s} = 7$ TeV




Normalized ΔS in $pp \rightarrow 4j$ in $|\eta| < 4.7$ at $\sqrt{s} = 7$ TeV




MINIMIZ.



Normalized ΔS in $pp \rightarrow 4j$ in $|\eta| < 4.7$ at $\sqrt{s} = 7$ TeV



Normalized $\Delta_{soft}^{rel} p_T$ in $pp \rightarrow 4j$ in $|\eta| < 4.7$ at $\sqrt{s} = 7$ TeV




$$\sigma_{eff} = 19.0^{+4.7}_{-3.0} \text{ mb}$$

LEFT: ΔS
RIGHT: $\Delta_{soft}^{rel} p_T$

CMS-GEN-14-001

Where do we stand now?


- UE measurements sensitive to soft MPI
- Observables sensitive to DPS measured in various final states
- Values of σ_{eff} extracted in W+dijet and four-jet
- Ongoing extraction for the other channels



It is not all..

- CMS Coll. *Measurement of Prompt Double J/ ψ Production at 7 TeV* JHEP1409(2014)094
- CMS Coll. *Measurement of the cross section and angular correlations for associated production of a Z boson with b hadrons* JHEP12(2013)039
- CMS Coll. *Measurement of the production cross section for a W boson and two b jets at 7 TeV* Phys.Lett.B735(2014)
- ATLAS Coll. *Associated production of prompt J/ ψ mesons and W boson* JHEP04(2014)172
- ATLAS Coll. *Measurement of the cross-section for W boson production in association with b-jets* New J.Phys.15(2013)033038


No extraction of
a value of σ_{eff}
but indication of
need for DPS !




Angular correlations in Z+b-hadrons final states

Event selection




Presence of two leptons with $p_T > 20$ GeV in $|\eta| < 2.4$ with invariant mass close to the Z peak and two b-hadrons with $p_T > 15$ GeV in $|\eta| < 2$



$$\Delta\phi = |\Delta\phi_{b1} - \Delta\phi_{b2}|$$



$$\Delta R = \sqrt{\Delta\phi_{b1}^2 + \Delta\eta_{b2}^2}$$




JHEP 03
(2014) 032

Data compatible with predictions at parton level with
DPS contribution ($\sigma_{eff} \sim 25-30$ mb) included

What to do next?

→ Measurements for LHC Run 2



Joined effort between phenomenological and experimental community

- **Important to study first the sensitivity of the physics channel and the considered observables**
- **Important to produce unfolded results in order to be able to compare predictions from any model**
- Double parton scattering is essential for proton structure as well as for background to physics searches
- Several final states can be used for DPS detection
 - W+jets, four-jets, two b- + two other jets...
- **The measured final states clearly indicate the need for DPS for describing the experimental results**
- **Future: measure energy dependence
get a unified picture of DPS with UE- and MB-sensitive measurements**

- **Important to study first the sensitivity of the physics channel and the considered observables**
- **Important to produce unfolded results in order to be able to compare predictions from any model**
- Double parton scattering is essential for proton structure as well as for background to physics searches
- Several final states can be used for DPS detection
 - W+jets, four-jets, two b- + two other jets...
- **The measured final states clearly indicate the need for DPS for describing the experimental results**
- **Future: measure energy dependence
get a unified picture of DPS with UE- and MB-sensitive measurements**

THANK YOU!



BACK-UP SLIDES



Determination of σ_{eff} in the four-jet channel

Tuning the four-jet observables (Phys.Rev., D89, 2014) with PYTHIA8

Parameter	CDPSTP8S1-4j	CDPSTP8S2-4j	4C
MultipleInteractions:expPow	1.16	0.6921	2.0
MultipleInteractions:ecmPow	0.19*	0.345	0.19
MultipleInteractions:pT0ref	2.09*	2.125	2.09
BeamRemnants:reconnectRange	1.5* *=unchanged wrt 4C	6.526	1.5
χ^2/NdF	0.75	0.42	-
σ_{eff} (mb)	$21.3^{+1.7}_{-1.3}$	$19.0^{+4.7}_{-3.0}$	30.3

$$\sigma_{\text{eff}} = 19.0^{+4.7}_{-3.0} \text{ mb}$$

$\rightarrow \sigma_{\text{eff}}$ (Tune 4C) ~ 30.3 mb



DPS-based tune
propagated to
MADGRAPH ME


Simulation of UE
independent on the
used matrix element

LEFT: ΔS
RIGHT: $\Delta_{\text{soft},PT}^{\text{rel}}$


CMS-GEN-14-001

Choice of sensitive observables


Normalized ΔS in $p\bar{p} \rightarrow 4j$ in $|\eta| < 4.7$ at $\sqrt{s} = 7$ TeV



Normalized Δ_{soft}^{rel} in $p\bar{p} \rightarrow 4j$ in $|\eta| < 4.7$ at $\sqrt{s} = 7$ TeV




Normalized $\Delta\phi^{soft}$ in $p\bar{p} \rightarrow 4j$ in $|\eta| < 4.7$ at $\sqrt{s} = 7$ TeV



...but also the phase space thresholds matter!!


Normalized ΔS in $p\bar{p} \rightarrow 2b2j$ at $\sqrt{s} = 7$ TeV



LEFT: four jets selected applying
the same p_T

RIGHT: four jets with a rapidity
cut applied between the most
remote jets

arXiv 1503.08022



D0 DPS analysis: $\gamma+3\text{jets}$ and $\gamma+b/c\text{ jet}+2\text{jets}$

SELECTION 1: $p_T^\gamma > 26 \text{ GeV}$, $p_T^{\text{lead}} > 35 \text{ GeV}$, $15 < p_T^{\text{others}} < 35 \text{ GeV}$ in $|\eta| < 2.5$



SELECTION 2: $p_T^\gamma > 26 \text{ GeV}$, $p_T^b > 35 \text{ GeV}$, $15 < p_T^{\text{others}} < 35 \text{ GeV}$ in $|\eta| < 2.5$

CONSIDERED OBSERVABLES: normalized ΔS btw γ -j and dijet systems

BACKGROUND: SHERPA sample with MPI simulation off

SIGNAL: Two independent events recorded from data

DRIVING UNCERTAINTY: model dependence (only samples with MPI off!)



$$\sigma_{\text{eff}} \propto \frac{N_{DI}}{N_{DP}} \cdot \frac{\epsilon_{DP}}{\epsilon_{DI}} \cdot \sigma_{\text{hard}}$$

with $f_{\gamma+3j}^{DP} = 21\%$ and

$f_{\gamma+b/c\text{ jet}+2j}^{DP} = 17\%$

$\gamma+3\text{jets}$

$$\sigma_{\text{eff}} = 12.7 \pm 0.2 \pm 1.3 \text{ mb}$$

$\gamma+b/c\text{ jet}+2\text{jets}$

$$\sigma_{\text{eff}} = 14.6 \pm 0.6 \pm 3.2 \text{ mb}$$


Phys. Rev. D 89, 072006 (2014)

Recommendations for DPS extraction

	CMS	ATLAS	D0/CDF
Background and signal should cover the full phase space	✓	✓	✗
Use more than one MC event generator to correctly evaluate the model dependence and the systematic uncertainty	✓	✓	✓
Use more than one variable for the DPS determination	✓	✗	✗

BUT..difficult to define the background template in the same way with different generators!

The proposed new approach





A FEW REMARKS WHEN USING THE TUNING METHOD:

- ① Investigation of the contribution of different matrix elements used with the same UE simulation
- ② Use more than one MC event generator to study the DPS contribution needed in different models
- ③ Use more than one variable for the DPS determination
- ④ Check if the new set of parameters spoil description of more inclusive distribution

How does the new tune perform in the UE description?

Measurement of charged particle mult. and p_T sum in hadronic events

ATLAS Coll. Phys.Rev. D83 (2011) 112001



Tune	σ_{eff} (mb)
P8 4C	30.3
CDPSP8S2	$19.0^{+4.7}_{-3.0}$

A tension appears between the description of "softer" and "harder" MPI within the same framework



Charged particle multiplicity (top) and p_T sum (bottom) for transverse (left) and toward (right) regions



CMS-GEN-14-001

How to fix this?

→ Attempt to implement in a tune a value of σ_{eff} compatible with experimental measurements

HERWIG++ case: $\sigma_{\text{eff}} = \frac{28\pi}{\mu}$, with μ inverse proton radius

Tune UE-EE-5C ([arXiv:1307.5015](https://arxiv.org/abs/1307.5015)) : $\sigma_{\text{eff}} = 15$ mb (CDF)



Slight underestimation
of the low ΔS region

LEFT: ΔS
RIGHT: $\Delta_{\text{soft}}^{\text{rel}} p_T$

Another approach:
Dynamical approach to MPI contribution
([arXiv:1503.08246](https://arxiv.org/abs/1503.08246))

- Introduction of x - and scale-dependence for values of σ_{eff}
- Inclusion of 1×2 mechanisms


It is not all..

- ATLAS Coll. *Associated production of prompt J/ψ mesons and W boson* JHEP 04 (2014) 172
- LHCb Coll. *Prompt charm production in pp collisions* HEP 1206 (2012) 141
- ATLAS Coll. *Measurement of the cross-section for W boson production in association with b -jets* New J. Phys. 15 (2013) 033038
- LHCb Coll. *Study of forward Z +jet production in pp collisions* JHEP 01 (2014) 033
- CMS Coll. *Measurement of the cross section and angular correlations for associated production of a Z boson with b hadrons* JHEP 12 (2013) 039
- CMS Coll. *Measurement of Prompt Double J/ψ Production in pp Collisions* JHEP 1409 (2014) 094
- ALICE Coll. *J/ψ production as a function of charged particle multiplicity in pp collisions at 7 TeV* Phys.Lett.B 712, 165 (2012)


No extraction of a value of σ_{eff} but clear indication of need for DPS !

Cross section measurements sensitive to DPS

ATLAS Collaboration:
"Measurements of
 $W + \text{prompt } J/\psi$ in pp
collisions at 7 TeV"
JHEP 04 (2014) 172



ATLAS Collaboration:
"Measurement of the cross-section
for W boson production in
association with b -jets"
New J. Phys. 15 (2013) 033038



Measurements compatible with a DPS contribution with $\sigma_{\text{eff}} \sim 15-20$ mb

Keypoints of the choice of variables

- Observables which consider the whole final state are more sensitive to DPS
 - ΔS , sum of transverse momenta, energy of the four objects
- A large phase space for additional radiation reduces the DPS sensitivity
 - Better selection with objects close in transverse momentum
 - BUT..more complicated migration effects (and unfolding procedure)

CMS strategy for the DPS measurement

1st step




Corrected
distributions
DPS-sensitive
variables

2nd step

Data interpretation
and unambiguous
definition
of signal and
background templates

3rd step

Extraction of the DPS
fraction and study of
the process
dependence




- Compare the data to your own favourite predictions!

4th (future) step: differential distributions with high luminosities..

Cross section measurements sensitive to DPS (I)


Event selection

Presence of two pairs of same-sign muons in $|\eta| < 2.2$; the two pairs must have invariant mass close to J/ψ



$$\sigma(J/\psi J\psi + X)$$

$$1.49 \pm 0.07 \pm 0.13 \text{ nb}$$



Correction and phase-space extrapolation assuming unpolarized production


SPS background should dominate the fall at low Δy
DPS expected to fill the high Δy region

Useful baseline for building reliable models of J/ψ production before extracting DPS signal

Cross section measurements sensitive to DPS (III)

Event selection

Presence of a muon with $p_T > 25$ GeV in $|\eta| < 2.1$, $E_T^{miss} > 45$ GeV and two b-tagged jets with $p_T > 25$ GeV in $|\eta| < 2.4$





$$\Delta R = \sqrt{\Delta\phi_b^2 + \Delta\eta_b^2}$$

Good agreement with SM predictions
(MadGraph+Pythia8)

$$\sigma(W + b\bar{b}) = 0.53 \pm 0.05 \pm 0.09 \pm 0.06 \pm 0.01 \text{ pb}$$





Good agreement with MCFM
predictions corrected with DPS
contribution ($\sigma_{DPS} \sim 0.08$ pb)



Measurement of a final state with $\gamma + 3$ jets

Event selection

Selection of a photon and at least three jets in $|\eta| < 2.5$:
 $\gamma+1$ jet: $p_T > 75$ GeV, 2 jets: $p_T > 20$ GeV



Soft jets are expected to be produced also by a 2nd scattering

$$\Delta\phi(j_i, j_k) = \phi_i - \phi_k$$

$$\Delta_{\text{soft } PT}^{\text{rel}} = \frac{|\vec{p}_T(j_i, j_k)|}{|\vec{p}_T(j_i)| + |\vec{p}_T(j_k)|}$$

$$\Delta S = \arccos \left(\frac{\vec{p}_T(\gamma, j^k) \cdot \vec{p}_T(j^l, j^m)}{|\vec{p}_T(\gamma, j^k)| \cdot |\vec{p}_T(j^l, j^m)|} \right)$$

No difference between predictions with and w/o MPI


Little DPS sensitivity!

Choice of the physics channel


$$\sigma_{AB}^{DPS} = \frac{m}{2} \frac{\sigma_A \sigma_B}{\sigma_{eff}}$$

Internal structure of the proton
DPS background for any physics channel

→ Which channels can be used to look for DPS signals?



The Compact Muon Solenoid experiment



- Silicon tracker immersed in a 3.8 T magnetic field
- Wide calorimeter coverage
- Excellent jet energy resolution and muon detection efficiency
- Particle Flow technique for jet reconstruction

