


Second Stage of WISPDIMX Measurements



Andrei Lobanov
MPIfR Bonn / University of Hamburg



Two+ Clouds of SM

- ❑ Standard Model: $SU(3) \times SU(2) \times U(1)$ gives us (nearly) all things we may need in life.
- ❑ „*The beauty and clearness of the dynamical theory, [...], is at present obscured by two clouds [...]*” (Lord Kelvin, 1900) ... still true today?
 - gravitation and dark energy
 - ...plus some „lesser evils“ such as dark matter, strong CP problem, etc...
- ❑ Most of the solutions proposed invoke a „hidden sector“ of the global parameter space, weakly coupled to „normal matter“ of the SM through weakly interacting massive (WIMP) or slim (WISP) particles.

Three generations of matter (fermions)


	I	II	III	
mass	2.4 MeV/c ²	1.27 GeV/c ²	171.2 GeV/c ²	0
charge	2/3	2/3	2/3	0
spin	1/2	1/2	1/2	1
name	u up	c charm	t top	γ photon
	4.8 MeV/c ²	104 MeV/c ²	4.2 GeV/c ²	0
	-1/3	-1/3	-1/3	0
	1/2	1/2	1/2	1
Quarks	d down	s strange	b bottom	g gluon
	<2.2 eV/c ²	<0.17 MeV/c ²	<15.5 MeV/c ²	91.2 GeV/c ²
	0	0	0	0
	1/2	1/2	1/2	1
	ν _e electron neutrino	ν _μ muon neutrino	ν _τ tau neutrino	Z ⁰ Z boson
	0.511 MeV/c ²	105.7 MeV/c ²	1.777 GeV/c ²	80.4 GeV/c ²
	-1	-1	-1	±2
	1/2	1/2	1/2	1
Leptons	e electron	μ muon	τ tau	W [±] W boson

Gauge bosons



Many Faces of WISP

- ❑ WISP, and *axions* and *hidden photons* in particular, are strong dark matter candidates. Direct detection of WISP or putting bounds on their properties are important tasks for cosmology and particle physics.
- ❑ A number of experimental methods have been employed, both for laboratory and astrophysical searches – all relying on WISP interaction (coupling, kinetic mixing) with ordinary matter (most often: photons).
- ❑ Radio (24 MHz—2.4 THz): excellent sensitivity to WISP signal and access to DM/DE relevant particle mass ranges ($0.1\mu\text{eV}$ – 10meV)





WISP Dark Matter eXperiment

Direct WISP dark matter searches in the $0.8\text{--}2.0\ \mu\text{eV}$ mass range

Sebastian Baum¹, Babette Döbrich², Dieter Horns¹, Joerg Jaeckel³,
Reinhard Keller⁴, Denis Kostin², Michael Kramer⁴, Axel Lindner²,
Andrei Lobanov^{4,1}, Wolf-Dietrich Möller², Javier Redondo^{5,6}, Andreas Ringwald²,
Jacek Sekutowicz², Alexey Sulimov², Dieter Trines², Alexander Westphal²

¹ *Institut für Experimentalphysik, Universität Hamburg, Luruper Chaussee 149, D-22761 Hamburg, Germany*

² *Deutsches Elektronen Synchrotron DESY, Notkestrasse 85, D-22607 Hamburg, Germany*

³ *Institut für Theoretische Physik, Universität Heidelberg, Philosophenweg 16, D-69120 Heidelberg, Germany*

⁴ *Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany*


⁵ *Ludwig-Maximilians-Universität, Theresienstr. 37, D-80333 München, Germany*

⁶ *Max-Planck-Institut für Physik, Föhringer Ring 6, D-80805 München, Germany*

WISPDMMX Overview

- ❑ WISP Dark Matter eXperiment (WISPDMMX) is a pioneering search for hidden photon and axion dark matter in the $0.8\text{-}2.0\text{ }\mu\text{eV}$ range, exploring the particle masses below the mass range covered by ADMX.

- ❑ WISPDMMX utilizes a HERA 208-MHz resonant cavity and a 40 dB amplifier chain, and plans to make use of a strong magnet (e.g. 1.15 T H1 magnet).
- ❑ Uses multiple resonant modes in the 200-600 MHz range.
- ❑ Completed Phase 1: hidden photon searches at nominal resonances of the cavity.
- ❑ Currently in Phase 2: HP searches with cavity tuning
- ❑ Phase 3: ALP searches



1 – 208 MHz HERA cavity; 2 – cavity ports; 3 – antenna probes; 4 – WantCom 22 dB amplifier; 5 – MITEQ 18 dB amplifier; 6 – network analyzer (HP 85047A); 7 -- control computer, with onboard digitizer (Alazar ATS-9360, 1.8Gs/s)

Searching for WISP DM ...

❑ Hidden photons (γ'):

-- spontaneous photon conversion (kinetic mixing), $\gamma \leftrightarrow \gamma'$

„Haloscope“ experiments: Coupling strength (mixing angle):

$$\chi \propto t_{\text{mes}}^{-1/4} \text{SNR}^{1/2} T_n^{1/2} V_0^{-1/2} Q_0^{-1/2} \mathcal{G}_{\gamma'}^{-1/2} \rho_0^{-1/2} Q_{\gamma'}^{-1/4} m_{\gamma'}^{-1/4}$$

❑ Axions and axion-like particles (ϕ):

-- two-photon coupling (Primakoff process), $\phi \leftrightarrow \gamma + \gamma$, with B-field as a virtual photon


„Haloscope“ experiments: Coupling strength:

$$g[\text{GeV}^{-1}] \propto t_{\text{mes}}^{-1/4} \text{SNR}^{1/2} T_n^{1/2} B_0^{-1/4} V_0^{-1/2} Q_0^{-1/2} \mathcal{G}_{\phi}^{-1/2} \rho_0^{-1/2} Q_{\phi}^{-1/4} m_{\phi}^{3/4}$$

t_{mes} , SNR – measurement time and SNR; T_n – noise temperature; V_0 , Q_0 – cavity volume and quality factors; B_0 – magnetic field strength; $\mathcal{G}_{\phi/\gamma}$ – form factor; ρ_0 – DM density; $Q_{\phi/\gamma}$ – quality factor of DM signal; $m_{\phi/\gamma}$ – particle mass

Accessible Resonant Modes

- Five resonant modes identified which have non-zero form factors for hidden photon measurements.
- Outside resonance:
 $G_f \approx 0.0018$ –
hence measurements in the entire spectral range could also be used for constraining χ .



Mode	TM ₀₁₀	TE ₁₁₁	TM ₀₂₀	TE ₁₁₂	TE ₁₂₀	TM ₀₃₀
f/MHz	199	308	428	508	560	580
Q	53000	61000	47000	73000	72000	125000
\mathcal{G}	0.43	0.67	0.32	0.019	0.036	0.061


Phase 1: Nominal Resonances

- ❑ Recording broadband (600 MHz) signal; useful range: 180--600 MHz; frequency resolution $\Delta\nu = 572$ Hz.
- ❑ 40.3 dB amplification; effective measurement time of 1.7 hours.
- ❑ No HP signal detected. Gaussian distribution of measured power around rms; no daily modulation; no significant RFI signals.
- ❑ Limits, assuming $\rho_0 = 0.39$ GeV/cm³ and $Q_{\phi/\gamma} = 2.2 \cdot 10^6$:

	κ	f/MHz	Q	\mathcal{G}	$P/W(95\% \text{ CL})$	$m_{\gamma'}/\mu\text{eV}$	$\chi(95\% \text{ CL})$
TM ₀₁₀	0.1	207.87961	55405	0.429	$1.08 \cdot 10^{-14}$	0.85972093	$5.4 \cdot 10^{-13}$
TE ₁₁₁	0.01	321.45113	59770	0.674	$1.08 \cdot 10^{-14}$	1.3294150	$8.4 \cdot 10^{-13}$
TE ₁₁₁	0.01	322.74845	58900	0.671	$1.08 \cdot 10^{-14}$	1.3347803	$8.5 \cdot 10^{-13}$
TM ₀₂₀	0.01	454.42411	44340	0.317	$1.08 \cdot 10^{-14}$	1.8793470	$10.1 \cdot 10^{-13}$
TE ₁₁₂	0.01	510.62681	71597	0.020	$1.09 \cdot 10^{-14}$	2.1117827	$28.2 \cdot 10^{-13}$
TE ₁₁₂	0.01	515.97110	67840	0.019	$1.09 \cdot 10^{-14}$	2.1338849	$29.5 \cdot 10^{-13}$
TE ₁₂₀	0.01	577.59175	60350	0.036	$1.10 \cdot 10^{-14}$	2.3887274	$20.4 \cdot 10^{-13}$
TE ₁₂₀	0.01	579.25126	66520	0.037	$1.10 \cdot 10^{-14}$	2.3955906	$19.1 \cdot 10^{-13}$



Phase 1: HP Exclusion Limits

- ❑ Exclusion limits from WISPDMMX Phase 1 measurements: evaluating the broadband signal.
- ❑ Further improvements (factor $\sim 10^2$) will come from stronger amplification, improving the frequency resolution, optimizing the antenna probes and cooling the apparatus.







Phase 2: Tuning the Cavity

- ❑ Tuning plunger assembly: one plunger ready, second being manufactured
- ❑ CST simulations of plunger assembly consisting of two plungers.
- ❑ The assembly should provide effective coverage of up to 56% of the 200-500 MHz range (up 70% with additional vacuum-pump tuning)
- ❑ It will also improve form factors of several modes
- ❑ Optimal antenna location is on the plunger frame




see poster by Le Hoang Nguyen

Phase 2: Testing the Tuning




Phase 2: Experiment Setup

❑ Cavity tuning,
500 MHz bandwidth,
~50 Hz resolution,
automated experim-
ent control




Phase 2: Expected HP Limits

- ❑ WISPDMPX: expected HP dark matter exclusion limits from tuned cavity measurements.






Phase 3: Expected ALP Limits

- ❑ WISPDMPX: expected ALP exclusion limits from measurements with tuned cavity combined with the solenoid magnet from H1 detector (1.15 Tesla)





Broadband: Spherical Reflectors

- ❑ Employing spherical reflectors enhance (focus) the near field EM signal from the reflector surface which arises due to its interaction with WISP dark matter (Horns et al. 2013). Promising for masses above $10 \mu\text{eV}$.
- ❑ Suzuki+ 2015, first results. Pilot study at DESY/Karlsruhe (Döbrich et al.)




Broadband: Radiometry Chambers

- ❑ „Squashing the cauliflower“ and going to $Q=1$ with a detection chamber „coated“ on the inside with fractal antennas.
- ❑ Should get a decent bandpass over a broad range of frequencies.
- ❑ Should get the sensitivity of the total inner surface area by adding (correlating) signals from individual fractal antenna elements.
- ❑ The correlation should also provide full 4π directional sensitivity of measurement.




Radiometry Chambers

- ❑ Time resolution of ~ 3 ns (L_{xyz}/m).
- ❑ Both time and spectral resolution (~ 1 Hz) are achievable with existing radioastronomy detector backends and correlators
- ❑ Coherent addition of signal – effective $Q \sim$ number of detector elements.
- ❑ Coherent addition of signal – full directional sensitivity
- ❑ Possible prototype: cylindrical chamber, with fractal antenna elements at both ends of the cylinder – works best for axions. Try IAXO?



Bright Broadband Future?

- ❑ IAXO (1 bore; 100 days; $T_{\text{sys}} = 4\text{K}$)
Spherical Reflector ($P_{\text{det}} = 10^{-22}\text{ W}$; $R_{\text{det}} = 10^{-2}\text{ Hz}$)



Summary

- ❑ WISP detection relies on low energy experiments; experiments in the radio regime are particularly promising
- ❑ WISPDMPX: First direct WISP dark matter searches in the 0.8-2.0 μeV range: completing measurements at nominal resonances (Phase 1).
- ❑ Next steps:
 - WISPDMPX: Definitive searches for hidden photon (Phase 2) and ALP (Phase 3) dark matter in the 0.8-2.0 μeV range.
 - Further design and implementation of broad-band approaches to WISP searches over the 10^{-2} – 10^{-7} eV mass range.
- ❑ This is an emerging field of study that has a great scientific potential.