

Coannihilating Dark Matter at the LHC

Michael J. Baker

with

Joachim Brod, Sonia El Hedri, Anna Kaminska, Joachim Kopp, Jia Liu,
Andrea Thamm, Maikel de Vries, Xiao-Ping Wang, Felix Yu, José Zurita

arXiv:1510.xxxxx

JGU Mainz

DESY Theory Workshop - 30 September 2015



Outline


- 1 Motivation
- 2 Classification of Simplified Models
- 3 Phenomenology

Outline

- 1 Motivation
- 2 Classification of Simplified Models
- 3 Phenomenology



Dark Matter

Begeman, Broeils & Sanders, 1991




Dark Matter

Planck 2013




Dark Matter



Viel, Becker, Bolton & Haehnelt, 2013


Dark Matter



PDG, 2014

$$\Omega_{\text{nbm}} h^2 = 0.1198 \pm 0.0026$$


Dark Matter





PDG, 2014

$$\Omega_{\text{nbm}} h^2 = 0.1198 \pm 0.0026 \sim \mathcal{O}(\Omega_{\text{bm}} h^2)$$


Theoretical Framework



Theoretical Framework



Theoretical Framework



Simplified Models

- Much recent work on simplified models of DM, e.g.,
 - Profumo *et al.* 1307.6277,
 - De Simone *et al.* 1402.6287,
 - Abdallah *et al.* 1506.03116, ...
- Various tensions, e.g., between relic density and direct/indirect constraints
- Coannihilating models can relieve these tensions

Simplified Models

- Much recent work on simplified models of DM, e.g.,
 - Profumo *et al.* 1307.6277,
 - De Simone *et al.* 1402.6287,
 - Abdallah *et al.* 1506.03116, ...
- Various tensions, e.g., between relic density and direct/indirect constraints
- Coannihilating models can relieve these tensions

Simplified Models

- Much recent work on simplified models of DM, e.g.,
 - Profumo *et al.* 1307.6277,
 - De Simone *et al.* 1402.6287,
 - Abdallah *et al.* 1506.03116, ...
- Various tensions, e.g., between relic density and direct/indirect constraints
- Coannihilating models can relieve these tensions

Our Goal

A complete classification of simplified coannihilation models

Our Goal

A complete classification of simplified coannihilation models

The Coannihilation Codex

Our Goal

A complete classification of simplified coannihilation models

The Coannihilation Codex

This allows us to

- Study connections between experimental probes
- Discuss general phenomenology of models
- Identify lesser studied scenarios
- In the event of a signal, gives a framework for the inverse problem

Outline

- 1 Motivation
- 2 Classification of Simplified Models
- 3 Phenomenology

Assumptions

To complete a classification we need to make some assumptions





- DM is a thermal relic
- DM is a colourless, electrically neutral particle in $(1, N, \beta)$
- Coannihilation diagram is 2-to-2 via dimension four, tree-level couplings
- New particles have spin 0, 1/2 or 1

Assumptions

To complete a classification we need to make some assumptions

- DM is a thermal relic
- DM is a colourless, electrically neutral particle in $(1, N, \beta)$
- Coannihilation diagram is 2-to-2 via dimension four, tree-level couplings
- New particles have spin 0, 1/2 or 1

Coannihilation Diagrams



Classification Procedure

- Work in unbroken $SU(2)_L \times U(1)_Y$
- Given SM field content, iterate over SM_1 and SM_2 to find all possible X using
 - Gauge invariance
 - Lorentz invariance
 - \mathbb{Z}_2 parity (to prevent DM decay)
- Then find all s-channel and t-channel mediators, using same restrictions and
 - Dimension four, tree-level couplings
 - Gauge bosons only couple through kinetic terms

Classification Procedure

- Work in unbroken $SU(2)_L \times U(1)_Y$
- Given SM field content, iterate over SM_1 and SM_2 to find all possible X using
 - Gauge invariance
 - Lorentz invariance
 - \mathbb{Z}_2 parity (to prevent DM decay)
- Then find all s-channel and t-channel mediators, using same restrictions and
 - Dimension four, tree-level couplings
 - Gauge bosons only couple through kinetic terms

Classification Procedure

- Work in unbroken $SU(2)_L \times U(1)_Y$
- Given SM field content, iterate over SM_1 and SM_2 to find all possible X using
 - Gauge invariance
 - Lorentz invariance
 - \mathbb{Z}_2 parity (to prevent DM decay)
- Then find all s-channel and t-channel mediators, using same restrictions and
 - Dimension four, tree-level couplings
 - Gauge bosons only couple through kinetic terms

s-channel classification - sample



DM in $(1, N, \beta)$

ID	X	$\alpha + \beta$	M_s	Spin	$(SM_1 \ SM_2)$	SM_3	M-X-X
ST11	$(3, N \pm 1, \alpha)$	$\frac{7}{3}$	$(3, 2, \frac{7}{3})$	B	$(Q_L \bar{\ell}_R), (u_R \bar{L}_L)$		
ST12				F	$(u_R H)$		
ST13		$\frac{1}{3}$	$(3, 2, \frac{1}{3})$	B	$(d_R \bar{L}_L), (\bar{Q}_L d_R), (u_R L_L)$		
ST14				F	$(u_R H^\dagger), (d_R H)$	Q_L	
ST15		$-\frac{5}{3}$	$(3, 2, -\frac{5}{3})$	B	$(\bar{Q}_L \bar{u}_R), (Q_L \ell_R), (d_R L_L)$		
ST16				F	$(d_R H^\dagger)$		
ST17	$(3, N \pm 2, \alpha)$	$\frac{4}{3}$	$(3, 3, \frac{4}{3})$	B	$(Q_L \bar{L}_R)$		$\checkmark \alpha = -\frac{2}{3}$
ST18				F	$(Q_L H)$		
ST19		$-\frac{2}{3}$	$(3, 3, -\frac{2}{3})$	B	$(\bar{Q}_L \bar{Q}_L), (Q_L L_L)$		$\checkmark \alpha = \frac{1}{3}$
ST20				F	$(Q_L H^\dagger)$		

t-channel classification - sample

DM in $(1, N, \beta)$

ID	X	$\alpha + \beta$	M_t	Spin	$(SM_1 \ SM_2)$	SM_3
TU26	$(1, N \pm 2, \alpha)$	0	$(1, N \pm 1, \beta - 1)$	I	(HH^\dagger)	
TU27			$(1, N \pm 1, \beta + 1)$	II	$(L_L H)$	
TU28			$(1, N \pm 1, \beta - 1)$	III	(HL_L)	
TU29			$(\bar{3}, N \pm 1, \beta - \frac{1}{3})$	IV	$(Q_L \overline{Q_L})$	
TU30			$(1, N \pm 1, \beta + 1)$	IV	$(L_L \overline{L_L})$	
TU31		-2	$(1, N \pm 1, \beta + 1)$	I	$(H^\dagger H^\dagger)$	
TU32			$(1, N \pm 1, \beta + 1)$	II	$(L_L H^\dagger)$	
TU33			$(1, N \pm 1, \beta + 1)$	III	$(H^\dagger L_L)$	



Classification: hybrid models

ID	X	$\alpha + \beta$	SM partner	Extensions
H1	$(1, N, \alpha)$	0	$B, W_i^{N \geq 2}$	SU1, SU3, TU1, TU4–TU8
H2		−2	ℓ_R	SU6, SU8, TU10, TU11
H3	$(1, N \pm 1, \alpha)$	−1	H^\dagger	SU10, TU18–TU23
H4			L_L	SU11, TU16, TU17
H5	$(3, N, \alpha)$	$\frac{4}{3}$	u_R	ST3, ST5, TT3, TT4
H6		$-\frac{2}{3}$	d_R	ST7, ST9, TT10, TT11
H7	$(3, N \pm 1, \alpha)$	$\frac{1}{3}$	Q_L	ST14, TT28–TT31

7 models

Classification: s-channel

ID	X	$\alpha + \beta$	M_s	Spin	$(SM_1 \ SM_2)$	SM_3	M-X-X
SU1	(1, N, α)	0	(1, 1, 0)	B	$\{(u_R \bar{u}_R), (d_R \bar{d}_R), (Q_L \bar{Q}_L), (e_R \bar{e}_R), (L_L \bar{L}_L), (H H^\dagger)\}$	$B, W_i^{N \geq 2}$	✓
SU2				F	$(L_L H)$		
SU3			(1, 3, 0) $^{N \geq 2}$	B	$\{(Q_L \bar{Q}_L), (L_L \bar{L}_L), (H H^\dagger)\}$	B, W_i	✓
SU4				F	$(L_L H)$		
SU5		-2	(1, 1, -2)	B	$(d_R \bar{u}_R), (H^\dagger H^\dagger)$		✓
SU6				F	$(L_L H^\dagger)$	ℓ_R	
SU7			(1, 3, -2) $^{N \geq 2}$	B	$(H^\dagger H^\dagger), (L_L L_L)$	✓ ($\alpha = \pm 1$)	
SU8				F	$(L_L H^\dagger)$	ℓ_R	
SU9	(1, N $\pm 1, \alpha$)	-4	(1, 1, -4)	B	$(e_R \bar{e}_R)$	✓ ($\alpha = \pm 2$)	
SU10		-1	(1, 2, -1)	B	$\{(d_R \bar{Q}_L), (\bar{u}_R \bar{Q}_L), (\bar{L}_L \bar{e}_R)\}$	H^\dagger	
SU11				F	$(\ell_R H)$	L_L	
SU12		-3	(1, 2, -3)	B	$(L_L \ell_R)$		
SU13				F	$(\ell_R H^\dagger)$		
SU14		0	(1, 3, 0)	B	$(L_L \bar{L}_L), (Q_L \bar{Q}_L), (H H^\dagger)$	✓ ($\alpha = 0$)	
SU15				F	$(L_L H)$		
SU16		-2	(1, 3, -2)	B	$(H^\dagger H^\dagger), (L_L L_L)$	✓ ($\alpha = \pm 1$)	
SU17				F	$(L_L H^\dagger)$		

SU type - 17 models

ID	X	$\alpha + \beta$	M_s	Spin	$(SM_1 \ SM_2)$	SM_3	M-X-X
ST1	(3, N, α)	$\frac{10}{3}$	(3, 1, $\frac{10}{3}$)	B	$(u_R \bar{u}_R)$		✓ $\alpha = -\frac{5}{3}$
ST2				B	$(d_R \bar{e}_R), (Q_L \bar{L}_L), (d_R \bar{d}_R)$		✓ $\alpha = -\frac{5}{3}$
ST3			(3, 1, $\frac{4}{3}$)	F	$(Q_L H)$	u_R	
ST4				B	$(Q_L \bar{L}_L)$		✓ $\alpha = -\frac{5}{3}$
ST5		$-\frac{2}{3}$	(3, 3, $\frac{4}{3}$) $^{N \geq 2}$	B	$(Q_L H)$	u_R	
ST6				B	$(Q_L \bar{Q}_L), (\bar{u}_R \bar{d}_R), (\bar{u}_R \bar{e}_R), (Q_L \bar{L}_L)$		✓ $\alpha = \frac{1}{3}$
ST7			(3, 1, $-\frac{8}{3}$)	F	$(Q_L H^\dagger)$	d_R	
ST8				F	$(Q_L \bar{L}_L)$		✓ $\alpha = \frac{1}{3}$
ST9	(3, N $\pm 1, \alpha$)	$\frac{7}{3}$	(3, 3, $-\frac{7}{3}$) $^{N \geq 2}$	F	$(Q_L H^\dagger)$	d_R	
ST10				B	$(\bar{u}_R \bar{u}_R), (d_R \bar{d}_R)$		✓ $\alpha = \frac{1}{3}$
ST11			(3, 2, $\frac{5}{3}$)	B	$(Q_L \bar{e}_R), (\bar{u}_R \bar{L}_L)$		
ST12				F	$(u_R H)$		
ST13		$-\frac{5}{3}$	(3, 2, $\frac{1}{3}$)	B	$(d_R \bar{L}_L), (Q_L \bar{d}_R), (\bar{u}_R \bar{L}_L)$		
ST14				F	$(u_R H^\dagger), (d_R H)$	Q_L	
ST15			(3, 2, $-\frac{5}{3}$)	B	$(Q_L \bar{u}_R), (Q_L \bar{e}_R), (d_R \bar{L}_L)$		
ST16				F	$(d_R H^\dagger)$		
ST17	(3, N $\pm 2, \alpha$)	$\frac{4}{3}$	(3, 3, $\frac{4}{3}$)	B	$(Q_L \bar{L}_R)$		✓ $\alpha = -\frac{2}{3}$
ST18				F	$(Q_L H)$		✓ $\alpha = -\frac{2}{3}$
ST19				B	$(Q_L \bar{Q}_L), (Q_L \bar{L}_L)$		✓ $\alpha = \frac{2}{3}$
ST20		$-\frac{2}{3}$	(3, 3, $-\frac{2}{3}$)	B			
				F	$(Q_L H^\dagger)$		

ST type - 20 models

ID	X	$\alpha + \beta$	M_s	Spin	$(SM_1 \ SM_2)$	SM_3	M-X-X
SO1	(8, N, α)	0	(8, 1, 0) $^{\neq 0 \neq 2}$	B	$(d_R \bar{u}_R), (\bar{u}_R \bar{u}_R), (Q_L \bar{Q}_L)$		✓ $\alpha = 0$
SO2			(8, 3, 0) $^{N \geq 2}$	B	$(Q_L \bar{Q}_L)$		✓ $\alpha = 0$
SO3		-2	(8, 1, -2)	B	$(d_R \bar{u}_R)$		✓ $\alpha = \pm 1$
SO4	(8, N $\pm 1, \alpha$)	-1	(8, 2, -1)	B	$(d_R \bar{Q}_L), (Q_L \bar{u}_R)$		
SE1	(6, N, α)	0	(8, 3, 0)	B	$(Q_L \bar{Q}_L)$		✓ $\alpha = 0$
SE2			(6, 1, $\frac{5}{3}$)	B	$(u_R \bar{u}_R)$		✓ $\alpha = -\frac{2}{3}$
SE3			(6, 1, $\frac{2}{3}$)	B	$(Q_L \bar{Q}_L), (\bar{u}_R \bar{d}_R)$		✓ $\alpha = -\frac{1}{3}$
SE4		$\frac{2}{3}$	(6, 3, $\frac{8}{3}$) $^{N \geq 2}$	B	$(Q_L \bar{Q}_L)$		✓ $\alpha = -\frac{1}{3}$
SE5			(6, 1, $-\frac{4}{3}$)	B	$(d_R \bar{d}_R)$		✓ $\alpha = \frac{2}{3}$
SE6		$\frac{5}{3}$	(6, 2, $\frac{5}{3}$)	B	$(Q_L \bar{u}_R)$		
SE7		$-\frac{1}{3}$	(6, 2, $-\frac{1}{3}$)	B	$(Q_L \bar{d}_R)$		
	(6, N $\pm 2, \alpha$)	$\frac{4}{3}$	(6, 3, $\frac{4}{3}$)	B	$(Q_L \bar{Q}_L)$		✓ $\alpha = -\frac{1}{3}$

SO and SE type - 5 and 7 models

U: X uncoloured

T: X $SU(3)$ triplet

O: X $SU(3)$ octet

E: X $SU(3)$ exotic

Classification: t-channel

ID	X	$\alpha + \beta$	M_t	Spin	(SM1 SM2)	SM3
TU1	(1, N, α)	0	$(1, N \pm 1, \beta - 1)$	I	(HH^\dagger)	$B, W_1^{N \geq 2}$
TU2			$(1, N \pm 1, \beta + 1)$	II	$(L_L H)$	
TU3			$(1, N \pm 1, \beta - 1)$	III	$(H L_L)$	
TU4			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	$B, W_1^{N \geq 2}$
TU5			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(u_R \bar{u}_R)$	$B, W_1^{N \geq 2}$
TU6			$(\bar{3}, N, \beta + \frac{1}{2})$	IV	$(d_R \bar{d}_R)$	$B, W_1^{N \geq 2}$
TU7		-2	$(1, N \pm 1, \beta + 1)$	IV	$(L_L \bar{L}_L)$	$B, W_1^{N \geq 2}$
TU8			$(1, N, \beta + 2)$	IV	$(\ell_R \bar{\ell}_R)$	$B, W_1^{N \geq 2}$
TU9			$(1, N \pm 1, \beta + 1)$	I	$(H^\dagger H^\dagger)$	
TU10			$(1, N \pm 1, \beta + 1)$	II	$(L_L H^\dagger)$	ℓ_R
TU11	(1, N ± 1 , α)	-2	$(1, N \pm 1, \beta + 1)$	III	$(H^\dagger L_L)$	ℓ_R
TU12			$(1, N \pm 1, \beta + 1)$	IV	$(L_L L_L)$	
TU13			$(\bar{3}, N, \beta + \frac{1}{2})$	IV	$(\bar{u}_R d_R)$	
TU14			$(\bar{3}, N, \beta + \frac{1}{2})$	IV	$(d_R \bar{u}_R)$	
TU15		-4	$(1, N, \beta + 2)$	IV	$(\ell_R \bar{\ell}_R)$	
TU16			$(1, N, \beta + 2)$	II	$(\ell_R H)$	L_L
TU17		-1	$(1, N \pm 1, \beta - 1)$	III	$(H H)$	L_L
TU18			$(1, N, \beta + 2)$	IV	$(\ell_R \bar{L}_L)$	H^\dagger
TU19			$(1, N \pm 1, \beta - 1)$	IV	$(L_L L_L)$	H^\dagger
TU20			$(\bar{3}, N, \beta + \frac{1}{2})$	IV	$(d_R \bar{Q}_L)$	H^\dagger
TU21			$(\bar{3}, N \pm 1, \beta + \frac{1}{2})$	IV	$(Q_L d_R)$	H^\dagger
TU22			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{u}_R)$	H^\dagger
TU23			$(\bar{3}, N, \beta + \frac{1}{2})$	IV	$(\bar{u}_R Q_L)$	H^\dagger
TU24		-3	$(1, N \pm 1, \beta + 1)$	IV	$(L_L \ell_R)$	
TU25			$(1, N, \beta + 2)$	IV	$(\ell_R L_L)$	
TU26	(1, N ± 2 , α)	0	$(1, N \pm 1, \beta - 1)$	I	(HH^\dagger)	
TU27			$(1, N \pm 1, \beta + 1)$	II	$(L_L H)$	
TU28			$(1, N \pm 1, \beta - 1)$	III	$(H L_L)$	
TU29			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	
TU30		-2	$(1, N \pm 1, \beta + 1)$	IV	$(L_L L_L)$	
TU31			$(1, N \pm 1, \beta + 1)$	I	$(H^\dagger H^\dagger)$	
TU32			$(1, N \pm 1, \beta + 1)$	II	$(L_L H^\dagger)$	
TU33			$(1, N \pm 1, \beta + 1)$	III	$(H^\dagger L_L)$	

TU type - 33 models

TT type - 52 models

ID	X	$\alpha + \beta$	M_t	Spin	(SM1 SM2)	SM3
TO1	(8, N, α)	0	$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	
TO2			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(u_R \bar{u}_R)$	
TO3			$(\bar{3}, N, \beta + \frac{1}{2})$	IV	$(d_R \bar{d}_R)$	
TO4		-2	$(\bar{3}, N, \beta + \frac{1}{2})$	IV	$(d_R \bar{u}_R)$	
TO5			$(\bar{3}, N, \beta + \frac{1}{2})$	IV	$(\bar{u}_R d_R)$	
TO6	(8, N ± 1 , α)	-1	$(\bar{3}, N, \beta + \frac{1}{2})$	IV	$(d_R \bar{Q}_L)$	
TO7			$(\bar{3}, N \pm 1, \beta + \frac{1}{2})$	IV	$(Q_L d_R)$	
TO8			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{u}_R)$	
TO9			$(\bar{3}, N, \beta + \frac{1}{2})$	IV	$(\bar{u}_R Q_L)$	
TO10			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	
TE1	(6, N, α)	$\frac{4\alpha}{3}$	$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(u_R \bar{u}_R)$	
TE2			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	
TE3			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(u_R d_R)$	
TE4			$(\bar{3}, N, \beta + \frac{1}{2})$	IV	$(d_R u_R)$	
TE5			$(\bar{3}, N, \beta + \frac{1}{2})$	IV	$(d_R d_R)$	
TE6	(6, N ± 1 , α)	$\frac{4\alpha}{3}$	$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(u_R Q_L)$	
TE7			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L u_R)$	
TE8			$(\bar{3}, N, \beta + \frac{1}{2})$	IV	$(d_R Q_L)$	
TE9			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L d_R)$	
TE10			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	

TO and TE type - 10 and 10 models

ID	X	$\alpha + \beta$	M_t	Spin	(SM1 SM2)	SM3
TT1	(3, N, α)	$\frac{4\alpha}{3}$	$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(u_R \bar{u}_R)$	u_R
TT2			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(d_R \bar{d}_R)$	u_R
TT3			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	III	$(H Q_L)$	u_R
TT4			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(d_R \bar{u}_R)$	u_R
TT5			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT6		-1	$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT7			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT8			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT9			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT10			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT11	(3, N, α)	-1	$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(u_R \bar{u}_R)$	u_R
TT12			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(d_R \bar{d}_R)$	u_R
TT13			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	III	$(H Q_L)$	u_R
TT14			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(d_R \bar{u}_R)$	u_R
TT15			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT16		-1	$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT17			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT18			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT19			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT20			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT21	(3, N, α)	$\frac{4\alpha}{3}$	$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(u_R \bar{u}_R)$	u_R
TT22			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(d_R \bar{d}_R)$	u_R
TT23			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	III	$(H Q_L)$	u_R
TT24			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(d_R \bar{u}_R)$	u_R
TT25			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT26		-1	$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT27			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT28			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT29			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT30			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT31	(3, N, α)	$\frac{4\alpha}{3}$	$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(u_R \bar{u}_R)$	u_R
TT32			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(d_R \bar{d}_R)$	u_R
TT33			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	III	$(H Q_L)$	u_R
TT34			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(d_R \bar{u}_R)$	u_R
TT35			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT36		-1	$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT37			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT38			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT39			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT40			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT41	(3, N, α)	$\frac{4\alpha}{3}$	$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(u_R \bar{u}_R)$	u_R
TT42			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(d_R \bar{d}_R)$	u_R
TT43			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	III	$(H Q_L)$	u_R
TT44			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(d_R \bar{u}_R)$	u_R
TT45			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT46		-1	$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT47			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT48			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT49			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT50			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT51	(3, N, α)	$\frac{4\alpha}{3}$	$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(u_R \bar{u}_R)$	u_R
TT52			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(d_R \bar{d}_R)$	u_R
TT53			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	III	$(H Q_L)$	u_R
TT54			$(\bar{3}, N, \beta - \frac{1}{2})$	IV	$(d_R \bar{u}_R)$	u_R
TT55			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT56		-1	$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT57			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT58			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT59			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R
TT60			$(\bar{3}, N \pm 1, \beta - \frac{1}{2})$	IV	$(Q_L \bar{Q}_L)$	u_R







Complete Classification

Given our assumptions, one of these simplified models of coannihilating dark matter is the one chosen by Nature!







Outline

- 1 Motivation
- 2 Classification of Simplified Models
- 3 Phenomenology**




Production: s-channel





Production: s-channel






Decay: s-channel



Decay: s-channel



Decay: s-channel



Generic Signatures: s-channel

- **Mono-Y (Y=jet, photon, Z,...) + \cancel{E}_T** from DM DM, XX,...
 - classic signature
- **Single and Double Resonances** from M and MM
 - ATLAS/CMS Exotics
- **Mono-Y + \cancel{E}_T + soft** from XX,MM,...
 - has been motivated, no searches yet
- **Resonance + \cancel{E}_T + soft** from MM
 - new signature to explore!

Generic Signatures: s-channel

- **Mono-Y (Y=jet, photon, Z,...) + \cancel{E}_T** from DM DM, XX,...
 - classic signature
- **Single and Double Resonances** from M and MM
 - ATLAS/CMS Exotics
- **Mono-Y + \cancel{E}_T + soft** from XX,MM,...
 - has been motivated, no searches yet
- **Resonance + \cancel{E}_T + soft** from MM
 - new signature to explore!

Generic Signatures: s-channel

- **Mono-Y** ($Y=\text{jet, photon, } Z, \dots$) + \cancel{E}_T from DM DM, XX, ...
 - classic signature
- **Single and Double Resonances** from M and MM
 - ATLAS/CMS Exotics
- **Mono-Y + \cancel{E}_T + soft** from XX, MM, ...
 - has been motivated, no searches yet
- **Resonance + \cancel{E}_T + soft** from MM
 - new signature to explore!

Generic Signatures: s-channel

- **Mono-Y (Y=jet, photon, Z,...) + \cancel{E}_T** from DM DM, XX,...
 - classic signature
- **Single and Double Resonances** from M and MM
 - ATLAS/CMS Exotics
- **Mono-Y + \cancel{E}_T + soft** from XX,MM,...
 - has been motivated, no searches yet
- **Resonance + \cancel{E}_T + soft** from MM
 - new signature to explore!

Example - ST11

ID	X	$\alpha + \beta$	M_s	Spin	(SM ₁ SM ₂)	SM ₃	M-X-X
ST11	$(3, N \pm 1, \alpha)$	$\frac{7}{3}$	$(3, 2, \frac{7}{3})$	B	$(Q_L \overline{\ell_R}), (u_R \overline{L_L})$		

DM in $(1, N, \beta)$

Field	Rep.	Spin and mass assignment
DM	$(1, 1, 0)$	Majorana fermion
X	$(3, 2, 7/3)$	Dirac fermion
M	$(3, 2, 7/3)$	Scalar

Example - ST11


ID	X	$\alpha + \beta$	M_s	Spin	$(SM_1 \ SM_2)$	SM_3	M-X-X
ST11	$(3, N \pm 1, \alpha)$	$\frac{7}{3}$	$(3, 2, \frac{7}{3})$	B	$(Q_L \overline{\ell_R}), (u_R \overline{L_L})$		

DM in $(1, N, \beta)$

Field	Rep.	Spin and mass assignment
DM	$(1, 1, 0)$	Majorana fermion
X	$(3, 2, 7/3)$	Dirac fermion
M	$(3, 2, 7/3)$	Scalar

Example - ST11


Field	Rep.	Spin and mass assignment
DM	(1,1,0)	Majorana fermion
X	(3,2,7/3)	Dirac fermion
M	(3,2,7/3)	Scalar



$$\mathcal{L} \supset \mathcal{L}_{\text{kin}} + y_D \bar{X} M DM + y_{Q\ell} \overline{Q}_L M \ell_R + y_{Lu} \overline{L}_L M^c u_R + h.c.$$

Example - ST11


Field	Rep.	Spin and mass assignment
DM	(1,1,0)	Majorana fermion
X	(3,2,7/3)	Dirac fermion
M	(3,2,7/3)	Scalar



$$\mathcal{L} \supset \mathcal{L}_{\text{kin}} + y_D \bar{X} M DM + y_{Q\ell} \overline{Q_L} M \ell_R + y_{Lu} \overline{L_L} M^c u_R + h.c.$$


Example - ST11

Field	Rep.	Spin and mass assignment
DM	(1,1,0)	Majorana fermion
X	(3,2,7/3)	Dirac fermion
M	(3,2,7/3)	Scalar




$$\mathcal{L} \supset \mathcal{L}_{\text{kin}} + y_D \bar{X} M DM + y_{Q\ell} \overline{Q}_L M \ell_R + y_{Lu} \overline{L}_L M^c u_R + h.c.$$

Example - ST11




$$\mathcal{L} \supset \mathcal{L}_{\text{kin}} + y_D \bar{X} M DM + y_{Q\ell} \bar{Q}_L M \ell_R + y_{Lu} \bar{L}_L M^c u_R + h.c.$$




- Strong production
- Lepton-jet resonance + \cancel{E}_T + soft lepton & jet
- No dedicated LHC search

Example - ST11




$$\mathcal{L} \supset \mathcal{L}_{\text{kin}} + y_D \bar{X} M DM$$

$$+ y_{Q\ell} \bar{Q}_L M \ell_R + y_{Lu} \bar{L}_L M^c u_R + h.c.$$




- Strong production
- Lepton-jet resonance + \cancel{E}_T + soft lepton & jet
- No dedicated LHC search

Example - ST11



$$\mathcal{L} \supset \mathcal{L}_{\text{kin}} + y_D \bar{X} M DM$$

$$+ y_{Q\ell} \bar{Q}_L M \ell_R + y_{Lu} \bar{L}_L M^c u_R + h.c.$$



- Strong production
- Lepton-jet resonance + \cancel{E}_T + soft lepton & jet
- No dedicated LHC search

Summary

- Given assumptions, the Coannihilation Codex contains the real model of Nature!
- Guaranteed kinetic & coannihilation vertices → signatures
- Classify general signatures
 - Identify new signatures
 - Identify interesting models, e.g., leptoquarks and DM
- Huge number of DM models
 - collider signatures
 - direct and indirect detection
 - flavour bounds
 - cosmology
 - ...

Summary

- Given assumptions, the Coannihilation Codex contains the real model of Nature!
- Guaranteed kinetic & coannihilation vertices \rightarrow signatures
- Classify general signatures
 - Identify new signatures
 - Identify interesting models, e.g., leptoquarks and DM
- Huge number of DM models
 - collider signatures
 - direct and indirect detection
 - flavour bounds
 - cosmology
 - ...

Summary

- Given assumptions, the Coannihilation Codex contains the real model of Nature!
- Guaranteed kinetic & coannihilation vertices \rightarrow signatures
- Classify general signatures
 - Identify new signatures
 - Identify interesting models, e.g., leptoquarks and DM
- Huge number of DM models
 - collider signatures
 - direct and indirect detection
 - flavour bounds
 - cosmology
 - ...

Summary

- Given assumptions, the Coannihilation Codex contains the real model of Nature!
- Guaranteed kinetic & coannihilation vertices \rightarrow signatures
- Classify general signatures
 - Identify new signatures
 - Identify interesting models, e.g., leptoquarks and DM
- Huge number of DM models
 - collider signatures
 - direct and indirect detection
 - flavour bounds
 - cosmology
 - ...

Summary

- Given assumptions, the Coannihilation Codex contains the real model of Nature!
- Guaranteed kinetic & coannihilation vertices \rightarrow signatures
- Classify general signatures
 - Identify new signatures
 - Identify interesting models, e.g., leptoquarks and DM
- Huge number of DM models
 - collider signatures
 - direct and indirect detection
 - flavour bounds
 - cosmology
 - ...