

Determinations of α_s from $e^+e^- \rightarrow \text{Hadrons}$

J. H. Kühn

(KIT, Karlsruhe)

- 1) Status: BESS (BaBar, Belle)
- 2) Status and Perspectives for $e^+e^- \rightarrow \text{hadrons at } Z$
- 3) M_W from G_F, M_Z, α and the rest
- 4) Perspectives for $e^+e^- \rightarrow \text{hadrons above } Z$
- 5) Perspectives for $e^+e^- \rightarrow Z + H (\rightarrow \text{hadrons})$

1) Status: BESS (BaBar, Belle)

e^+e^- at low energies

BESS (PLB 641 (2006) 145)

$$R(3.650 \text{ GeV and } 3.6648 \text{ GeV}) = 2.224 \pm 0.019 \pm 0.089$$

$$R = 3 \left(Q_u^2 + Q_d^2 + Q_s^2 \right) \left(1 + a_s + 1.40923a_s^2 - 12.7671a_s^3 - 79.9806a_s^4 \right) \\ + 3 \underbrace{(Q_u + Q_d + Q_c)^2}_{=0} \left(-0.4138a_s^3 - 4.9841a_s^4 \right)$$

present experimental precision at BESS: $\frac{\delta R}{R} \approx 4\%$

$$\Rightarrow \alpha_s \approx 0.31^{+0.13}_{-0.14}$$

$$\Rightarrow \frac{\delta R}{R} \approx 2\% \quad \text{looks feasible}$$

BaBar

Belle





?

2) Status and Perspectives for $e^+e^- \rightarrow \text{Hadrons at the } Z$

Γ_{had} and $\Gamma_{had}/\Gamma_{lept}$ corrections known to $O(\alpha_s^4)$, N³LO

(Baikov, Chetyrkin, JK, Rittinger, arxiv: 0801.1821, 1201.5804)

non-singlet & singlet, vector & axial correlators



- theory uncertainty from $M_Z/3 < \mu < 3M_Z$

$$\left. \begin{aligned} \Rightarrow \delta\Gamma_{NS} &= 101\text{keV}; \\ \delta\Gamma_S^V &= 2.7\text{keV}; \\ \delta\Gamma_S^A &= 42\text{keV}; \end{aligned} \right\} \begin{aligned} \Sigma &= 145.7\text{keV} \\ &(\text{corresponds to } \delta\alpha_s \sim 3 \times 10^{-4}) \end{aligned}$$

TLEP: $\delta\Gamma_{had} \hat{=} 100 \text{ keV}$

- similar analysis of $\Gamma(W \rightarrow \text{had})$ only affected by non-singlet contributions!

- b-mass corrections under control: $m_b^2\alpha_s^4; m_b^4\alpha_s^3; \dots$


- one more loop?

$$\alpha_s^2(1979), \alpha_s^3(1991), \alpha_s^4(2008), \alpha_s^5(?),$$

guesses on α_s^5 based on

Mixed electroweak and QCD: light quarks (u,d,c,s)

terms of $O(\alpha\alpha_s)$, Czarnecki, JK; hep-ph/9608366



$$\Delta\Gamma \equiv \Gamma(\text{two loop (EW} \star \text{QCD)}) - \Gamma_{\text{Born}} \delta_{\text{EW}}^{\text{NLO}} \delta_{\text{QCD}}^{\text{NLO}} = -0.59(3) \text{ MeV}$$

aim (T-LEP paper): $\delta\Gamma \approx 0.1 \text{ MeV}$

three loop: reduction by $\# \cdot \frac{\alpha_s}{\pi} = \#0.04$

should not exceed 5!

corrections of $O(\alpha_w \alpha_s^2)$ (three loop) difficult

$$\Gamma(Z \rightarrow b\bar{b}) \equiv \Gamma_b$$

aim: $\delta R_b \equiv \frac{\delta\Gamma_b}{\Gamma_Z} = 2 - 5 \times 10^{-5}$ (LEP: $R_b = 0.21629 \pm 0.00066$, corresponds to $\delta\Gamma_b \approx 1.6$ MeV)

2×10^{-5} corresponds to 0.05 MeV!

corrections specific for $b\bar{b}$:

m_t^2 -enhancement: order $G_F m_t^2$ and $G_F m_t^2 \alpha_s$

$$\Delta\Gamma = \frac{G_F M^3}{16\pi^3} G_f m_t^2 \left(1 - \frac{2}{3} s_w^2\right) \left(1 - \frac{\pi^2 - 3}{3} \frac{\alpha_s}{\pi}\right) \quad (\text{Fleischer et al 1992})$$


Complete $\alpha_w \alpha_s$ result:

$$\Gamma_b - \Gamma_q = \left(-5.69 - 0.79 \quad O(\alpha) \right. \\ \left. + 0.50 + 0.06 \quad O(\alpha\alpha_s) \right) \text{ MeV}$$

separated into m_t^2 -enhanced and rest

(Harlander, Seidensticker, Steinhauser

hep-ph/9712228)



dressed with gluons

motivates the evaluation of m_t^2 -enhanced corrections of $O(G_F m_t^2 \alpha_s^2)$

(Chetyrkin, Steinhauser, hep-ph/990480)

$$\delta\Gamma_b(G_F m_t^2 \alpha_s^2) \approx 0.1 \text{ MeV} \quad (\text{non-singlet})$$

(absent in Z-fitter, G-fitter!)

General observation:

many top-induced corrections become significantly smaller, if m_t is expressed in \overline{MS} convention

$$\bar{m}_t(\bar{m}_t) = m_{pole} \left(1 - 1.33 \left(\frac{\alpha_s}{\pi} \right) - 6.46 \left(\frac{\alpha_s}{\pi} \right)^2 - 60.27 \left(\frac{\alpha_s}{\pi} \right)^3 - 704.28 \left(\frac{\alpha_s}{\pi} \right)^4 \right)$$

(Karlsruhe, 1999) (Marquard, Smirnov, Smirnov, Steinhauser, 2015)

$$= (173.34 - 7.96 - 1.33 - 0.43 - 0.17) \text{ GeV}$$

$$= \left(163.45 \pm 0.72|_{m_t} \pm 0.19|_{\alpha_s} \pm ?|_{th} \right) \text{ GeV}$$

top scan $\Rightarrow m(\text{potential subtracted})$

$$\delta m_t \sim 20 - 30 \text{ MeV}$$


$$\Gamma(Z \rightarrow b\bar{b})$$

Can we isolate the $Zb\bar{b}$ -vertex?

$$R_b = 0.21629 \pm 0.00066 \text{ (LEP)}; \quad 3\% \hat{=} \Gamma(Z \rightarrow b\bar{b})/\Gamma_{had} \equiv 1.2 \text{ MeV}$$

$$\text{TLEP: } \delta R = 2 - 5 \times 10^{-5} \hat{=} 50 - 120 \text{ keV}$$

conceptual problem: singlet-terms



mixed contributions, "singlet"

$$\Gamma_{b\bar{b}c\bar{c}}^{\text{singlet}} = \left(\frac{G_F M_Z^3}{8\sqrt{2}\pi} \right) 0.31 \left(\frac{\alpha_s}{\pi} \right)^2 \approx 340 \text{ keV}$$

(total hadronic rate more robust!)

3) M_W from G_F , M_Z , α and the rest

LEP: $\delta M_W \simeq 30$ MeV; LEP+Tevatron: $\delta M_W \simeq 15$ MeV; Fcc-ee: $\delta M_W \simeq 0.5 - 1$ MeV

Theory

$$M_W^2 = f(G_F, M_Z, m_t, \Delta\alpha, \dots) = \frac{M_Z^2}{2(1-\delta\rho)} \left(1 + \sqrt{1 - \frac{4\pi\alpha(1-\delta\rho)}{\sqrt{2}G_F M_Z^2} \left(\frac{1}{1-\Delta\alpha} + \dots \right)} \right);$$

m_t -dependence through $\delta\rho_t$

$$\delta M_W \approx M_W \frac{1}{2} \frac{\cos^2 \theta_w}{\cos^2 \theta_w - \sin^2 \theta_w} \delta\rho \approx 5.7 \times 10^4 \delta\rho \text{ [MeV]}$$

$$\delta\rho_t = 3X_t \left(1 - 2.8599 \left(\frac{\alpha_s}{\pi} \right) - 14.594 \left(\frac{\alpha_s}{\pi} \right)^2 - 93.1 \left(\frac{\alpha_s}{\pi} \right)^3 \right)$$

↓


$$\delta M_W = 9.5 \text{ MeV}$$

↓

$$\delta M_W = 2.1 \text{ MeV}$$

α_s^3 : 4 loop (Chetyrkin, JK, Maierhöfer, Sturm; Boughezal, Czakon, 2006)

mixed QCD \star electroweak



three loop

$(X_t \equiv G_F m_t^2)$

- X_t^3 (purely weak) $\Rightarrow 200\text{eV}$
- $\alpha_s X_t^2$ (mixed) $\Rightarrow 2.5\text{MeV}$
- $\alpha_s^2 X_t$ (QCD three loop) $\Rightarrow -9.5\text{MeV}$
- $\alpha_s^3 X_t$ (QCD four loop) $\Rightarrow 2.1\text{MeV}$

the future

individual uncalculated higher orders below 0.5 MeV, examples:

$\alpha_s^2 X_t^2$ presumably feasible (4 loop tadpoles), $\alpha_s^4 X_t$ 5 loop tadpoles?

dominant contribution from $m_t(\text{pole}) \Rightarrow \bar{m}_t$

crucial input: m_t also for stability of the universe

$$\delta M_W \approx 6 \times 10^{-3} \delta m_t$$


$$\delta m_t = 1 \text{ GeV}$$

$$\Rightarrow \delta M_W \approx 6 \text{ MeV (status)}$$

conversely:

$$\text{TLEP: } \delta M_W = 0.5 \text{ MeV}$$

$$\text{requires } \delta m_t = 100 \text{ MeV}$$





(Zoller)

TLEP: $\delta m_t = 10 - 20$ MeV

based on **bold** extrapolation of ILC study (ILC: 35 MeV, no theory error)

momentum distribution etc: LO only

σ_{tot} in N³LO just completed (Beneke, Kiyo Marquard, Piclum, Penin, Steinhauser)



robust location of threshold, extraction of λ_{Yuk} requires normalization!

important ingredient: $\bar{m}_t(\bar{m}_t) \Leftrightarrow m_{pole}$

example: $m_{pole} = 173.340 \pm 0.87 \text{ GeV}$, $\alpha_s \equiv \alpha_s^{(6)}(m_t) = 0.1088$

4 loop term is just completed ([Marquard, Smirnov, Smirnov, Steinhauser, 2015](#))


$$\begin{aligned} m_{pole} &= \bar{m}_t(\bar{m}_t) \left(1 + 0.4244 \alpha_s + 0.8345 \alpha_s^2 + 2.365 \alpha_s^3 + (8.49 \pm 0.25) \alpha_s^4 \right) \\ &= (163.643 + 7.557 + 1.617 + 0.501 + 0.195 \pm 0.05) \text{ GeV} \end{aligned}$$

four-loop term matters!

4) Perspectives for $e^+e^- \rightarrow \text{Hadrons above } Z$


pure QED

\Rightarrow absence of




(Yang Theorem)

corrections of $O(\alpha^2)$!




electroweak theory



$$\sum_q \left((g_A^e)^2 g_V^e \right) \left((g_A^q)^2 g_V^q \right) = \left((g_A^e)^2 g_V^e \right) \left((g_A^u)^2 \sum_q g_V^q \right) \neq 0$$


leading contribution $O(\alpha)$




$$\sum_q \left((g_A^q)^2 g_V^q \right) = \left((g_A^u)^2 \sum_q g_V^q \right) \neq 0$$

\Rightarrow interference of order α

similarly for




Important radiative tail from Z



precise predictions ($< \frac{1}{2}\%$) difficult:

large radiative tail from the Z (factor 3 compared to Born cross section)

5) Perspectives for $e^+e^- \rightarrow Z + H(\rightarrow \text{hadrons})$



Cross sections for the three major Higgs production processes as a function of center of mass energy (from arXiv:1306.6352)

example: $H \rightarrow b\bar{b}$ dominant decay mode, all branching ratios are affected!

TLEP: $\sigma_{HZ} \times Br(H \rightarrow b\bar{b})$: aim 0.2%

Higgs WG, arXiv:1307.1347 (Table 1) assumes $\alpha_s = 0.119 \pm 0.002$, $m_b|_{pole} = 4.49 \pm 0.06$ GeV:

$$\frac{\delta\Gamma(H \rightarrow b\bar{b})}{\Gamma(H \rightarrow b\bar{b})} = \pm 2.3\%|_{\alpha_s} \pm 3.2\%|_{m_b} \pm 2.0\%|_{th} \Rightarrow 7.5\%$$

Our estimate: $\Gamma(H \rightarrow b\bar{b}) = \frac{G_F M_H}{4\sqrt{2}\pi} m_b^2(M_H) R^S(s = M_H^2, \mu^2 = M_H^2)$

$$\begin{aligned} R^S(M_H) &= 1 + 5.667 \left(\frac{\alpha_s}{\pi}\right) + 29.147 \left(\frac{\alpha_s}{\pi}\right)^2 + 41.758 \left(\frac{\alpha_s}{\pi}\right)^3 - 825.7 \left(\frac{\alpha_s}{\pi}\right)^4 \\ &= 1 + 0.1948 + 0.03444 + 0.0017 - 0.0012 \\ &= 1.2298 \quad (\text{Chetyrkin, Baikov, JK, 2006}) \end{aligned}$$

for $\alpha_s(M_Z) = 0.118$, $\alpha_s(M_H) = 0.108$

Theory uncertainty ($M_H/3 < \mu < 3M_H$): 5‰ (four loop) reduced to 1.5‰ (five loop)

present parametric uncertainties:

$$m_b(10\text{GeV}) = 3610 - \frac{\alpha_s - 0.1189}{0.002} 12 \pm 11 \text{ MeV (Karlsruhe, arXiv:0907.2110)}$$

$$\left(\begin{array}{l} \text{Bodenstein+Dominguez: } 3623(9) \text{ MeV} \\ \text{HPQCD} \quad \quad \quad 3617(25) \text{ MeV} \end{array} \right)$$

(α_s uncertainties are presently dominant, assuming $\delta = 0.002$, they influence m_b -determination; running to M_H ; R^S)

running from 10 GeV to M_H depends on

anomalous mass dimension, β -function and α_s

$$m_b(M_H) = 2759 \pm 8 |_{m_b} \pm 27 |_{\alpha_s} \text{ MeV}$$

γ_4 (five loop): Baikov, Chetyrkin, J.H.K., 2012

β_4 under construction

$$\frac{\delta m_b^2(M_H)}{m_b^2(M_H)} = -1.4 \times 10^{-4} \left(\frac{\beta_4}{\beta_0} = 0 \right) \quad | \quad -4.3 \times 10^{-4} \left(\frac{\beta_4}{\beta_0} = 100 \right) \quad | \quad -7.3 \times 10^{-4} \left(\frac{\beta_4}{\beta_0} = 200 \right)$$

to be compared with $\delta\Gamma(H \rightarrow b\bar{b})/\Gamma(H \rightarrow b\bar{b}) = 2.0 \times 10^{-4}$ (FCC-ee)

with the just computed β_4 (Baikov, Chetyrkin, J.H.K, 2016)

$$\frac{\beta_4}{\beta_0} = 7.882 \text{ for } n_f = 5$$

and, as a result, the shift induced by the five loop term in $\beta(\alpha_s)$ amounts to

$$\frac{\delta m_b^2(M_H)}{m_b^2(M_H)} = -0.24 \times 10^{-4}$$

to be compared with $\delta\Gamma(H \rightarrow b\bar{b})/\Gamma(H \rightarrow b\bar{b}) = 2.0 \times 10^{-4}$ (FCC-ee)

perspectives: (assume $\delta\alpha_s = \delta\alpha_s(\text{now})/10 = 2 \times 10^{-4}$)

$\delta m_b(10\text{GeV})/m_b \sim 10^{-3}$ conceivable (dominated by $\delta\Gamma(\Upsilon \rightarrow e^+e^-)$)

$$\Rightarrow \frac{\delta\Gamma_{H \rightarrow b\bar{b}}}{\Gamma_{H \rightarrow b\bar{b}}} = \pm 2 \times 10^{-3} |_{m_b} \pm 1.3 \times 10^{-3} |_{\alpha_s, \text{running}} \pm 1 \times 10^{-3} |_{\text{theory}}$$

similarly: $\Gamma_{H \rightarrow c\bar{c}}$

$$\begin{aligned} \delta m_c(3 \text{ GeV})/m_c(3 \text{ GeV}) &= 13 \text{ MeV}/986 \text{ MeV} && (\text{now}) \\ &= 5 \text{ MeV}/986 \text{ MeV} && (\text{conceivable}) \end{aligned}$$

$$\begin{aligned} m_c(M_H) &= (609 \pm 8 |_{m_c} \pm 9 |_{\alpha_s}) \text{ MeV} && (\text{now}) \\ &\pm 3 \text{ MeV} && (\text{conceivable}) \end{aligned}$$

$$\begin{aligned} \Rightarrow \frac{\delta\Gamma_{H \rightarrow c\bar{c}}}{\Gamma_{H \rightarrow c\bar{c}}} &= \pm 5.5 \times 10^{-2} && (\text{now}) \\ &= \pm 1 \times 10^{-2} && (\text{conceivable}) \end{aligned}$$

Starting from order α_s^3 the separation of $H \rightarrow gg$ and $H \rightarrow b\bar{b}$

is no longer unambiguously possible. (Chetyrkin, Steinhauser, 1997)

$H \rightarrow gg$ to $O(\alpha_s^5)$ (hep-ph/0604194; Baikov, Chetyrkin)

(separation of gg , $b\bar{b}$, $c\bar{c}$ difficult in $O(\alpha_s^4)$ and higher)

$$\Gamma(H \rightarrow gg) = K \cdot \Gamma_{\text{Born}}(H \rightarrow gg)$$

and

$$K = 1 + 17.9167 a'_s + (156.81 - 5.7083 \ln \frac{M_t^2}{M_H^2}) (a'_s)^2 + (467.68 - 122.44 \ln \frac{M_t^2}{M_H^2} + 10.94 \ln^2 \frac{M_t^2}{M_H^2}) (a'_s)^3.$$

take $M_t = 175$ GeV, $M_H = 120$ GeV and $a'_s = \alpha_s^{(5)}(M_H)/\pi = 0.0363$:


$$\begin{aligned} K &= 1 + 17.9167 a'_s + 152.5 (a'_s)^2 + 381.5 (a'_s)^3 \\ &= 1 + 0.65038 + 0.20095 + 0.01825. \end{aligned}$$

Claim: experimental precision of $\sigma(HZ)$ BR ($H \rightarrow gg$) = 1.4%

~ approximately equal to last calculated correction

$$H \rightarrow \gamma\gamma$$

(arxiv:1212.6233; Maierhöfer, Marquard)



non-singlet and singlet terms; electroweak corrections (Passarino,...)

$$\Gamma_{H \rightarrow \gamma\gamma} = (9.398 - \frac{0.148}{\text{LO} \times \text{NLO-EW}} + \frac{0.168}{\text{LO} \times \text{NLO-QCD}} + 0.00793 \alpha_s^2) \text{ keV}$$

α_s^2 term dominated by singlet part of prediction,

prediction good to $O(1)$ permille!

SUMMARY

- Improved experimental precision at low energies (BESS, BELLE) would lead to precise value of α_s
- Z decays at LEP (and even more so at a future Fcc-ee or linear collider) gives a precise value of α_s . Present result: $\alpha_s = 0.1197 \pm 0.0028$
- Important implications for prediction of $M_W^2 = f(G_F, M_Z, \alpha_s, \dots)$
- Large radiative corrections for $e^+e^- \rightarrow$ hadrons above the Z;
precise predictions are not yet established
- m_b -determination from sum rules at low energies; impact on $\Gamma(H \rightarrow b\bar{b})!$
- attractive perspectives for $e^+e^- \rightarrow Z + H$, in particular for $\Gamma(H \rightarrow$ hadrons)