

# XXIV International Workshop on Deep-Inelastic Scattering and Related Subjects (DIS16)



Contribution ID: 29

Type: **not specified**

## Search for QCD Instanton-Induced Processes in DIS at HERA

*Tuesday, April 12, 2016 2:25 PM (20 minutes)*

Signals of QCD instanton-induced processes are searched for in deep-inelastic scattering (DIS) at the electron-proton collider HERA in the kinematic region defined by the Bjorken-scaling variable  $x > 0.001$ , the inelasticity  $0.2 < y < 0.7$  and the photon virtuality  $150 < Q^2 < 15000 \text{ GeV}^2$ . The search is performed using H1 data corresponding to an integrated luminosity of  $350 \text{ pb}^{-1}$ . Several observables of the hadronic final state of the events are exploited to identify a potentially instanton-enriched domain. Two Monte Carlo models, RAPGAP and ARIADNE, are used to estimate the background from the standard DIS processes, and the instanton-induced scattering processes are modeled by the program QCDINS. In order to extract the expected signal a multivariate data analysis technique is used. Exclusion limits on instanton production are reported, excluding cross sections larger than  $2 \text{ pb}$ . Limits are also reported as a function of parameters used to regularize the perturbative instanton model.

**Primary author:** Dr H1, Collaboration (DESY)

**Presenter:** MIKOCKI, Stanislaw (Institute of Nuclear Physics Polish Academy of Sciences)

**Session Classification:** WG2 QCD and Hadronic Final States

**Track Classification:** QCD and Hadronic Final States