



Contribution ID: 178

Type: not specified

New single- and double-parton scattering mechanisms for double charmed meson production

Tuesday, April 12, 2016 9:20 AM (15 minutes)

Some time ago two of us predicted that at large energies relevant for the LHC the production of double charm should be dominated by the double-parton scattering (DPS) mechanism [1]. Those studies of double $c\bar{c}$ production was extended next to the k_t -factorization approach which includes effectively higher-order QCD effects [2,3]. A relatively good description of the LHCb experimental data [4] was achieved for both the total yield and the dimeson correlation observables. The single-parton scattering (SPS) $gg \rightarrow c\bar{c}c\bar{c}$ contribution was discussed carefully in both collinear [3] and k_t -factorization [5] approaches. Their contribution to the $c\bar{c}c\bar{c}$ cross section was found to be rather small and was not able to describe details of the LHCb data.

Here we discuss production of $D^0 D^0$ (and $\bar{D}^0 \bar{D}^0$) pairs within an alternative approach where $g \rightarrow D$ fragmentation is included [6]. We consider double-parton scattering (DPS) mechanisms of double $c\bar{c}$ production and subsequent $cc \rightarrow D^0 D^0$ hadronization as well as double g and mixed $gc\bar{c}$ production with $gg \rightarrow D^0 D^0$ and $gc \rightarrow D^0 D^0$ hadronization calculated with the help of the scale dependent hadronization functions of Kniehl et al. Single-parton scattering (SPS) mechanism of digluon production is also taken into account. We compare our results with several correlation observables in azimuthal angle $\varphi_{D^0 D^0}$ between D^0 mesons or in dimeson invariant mass $M_{D^0 D^0}$. The inclusion of new mechanisms with $g \rightarrow D^0$ fragmentation leads to larger cross sections, than when including only DPS mechanism with standard scale independent $cc \rightarrow D^0 D^0$ fragmentation functions. Some consequences of the presence of the new mechanisms are discussed. In particular a larger σ_{eff} is needed to describe the LHCb data. There is a signature that σ_{eff} may depend on transverse momentum of c quarks and/or \bar{c} antiquarks.

[1] M. Luszczak, R. Maciula and A. Szczurek, Production of two $c\bar{c}$ pairs in double-parton scattering, Phys. Rev. D 85, 094034 (2012).

[2] R. Maciula and A. Szczurek, Production of $c\bar{c}c\bar{c}$ in double-parton scattering within k_t -factorization approach - meson-meson correlations, Phys. Rev. D 87, 074039 (2013).

[3] A. van Hameren, R. Maciula and A. Szczurek, Single-parton scattering versus double-parton scattering in the production of two $c\bar{c}$ pairs and charmed meson correlations at the LHC, Phys. Rev. D 89, 094019 (2014).

[4] R. Aaij et al. [LHCb Collaboration], Observation of double charm production involving open charm in pp collisions at $\sqrt{s} = 7$ TeV, J. High Energy Phys. 06, 141 (2012); [J. High Energy Phys. 03, 108 (2014)].

[5] A. van Hameren, R. Maciula and A. Szczurek, Production of two charm quark-antiquark pairs in single-parton scattering within the k_t -factorization approach, Phys. Lett. B 748, 167 (2015).

[6] R. Maciula and V. A. Saleev, A. V. Shipilova and A. Szczurek, arXiv:1601.06981 [hep-ph].

Primary authors: Prof. SZCZUREK, Antoni (the Institute of Nuclear Physics PAS); Dr MACIULA, Rafal (the Institute of Nuclear Physics PAS); Dr ALEXANDRA, Shipilova (Samara State Aerospace University); Prof. SALEEV, Vladimir (Samara State Aerospace University)

Presenter: Prof. SZCZUREK, Antoni (the Institute of Nuclear Physics PAS)

Session Classification: WG4 Heavy Flavours

Track Classification: Heavy Flavours (Charm, Beauty and Top)