

XXIV International Workshop on Deep-Inelastic Scattering and Related
Subjects (DIS16)



Contribution ID: 189

Type: **not specified**

Comparison of the F2 Structure Function in Iron as Measured by Charged Lepton and Neutrino Probes

Wednesday, April 13, 2016 11:20 AM (15 minutes)

World data for the F2 structure function for Iron, as measured by multiple charged lepton and neutrino scattering experiments, are compared. Data obtained from charged lepton and neutrino scattering at larger values of x are in remarkably good agreement with a simple invocation of the 18/5 rule, while a discrepancy in the behavior of the data obtained from the different probes well beyond the data uncertainties is observed in the shadowing/anti-shadowing transition region where the Bjorken scaling variable x is less than 0.15. The data are compared to theoretical calculations. Details and results of the data comparison will be presented.

Primary author: Dr CYNTHIA, Keppel (Thomas Jefferson National Accelerator Facility)

Co-authors: Prof. CHRISTY, M. Eric (Hampton University); Dr KALANTARIANS, Narbe (Hampton University)

Presenter: Dr CYNTHIA, Keppel (Thomas Jefferson National Accelerator Facility)

Session Classification: WG1 Structure Functions and Parton Densities

Track Classification: Structure Functions and Parton Densities