


# Searches for Dark Matter and Axion-Like Particles in Active Galactic Nuclei

Joseph Conlon  
University of Oxford

New Directions in String Phenomenology

DESY, February 15<sup>th</sup> 2017





Based on

1605.01043


Marcus Berg, JC, Francesca Day, Nick Jennings, Sven Krippendorf, Markus Rummel

1608.01684

JC, Francesca Day, Nick Jennings, Sven Krippendorf, Markus Rummel



# I. The Targets



# Axion-Like Particles in String Theory

- 30-year old result:

String compactifications lead to a plenitude of axions in the low-energy theory

- 'Model-dependent' axions number  $O(100)$  for typical compactifications
- Axion-like particles are one of **the most motivated targets** in looking for signatures of string compactifications




# Axion-like particles

- Light axion-like particles (ALPs) are one of the most motivated ways to extend the Standard Model
- They arise generically in string theory – e.g. a light ALP is always present in the Large Volume Scenario.
- Phenomenologically, they are parametrised by the coupling

$$a g_{a\gamma\gamma} \mathbf{E} \cdot \mathbf{B} \equiv \frac{a}{M} \mathbf{E} \cdot \mathbf{B}$$

- In the presence of a background  $\mathbf{B}$  field, the ALP  $a$  and photon  $\gamma$  eigenstates mix, leading to photon-ALP oscillations (cf neutrino oscillations)

# Dark Matter



Dark, unknown part of the energy budget of the universe

Constitutes approx 25% of total energy in the universe

# II. The Methods



# Two probes of fundamental physics

- I focus in this talk on two uses of the spectra of Active Galactic Nuclei to search for new physics residing in dark, weakly coupled hidden sectors
  1. Searches for ALPs using spectral irregularities induced by ALP-photon mixing (original motivation for this work)
  2. Search for dark matter absorption lines (secondary post-data motivation)

# How to search for ALPs?

- The basic physics used here to look for ALPs is very simple.
  1. Send photons from A to B
  2. Have a magnetic field inbetween A and B
  3. Photon-ALP interconversion causes some of these photons to oscillate into ALPs
  4. The photon spectrum on arrival at B will show modulations compared to the source photon spectrum at A.
- In our case, the source A will be the central AGN (Active Galactic Nucleus) of the Perseus galaxy cluster and B is the *Chandra* X-ray telescope



AGN



NGC1275



Milli- parsec

Hundred kilo-parsecs

Megaparsecs

Perseus cluster



68 Mpc

Chandra



# Active Galactic Nuclei are point sources

- X-ray emission from AGNs comes from extremely small physical region
- This follows from the time variability of AGN spectra: intensities fluctuate on hour to day timescales, implying emission originates from within a light-day
- Basic components to X-ray spectrum are
  1. Power-law
  2. Reflection spectrum (incident photons illuminate accretion disc, resulting in fluorescent emission) – in practice manifest as neutral Fe  $K\alpha$  line at 6.4 keV.
  3. Thermal soft excess (origin not entirely known)

# AGNs: the standard Unified Model



Credit ESA/NASA, AVO project, Paolo Padavani



# The Perseus Cluster

- The Perseus galaxy cluster is the brightest X-ray galaxy cluster in the sky, and is located at a redshift of 0.0176
- It is a cool-core cluster centred around the Seyfert galaxy NGC1275 and its Active Galactic Nucleus.
- The Milky Way column density along the line of sight to Perseus is high, at  $n_H = 1.5 \times 10^{21} \text{ cm}^{-2}$  (implies significant absorption of soft X-rays).
- The Perseus cluster is the subject of enormous observation time with the *Chandra* X-ray telescope, totalling 1.5 Ms – gives over 500,000 photon counts from the central AGN



Optical image of Perseus, credit *R. Jay GaBany*, [Cosmotography.com](https://www.cosmotography.com)





X-ray image of the  
Perseus cluster:  
NGC1275 AGN is the  
central white dot

The AGN jets blow  
bubbles into the  
surrounding intra-cluster  
medium

Perseus in X-rays (NASA, Chandra)

# Photon-ALP Conversion

- Source is NGC1275, destination is earth: intervening magnetic field is **magnetic field of the Perseus cluster**.
- Galaxy clusters are particularly good locations for photon-ALP interconversion
- Magnetic fields extend over approx. 1 Mpc regions, with coherence lengths in 1- 10kpc region.
- Magnetic field strengths are 1 – 10 microGauss.
- Photon-ALP couplings  $g_{a\gamma\gamma}$  of  $10^{-12}$  to  $10^{-11}$  GeV<sup>-1</sup> generate conversion probabilities of order 10 – 50%.
- No **exact knowledge** of Perseus magnetic field; central value should be in range 10 – 25 microGauss.


# Photon-ALP Conversion – why X-rays?

- Axion-photon interconversion (for  $m_a < 10^{-12} \text{eV}$ , effectively massless) in galaxy clusters:

$$P_{\gamma \rightarrow a} = \frac{1}{2} \frac{\Theta^2}{1 + \Theta^2} \sin^2 \left( \Delta \sqrt{1 + \Theta^2} \right)$$

$$\Theta = 0.28 \left( \frac{B_{\perp}}{1 \mu\text{G}} \right) \left( \frac{\omega}{1 \text{keV}} \right) \left( \frac{10^{-3} \text{cm}^{-3}}{n_e} \right) \left( \frac{10^{11} \text{GeV}}{M} \right) \quad \Delta = 0.54 \left( \frac{n_e}{10^{-3} \text{cm}^{-3}} \right) \left( \frac{L}{10 \text{kpc}} \right) \left( \frac{1 \text{keV}}{\omega} \right)$$


- Sweet spot at X-ray energies:






# ALPS


AGNs are bright point sources of photons



Photons pass through galaxy cluster magnetic field




ALP-Photon conversion induces irregularities in observed X-ray spectrum



Precise form of modulations depends on cluster magnetic field

# Simulated photon survival probability



This would modulate the true spectrum

# III. The Results

## A. ALP Searches





*Chandra* X-ray telescope

~1.5 billion USD


15 years operation

Mature, well understood instrument


Large public observational data archive

# The Observations

- NGC1275 observed by *Chandra* in 2002 and 2004 for 1Ms with ACIS-S and 0.5 Ms in 2009 with ACIS-I.
- In ACIS-S observations, NGC1275 is on-axis, in 2009 observations 300ks with NGC1275 around 4 arcmin off-axis and 200ks with NGC1275 around 8 arcmin off-axis.
- Treat these three sets separately, focus on last case.
- *Chandra* on-axis point spread function is around 0.5 arcsec diameter on-axis, broadening to around 10 arcsec diameter when source is around 8 arcmin off-axis.



# Complete extraction for ACIS-I edge



Fit to absorbed power law gives two main features – excess at 2 – 2.2 keV, deficit at 3.4 – 3.5 keV

# Features in ACIS-I Edge Data

Two main features:


1. Excess at 2 – 2.2 keV

Subtle because of effective area dip at these energies

Possible to generate fake excesses via energy mismeasurement

2. Deficit at 3.4 – 3.5 keV


No obvious systematic effects - will discuss in detail below!



# ALP Constraints

Unambiguous statement – there are no spectral irregularities greater than 10%

ALP couplings leading to 20-30% irregularities are excluded



# ALP Constraints

1. Reasonable ( $B_{\text{central}} = 25 \mu\text{G}$ , 100 domains between 3.5 and 10kpc)

$$g_{a\gamma\gamma} < 1.5 \times 10^{-12} \text{ GeV}^{-1}$$


2. Conservative: ( $B_{\text{central}} = 15 \mu\text{G}$ , 100 domains between 0.7 and 10kpc)

$$g_{a\gamma\gamma} < 3.8 \times 10^{-12} \text{ GeV}^{-1}$$

3. Ultra-conservative: ( $B_{\text{central}} = 10 \mu\text{G}$ , 100 domains between 0.7 and 10kpc)

$$g_{a\gamma\gamma} < 5.6 \times 10^{-12} \text{ GeV}^{-1}$$

Absence of any spectral modulations at 20-30% level gives leading bounds on ALP-photon coupling at small mass



Bounds would be stronger still if we knew that either

2 - 2.2 keV feature

or

3.4 - 3.5 feature


were definitely not real.....

# III. The Results

## B. Dark Matter and the 3.5 keV Line



# Complete extraction for ACIS-I edge



Two main features – excess at 2 – 2.2 keV, deficit at 3.4 – 3.5 keV

# Look at 3.4 – 3.5 keV feature more closely...


We fit from 0.8 to 5 keV and cut out 1.8 – 2.3 keV region to avoid biasing the fit.

Fit with `xswabs * (xspowerlw + xsbapec)`


i.e. `Absorption * (power law + thermal cluster emission)`

Use thermal emission cluster parameters determined by Hitomi

For convenience, only show fit from 2.5 – 4.5 keV



Good fit – chi squared of 273/250 dof – with dip clearly visible at 3.5 keV



Now include a negative Gaussian..... $\Delta\chi^2 = 20.0$  for 2 dof

Over 4 sigma preference for dip/absorption at  $(3.54 \pm 0.02)$  keV! (cluster frame)

# The 3.5 KeV Line....

**Exactly** the same energy as the 3.5 keV line excess....

## DETECTION OF AN UNIDENTIFIED EMISSION LINE IN THE STACKED X-RAY SPECTRUM OF GALAXY CLUSTERS

ESRA BULBUL<sup>1,2</sup>, MAXIM MARKEVITCH<sup>3</sup>, ADAM FOSTER<sup>1</sup>, RANDALL K. SMITH<sup>1</sup> MICHAEL LOEWENSTEIN<sup>2,4</sup>, AND SCOTT W. RANDALL<sup>1</sup>

<sup>1</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA, USA

<sup>2</sup> CRESST and X-ray Astrophysics Laboratory, NASA Goddard Space Flight Center, Greenbelt, MD, USA

<sup>3</sup> NASA Goddard Space Flight Center, Greenbelt, MD, USA

<sup>4</sup> Department of Astronomy, University of Maryland, College Park, MD, USA

*Submitted to ApJ, 2014 February 10, Accepted 2014 April 28*

### ABSTRACT

We detect a weak unidentified emission line at  $E = (3.55 - 3.57) \pm 0.03$  keV in a stacked *XMM-Newton* spectrum of 73 galaxy clusters spanning a redshift range 0.01 – 0.35. MOS and PN observations independently show the presence of the line at consistent energies. When the full sample is divided

## **An unidentified line in X-ray spectra of the Andromeda galaxy and Perseus galaxy cluster**

A. Boyarsky<sup>1</sup>, O. Ruchayskiy<sup>2</sup>, D. Iakubovskiy<sup>3,4</sup> and J. Franse<sup>1,5</sup>

<sup>1</sup>Instituut-Lorentz for Theoretical Physics, Universiteit Leiden, Niels Bohrweg 2, Leiden, The Netherlands


<sup>2</sup>Ecole Polytechnique Fédérale de Lausanne, FSB/ITP/LPPC, BSP, CH-1015, Lausanne, Switzerland

<sup>3</sup>Bogolyubov Institute of Theoretical Physics, Metrologichna Str. 14-b, 03680, Kyiv, Ukraine

<sup>4</sup>National University “Kyiv-Mohyla Academy”, Skovorody Str. 2, 04070, Kyiv, Ukraine

<sup>5</sup>Leiden Observatory, Leiden University, Niels Bohrweg 2, Leiden, The Netherlands

We report a weak line at  $3.52 \pm 0.02$  keV in X-ray spectra of M31 galaxy and the Perseus galaxy cluster observed by MOS and PN cameras of *XMM-Newton* telescope. This line is not known as an atomic line in the



3.5keV line was expected to be resolved by Hitomi.....



Launch of Hitomi from Tanegashima Space Centre

17th February 2016




## The quiescent intracluster medium in the core of the Perseus cluster

The Hitomi collaboration\*

Clusters of galaxies are the most massive gravitational objects in the Universe and are still forming. They are

Hitomi returned a ground-breaking spectrum of Perseus before its tragic loss in March 2016



Energy resolution around 5eV, 20x better than Chandra or XMM!




# Images of the centre of Perseus

CHANDRA




Best angular resolution

HITOMI




Best energy resolution

# Hitomi Spectrum of Perseus Cluster



But look  
closely near  
3.5 keV.....

# Hitomi best-fit line properties




**No Excess!**

Overall best-fit line is negative  
at 3.54 keV with normalisation  
of

$-8 \times 10^{-6}$  photons  $\text{cm}^{-2}$   $\text{s}^{-1}$

2.5 sigma significance

# Hitomi view of Perseus



Hitomi cannot separately resolve AGN and thermal cluster emission

Its best-fit value

$(-8 \times 10^{-6} \text{ photons cm}^{-2} \text{ s}^{-1})$


is sensitive only to the SUM of

(3.54 keV features in cluster emission)

PLUS

(3.54 keV features in AGN spectrum)

# Hitomi view of Perseus



Hitomi best-fit value at 3.54 keV:

$$(-8 \times 10^{-6} \text{ photons cm}^{-2} \text{ s}^{-1})$$

XMM Excess (excluding AGN) in Hitomi Field of View

$$(9.0 \pm 2.9) \times 10^{-6} \text{ photons cm}^{-2} \text{ s}^{-1}$$

Deficit in AGN from Chandra (rescaled from 2009 to 2016 AGN luminosity)

$$(-16.7 \pm 3.6) \times 10^{-6} \text{ photons cm}^{-2} \text{ s}^{-1}$$

All consistent!

## 3.5 keV line in Perseus

**Deficit/absorption** in the spectrum of very bright point source


**Excess/emission** in the diffuse spectrum throughout the cluster

**At the same energy** – how to explain this in one model?



# Fluorescent Dark Matter

Dark matter absorbs and re-emits 3.5 keV photons; generates both AGN deficit and diffuse excess



# Fluorescent Dark Matter

$$\mathcal{L} \supset \frac{1}{M} \bar{\chi}_2 \sigma_{\mu\nu} \chi_1 F^{\mu\nu},$$

Simplest model involves two states ( $\chi_1$  and  $\chi_2$ )

Dark matter is in ground state  $\chi_1$

Absorption of real 3.5 keV photons takes it to excited state  $\chi_2$

Instant decay  $\chi_2 \rightarrow \chi_1 \gamma$  leads to diffuse 3.5 keV excess

(Lots of work on excitation via dark matter collision, but this scenario is surprisingly little studied)

# Conclusions

1. X-ray astronomy is a powerful probe of fundamental physics
2. Existing, archival *Chandra* observations of Perseus constraint offer leading constraints on  $g_{a\gamma\gamma}$  for light ALPs with  $m < 10^{-12}$  eV
3. Data contains a striking dip in the AGN spectrum at  $(3.54 \pm 0.02)$  keV – dark matter absorption?
4. 3.5 keV line is **compelling evidence** for new physics

THANK YOU!