

# News on the Measurement of the W Boson Mass






# Electroweak Precision Physics

# Summary of the Electroweak Sector

- The electroweak sector of the Standard Model has five parameters
  - $a_{em}$ ,  $G_F$ ,  $m_W$ ,  $m_Z$ ,  $\sin^2\theta_W$
  - (+  $m_H$  for the scalar sector)
- However, they are not independent, but related by theory


$$\sin^2\theta_W = 1 - \frac{m_W^2}{m_Z^2}$$

$$m_W^2 \sin^2\theta_W = \frac{\pi\alpha}{\sqrt{2}G_F}$$



# Radiative Corrections

- Tree-level not sufficient
  - The impact of corrections stored in **EW form factors**
- The relation between SM parameters appear with **quadratic dependence on  $m_{\text{top}}$ , logarithmic dependence on  $M_H$**
- Idea of electroweak fits
  - Measure many different observables
  - Calculate the relations between all observables
  - **Probe the consistency of the SM / Predict observables**



$$\sin^2 \theta_{\text{eff}}^f = \kappa_Z^f \sin^2 \theta_W$$

$$g_{V,f} = \sqrt{\rho_Z^f} (I_3^f - 2Q^f \sin^2 \theta_{\text{eff}}^f)$$

$$g_{A,f} = \sqrt{\rho_Z^f} I_3^f$$




$$M_W^2 = \frac{M_Z^2}{2} \left( 1 + \sqrt{1 - \frac{\sqrt{8}\pi\alpha(1 + \Delta r)}{G_F M_Z^2}} \right)$$

$$M_W \left( \ln(M_H), m_t^2, M_Z, \Delta\alpha_{\text{had}}^{(5)}(M_Z^2), \alpha_s(M_Z^2) \right)$$

$$\sin^2 \theta_{\text{eff}}^f \left( \ln(M_H), M_H, m_t^2, M_Z, \Delta\alpha_{\text{had}}^{(5)}(M_Z^2), \alpha_s(M_Z^2) \right)$$


# Input and Outcome to the Electroweak Fit

- Input for the global electroweak fit mostly from
  - LEP: Z boson observables
  - Tevatron: W boson, top quark mass
  - LHC: **Higgs Boson, top quark mass**
- Result: **Amazing predictions!**
  - Top-Quark mass before its discovery
  - Higgs-Boson mass before its discovery and the funding argument for the LHC





# Z Boson Measurements at LEP

- Z bosons can be produced directly in  $e^+e^-$  collisions, i.e. we can scan the Breit-Wigner lineshape
- We do **not need to reconstruct energy** of the decay particles
  - Mass measurement is reduced to a **pure counting experiment**
- Reach enormous precision
  - **Mass known to 2 MeV!**
- Not only crucial input for the electroweak fit, but also for the LHC detector calibration!




# Where do we stand and what do we need?

- Z boson observables: **LEP was great**; we can't do better (now)
- Top-quark mass: huge progress; **new approaches** to overcome theoretical unc. on  $m_{\text{top}}$ 
  - (SM prediction  $\pm 2.5$  GeV, measurement unc.  $< 0.7$  GeV)
- Higgs mass: Only **logarithmic dependence**, hence not important for the fit
- **W mass:** SM prediction at 5 MeV precision, known only to 19 MeV!
  - **Here we can gain a lot!**




# Why do we need that?

- So far a “simple” thing: test consistency of the SM
  - Current **p-value = 0.22**
- But electroweak precision measurements are **sensitive to several new physics scenarios**, e.g. SUSY
  - Radiative correction depends on mass splitting ( $\Delta m^2$ ) between squarks in SU(2) doublet
  - Precision on  $m_W$  could significantly limit the allowed MSSM space



Inspired by [S. Heinemeyer et. al. arXiv:1311.1663]





# Measurement Strategy


# Mass Sensitive Variables

- **Main signature:** final state lepton (electron or muon):  $p_T(\text{lepton})$
- **Recoil:** sum of “everything else” reconstructed in the calorimeters
  - a measure of  $p_T(W, Z)$
  - gives us also missing transverse energy!
- **Derived quantities**

$$\vec{u}_T = \sum_i \vec{E}_{T,i}$$



$$\vec{p}_T^{\text{miss}} = -\left( \vec{p}_T^l + \vec{u}_T \right)$$

$$m_T = \sqrt{2 p_T^l p_T^{\text{miss}} (1 - \cos \Delta\phi)}$$




# Mass Sensitive Variables

- Sensitive final state distributions:
  - Lepton transverse momentum  $\mathbf{p}_T(l)$
  - Transverse mass:  $\mathbf{m}_T$
  - Missing transverse energy (“neutrino  $\mathbf{p}_T$ ”):  $\mathbf{p}_T^{\text{miss}}$
- Template-Fit approach
  - Assume various W boson mass values in MC event generator and predict the  $\mathbf{p}_T(l)$  ,  $\mathbf{m}_T$  ,  $\mathbf{p}_T^{\text{miss}}$  distributions
  - Compare to data
  - Mass determination by  $\chi^2$  minimization



# Why is this measurement complicated?





JOHANNES GUTENBERG  
UNIVERSITÄT MAINZ



Simple Life in the  
USA

# Measurement Strategy at LHC and Tevatron





- **Parameterized** (not full) **simulation** that includes all corrections
- Low pile-up
- CDF-Experiment: e/mu channel
  - Only 20% of data-set used
  - Calibration: J/Psi, Upsilon, Z
- D0-Experiment: e-channel
  - Acceptance up to  $\eta < 1.0$
  - Calibration: Z boson only

- Analysis based on **full simulation**
  - PowhegPythia (NLO QCD+PS);
  - QED FSR using PHOTOS
  - **Reweighting to correct for physics and detector modelling**
- Data-Set: Run 1 (2011)
  - 7 TeV,  $4.6 \text{ fb}^{-1}$  (e),  $4.1 \text{ fb}^{-1}$  ( $\mu$ )
  - Mature, well understood data; moderate (but still significant) pile-up

# Production of W Bosons at the LHC and Tevatron

- **Tevatron:** Proton / anti-protons
  - Main production involves up and down quarks
  - small impact of heavy quarks
  - No differences between  $W^+/W^-$
  - Similar production of Z bosons
  
- **LHC:** Proton-proton collisions
  - Heavy quarks become important in the production
  - Different production modes of  $W^+$  and  $W^-$
  - Z Boson production still dominated by light quarks



# Measurements at the Tevatron

- Do not worry about tables
  - These tables have been my life for 5 years!
- Experimental uncertainties
  - CDF: 10 MeV (Lepton Scale, Hadronic Recoil)
  - D0: 18 MeV (Lepton Scale)
- Physics modelling dominated by PDFs: **10-11 MeV for both**
- Precision after >10 years
  - **CDF: 19 MeV**
  - **D0: 26 MeV**


[*Phys Rev D88 (2013)*]

Source	Uncertainty (MeV)
Lepton energy scale and resolution	7
Recoil energy scale and resolution	6
Lepton removal from recoil	2
Backgrounds	3
Experimental subtotal	10
Parton distribution functions	10
QED radiation	4
$p_T(W)$ model	5
Production subtotal	12
Total systematic uncertainty	15
$W$ -boson event yield	12
Total uncertainty	19

Source	Uncertainty (MeV)
Electron energy calibration	16
Electron resolution model	2
Electron shower modeling	4
Electron energy loss model	4
Recoil energy scale and resolution	5
Electron efficiencies	2
Backgrounds	2
Experimental subtotal	18
Parton distribution functions	11
QED radiation	7
$p_T(W)$ model	2
Production subtotal	13
Total systematic uncertainty	22
$W$ -boson event yield	13
Total uncertainty	26

# Signal Selection and Measurement Regions

- Lepton selections
  - Muons :  $|\eta| < 2.4$ ; isolated
  - Electrons :  $0 < |\eta| < 1.2$  or  $1.8 < |\eta| < 2.4$ ; isolated
- Kinematic requirements
  - $p_T > 30\text{GeV}$   $p_T^{\text{miss}} > 30\text{GeV}$
  - $m_T > 60\text{GeV}$   $\ell_T < 30\text{GeV}$
- **Measurement categories**
  - Electron/muon channel,  $p_T$ ,  $m_T$ -Fits, 3/4 rapidity regions, W boson charge
  - Muon Channel: 7.8 M events
  - Electron Channel: 5.9 M events






Physics Modelling

# Physics Modelling

- **No available generator can describe all observed features:**  
 $p_T(Z)/p_T(W)$ ,  $A_i$ , ...
  - Variation of  $d\sigma/dm$  modeled with a Breit-Wigner + EW cor.
  - $d\sigma/dp_T$  is modeled with PS MC
  - $d\sigma/dy$  modeled at NNLO
  - $A_i(y, pt)$  modeled at NNLO
- **QCD aspects**
  - Rapidity,  $p_T$  distributions; angular distributions
- **EW aspects**
  - ISR and FSR QED corrections
  - Missing higher-order effects





$$\frac{d\sigma}{dp_1 dp_2} = \left( \frac{d\sigma}{dm} \right) \left( \frac{d\sigma}{dy} \right) \left( \frac{d\sigma(p_t, y)}{dpt} \frac{1}{\sigma(y)} \right).$$

$$+ \left( \sum_i A_i(y, p_t) P_i(\cos\theta, \phi) \right)$$



# Transverse Momentum (A several years effort)

- **Traditional approach:** fit predictions to Z data, apply to W
  - primordial  $k_T$ ;  $\alpha_S^{\text{ISR}}$ ; ISR cut-off
  - Tested with Powheg+Pythia8, and **Pythia8 standalone**
- **Associated Uncertainties:** Z Boson Data, Parton Show Variations and
  - $Z \rightarrow W$  extrapolation : factorization scale variations (separately for light- and heavy-quark induced production), heavy quark masses



# Transverse Momentum (A several years effort)

- Theoretically more advanced calculations were also attempted
  - DYRES (and other resummation codes : ResBos, CuTe)
  - Powheg MiNLO + Pythia8
- All predict a **harder  $p_T(W)$  spectrum for given  $p_T(Z)$**  distribution
  - Behaviour is **disfavoured by data** (see later)



# Overview of QCD Uncertainties

- CT10nnlo uncertainties (synchronized in DYNNLO and Pythia) + envelope comparing CT10 to CT14 and MMHT.
  - Strong **anti-correlation of uncertainties for  $W_+$  and  $W^-$ !**
- AZ tune uncertainty; parton shower PDF and factorization scale; heavy-quark mass effects
- **$A_i$  uncertainties from Z data**; envelope for  $A_2$  discrepancy

W-boson charge Kinematic distribution	$W^+$		$W^-$		Combined	
	$p_T^\ell$	$m_T$	$p_T^\ell$	$m_T$	$p_T^\ell$	$m_T$
$\delta m_W$ [MeV]						
Fixed-order PDF uncertainty	13.1	14.9	12.0	14.2	8.0	8.7
AZ tune	3.0	3.4	3.0	3.4	3.0	3.4
Charm-quark mass	1.2	1.5	1.2	1.5	1.2	1.5
Parton shower $\mu_F$ with heavy-flavour decorrelation	5.0	6.9	5.0	6.9	5.0	6.9
Parton shower PDF uncertainty	3.6	4.0	2.6	2.4	1.0	1.6
Angular coefficients	5.8	5.3	5.8	5.3	5.8	5.3
Total	15.9	18.1	14.8	17.2	11.6	12.9



# Detector Calibration


# Mass Sensitive Variables

## ■ Lepton calibration

- momentum calibration using the Z peak
- efficiency corrections (reconstruction, identification, trigger) rederived via tag- and probe-method in 3 dimensions

## ■ Recoil calibration

- Event activity corrections
- Recoil response calibration using expected  $p_T$  balance between lepton pairs and  $u_T$  in Z events





# Muon Momentum and Efficiency Calibration

- Parameterization of momentum corrections:

$$p_T^{MC,corr} = p_T^{MC} \times [1 + \alpha(\eta, \phi)] \times [1 + \beta_{curr}(\eta) \cdot G(0,1) \cdot p_T^{MC}]$$



- Momentum scale and resolution corrections probed using  $Z \rightarrow \mu\mu$  events
- Also to correct: **Track sagitta** bias from rotational detector deformations
  - Using  $W \rightarrow \mu\nu$  events for overall effect and  $Z \rightarrow \mu\mu$  for eta-calibration



# Hadronic Recoil Calibration


## ■ Recoil correction steps

- Step 1: Equalize pile-up multiplicity distribution in data and MC
  - Step 2: Correct for residual differences in the  $\Sigma E_T$  distribution
    - removes  $u_T$  resolution discrepancy due to imperfect event activity mis-modeling
  - Can be done directly in W events – no extrapolation systematics
- ## ■ MC based closure test!



# A distribution which took us months

- Typically one expects a  $\Phi$  symmetry of the detector response (and the physics)
- We observed significant differences to MC
  - **offset of the interaction point** with respect to the detector center in the transverse plane
  - **Non-zero crossing angle** between the proton beams
  - **$\phi$ -dependent response of the calorimeters**



# 3 tables after 3 years of work

- Experimental uncertainty due to muon detector calibration on the 10 MeV level
  - In terms of average accuracy on the position resolution, this means  $\mu\text{m}$ -precision!
- Not even discussed here: **How to estimate backgrounds**
  - We control the background contributions on **a rel. 5% level!**
  - Final background related uncertainties
    - $p_T$ -fit: 3-5 MeV
    - $m_T$ -fit: 8-9 MeV (elec.)
    - $m_T$ -fit: 3-5 MeV (muon)

$ \eta_\ell $ range	[0.0, 0.8]		[0.8, 1.4]		[1.4, 2.0]		[2.0, 2.4]		Combined	
Kinematic distribution	$p_T^\ell$	$m_T$								
$\delta m_W$ [MeV]										
Momentum scale	8.9	9.3	14.2	15.6	27.4	29.2	111.0	115.4	8.4	8.8
Momentum resolution	1.8	2.0	1.9	1.7	1.5	2.2	3.4	3.8	1.0	1.2
Sagitta bias	0.7	0.8	1.7	1.7	3.1	3.1	4.5	4.3	0.6	0.6
Reconstruction and isolation efficiencies	4.0	3.6	5.1	3.7	4.7	3.5	6.4	5.5	2.7	2.2
Trigger efficiency	5.6	5.0	7.1	5.0	11.8	9.1	12.1	9.9	4.1	3.2
Total	11.4	11.4	16.9	17.0	30.4	31.0	112.0	116.1	9.8	9.7

$ \eta_\ell $ range	[0.0, 0.6]		[0.6, 1.2]		[1.82, 2.4]		Combined	
Kinematic distribution	$p_T^\ell$	$m_T$	$p_T^\ell$	$m_T$	$p_T^\ell$	$m_T$	$p_T^\ell$	$m_T$
$\delta m_W$ [MeV]								
Energy scale	10.4	10.3	10.8	10.1	16.1	17.1	8.1	8.0
Energy resolution	5.0	6.0	7.3	6.7	10.4	15.5	3.5	5.5
Energy linearity	2.2	4.2	5.8	8.9	8.6	10.6	3.4	5.5
Energy tails	2.3	3.3	2.3	3.3	2.3	3.3	2.3	3.3
Reconstruction efficiency	10.5	8.8	9.9	7.8	14.5	11.0	7.2	6.0
Identification efficiency	10.4	7.7	11.7	8.8	16.7	12.1	7.3	5.6
Trigger and isolation efficiencies	0.2	0.5	0.3	0.5	2.0	2.2	0.8	0.9
Charge mis-measurement	0.2	0.2	0.2	0.2	1.5	1.5	0.1	0.1
Total	19.0	17.5	21.1	19.4	30.7	30.5	14.2	14.3


$W$ -boson charge	$W^+$		$W^-$		Combined	
Kinematic distribution	$p_T^\ell$	$m_T$	$p_T^\ell$	$m_T$	$p_T^\ell$	$m_T$
$\delta m_W$ [MeV]						
$\langle \mu \rangle$ scale factor			0.2	1.0	0.2	1.0
$\Sigma \bar{E}_T$ correction			0.9	12.2	1.1	10.2
Residual corrections (statistics)			2.0	2.7	2.0	2.7
Residual corrections (interpolation)			1.4	3.1	1.4	3.1
Residual corrections ( $Z \rightarrow W$ extrapolation)			0.2	5.8	0.2	4.3
Total			2.6	14.2	2.7	11.8
					2.6	13.0



# Z Boson Cross- Checks


# Can we learn even more from the Z Boson?

- Measurement of the Z boson mass via the invariant mass distribution closes by construction
- Treat one decay lepton as neutrino and mimic W boson decay
  - $p_T$  and  $m_T$  fits
- Allows to test the full correction procedure, however without the transfer uncertainties to W bosons




# Results of the Z Boson Mass Fits

- Results consistent with  $m_Z$ , within uncertainties
- Fitted values a bit low on average, but these are all from the same events
  - if the decay angle is a bit more forward in data, all fit results decrease simultaneously.




Lepton charge Distribution	$\ell^+$		$\ell^-$		Combined	
	$p_T^\ell$	$m_T$	$p_T^\ell$	$m_T$	$p_T^\ell$	$m_T$
$\Delta m_Z$ [MeV]						
$Z \rightarrow ee$	$13 \pm 31 \pm 10$	$-93 \pm 38 \pm 15$	$-20 \pm 31 \pm 10$	$4 \pm 38 \pm 15$	$-3 \pm 21 \pm 10$	$-45 \pm 27 \pm 15$
$Z \rightarrow \mu\mu$	$1 \pm 22 \pm 8$	$-35 \pm 28 \pm 13$	$-36 \pm 22 \pm 8$	$-1 \pm 27 \pm 13$	$-17 \pm 14 \pm 8$	$-18 \pm 19 \pm 13$
Combined	$5 \pm 18 \pm 6$	$-58 \pm 23 \pm 12$	$-31 \pm 18 \pm 6$	$1 \pm 22 \pm 12$	$-12 \pm 12 \pm 6$	$-29 \pm 16 \pm 12$





W Boson Analysis

# Control Distributions (non $m_W$ sensitive)







# Crucial Test of $p_T(W)$ modelling

- Remember the problem with the  $p_T(W)$  description?
  - How do we know, which MC generator to trust?
  - How do we know, that our assigned uncertainty makes sense?
- The  $u_{||}(l)$  distribution is very sensitive to the underlying  $p_T(W)$  distribution
  - Can exploit this feature to verify the accuracy of our baseline model, and compare to alternative calculations





# W-Mass Distributions: Electrons

- Predictions set to final combined  $m_W$  value
- Dip at 40 GeV was studied thoroughly
  - No striking effects: stays at  $2\sigma$
  - Only mild impact on final  $m_W$





# W-Mass Distributions: Muons

- Very good agreement for muons
- Overall:  $\chi^2/n_{\text{dof}}$  probability distribution from 84 data/prediction comparison
  - $\langle P \rangle = 0.54$



# A Little Bit of History



- Over many years we investigated differences in blinded  $m_W$  mass-fits in different channels, templates, categories
  - Only after all corrections applied (and all bugs where found), we achieved consistent results







Final Measurement

# $m_W$ Fit Results in Various Categories




- Illustration of fit-results in all measurement categories based on  $p_T$  and  $m_T$  templates for  $W^+$  and  $W^-$  in the electron and muon channel
- Compatibility tests **performed before unblinding:  $\chi^2/n_{\text{dof}} = 29 / 27$**

# Testing stability for different Fit-Ranges



- The fitting ranges have been optimized on MC predictions for reducing overall systematic uncertainty
- Stability of fit-results was studied**, taking into account (de-)correlations

# $m_W$ Fit Results in Various Combinations



*Nobody cares about  
 your method. People  
 remember only your  
 last number!*

Final measured mass of the W boson




$$= 80.370 \pm 0.007(\text{stat.}) \pm 0.011(\text{exp.}) \pm 0.014 \text{ (mod) GeV}$$

$$= \mathbf{80.370 \pm 0.019 \text{ GeV}}$$




# Comparison to previous measurements and the Global Electroweak Fit

- **Good news:** New measurement reaches precision of CDF and is now the world leading measurement
- Bad news: **We are even more Standard Model ...**



# Outlook

- LHC
  - CMS will hopefully also publish soon a  $m_W$  measurement
  - ATLAS still has data-sets of 2012/2015/2016
  - **Room for improvements** on the theory and calibration
    - **IMHO: <10 MeV is feasible**
  
- Tevatron
  - x2-5 more statistics available (+ **forward detectors**)
  - **Use improved PDFs based on LHC measurements**



# Summary

- Five years effort from a large team and the full collaboration comes to an end
  - Many important discussions with colleagues from Theory and CMS
- First W mass measurement at the LHC unfortunately shows no signs of BSM
- Need some “break-through” in the MC event generator developments
  - We have quite some ideas to improve the achieved precision significantly!