



The Galactic Center Gamma-Ray Excess and its Interpretations


Dan Hooper – Fermilab and the University of Chicago
TeV Particle Astrophysics Workshop (TeVPA) 2018
August 30, 2018

The Galactic Center GeV Excess

- A bright and highly statistically significant excess of gamma-rays has been observed from the region surrounding the Galactic Center
- Although a consensus has formed regarding the basic features of this signal, its origin is still a topic of considerable debate



Among other references, see:

DH, Goodenough (2009, 2010)
DH, Linden (2011)
Abazajian, Kaplinghat (2012)
Gordon, Macias (2013)
Daylan, et al. (2014)
Calore, Cholis, Weniger (2014)
Murgia, et al. (2015)
Ackermann et al. (2017)



What Produces the Excess?


- Annihilating dark matter?
- A large population of centrally located millisecond pulsars?
- A recent outburst of cosmic rays?



Dark Matter and the GeV Excess

The observed characteristics of the excess are in good agreement with the expectations of annihilating dark matter

- **Spectrum:** Well fit by a $\sim 40\text{-}70$ GeV particle annihilating to quarks, and is uniform across the Inner Galaxy
- **Morphology:** Approximately spherically symmetric, with a flux that falls as $\sim r^{-2.4}$ out to at least $\sim 10^\circ$, consistent with a DM halo only slightly steeper than NFW
- **Intensity:** Requires an annihilation cross section of $\sigma v \sim 10^{-26}$ cm³/s, near the value of a thermal relic




Daylan, Finkbeiner, Hooper, Linden, Rodd, Slatyer (2014)
 Calore, Cholis, Weniger; Calore, Cholis, McCabe, Weinger (2014);

Millisecond Pulsars and The Galactic Center Gamma-Ray Excess

The Two Main Arguments in Favor of Pulsars:

- The gamma-ray spectrum of observed pulsars
- Small-scale power in the gamma-ray emission from the Inner Galaxy



Small Scale Power Among Inner Galaxy γ -Rays

- In 2015, two groups found that the \sim GeV photons from the direction of the Inner Galaxy are more clustered than predicted from smooth backgrounds, suggesting that the GeV excess might be generated by a population of unresolved point sources
- Lee et al. used a non-Poissonian template technique to show that the photon distribution within $\sim 10^\circ$ of the Galactic Center (masking within 2° of the Galactic Plane) is *clumpy*, potentially indicative of an unresolved point source population
- Bartels et al. reach a similar conclusion employing a wavelet technique

Lee, Lisanti, Safdi, Slatyer, Xue, arXiv:1506.05124


Bartels, Krishnamurthy, Weniger, arXiv:1506.05104

Small Scale Power Among Inner Galaxy γ -Rays


- A typical Fermi Inner Galaxy analysis might include the following spatial templates:
 - 1) Galactic diffuse emission
 - 2) Fermi Bubbles
 - 3) Isotropic background
 - 4) Known point sources
 - 5) Dark matter annihilation products (generalized NFW²)

- Lee et al. then add a number of non-Poissonian templates to model the distribution of unresolved point sources:
 - 5) Isotropically distributed point sources
 - 6) Disk-correlated point sources
 - 7) NFW² correlated point sources

Small Scale Power Among Inner Galaxy γ -Rays



Small Scale Power Among Inner Galaxy γ -Rays




Bottom Line: A population of $\sim 10^3$ point sources with luminosities near Fermi's detection threshold could potentially account for the GeV Excess

Evidence For Unresolved Point Sources?

- It is difficult to tell whether these clustered gamma-rays result from unresolved sources, or from backgrounds that are less smooth than are being modeled



Evidence For Unresolved Point Sources?

- It is difficult to tell whether these clustered gamma-rays result from unresolved sources, or from backgrounds that are less smooth than are being modeled



Evidence For Unresolved Point Sources?

- It is difficult to tell whether these clustered gamma-rays result from unresolved sources, or from backgrounds that are less smooth than are being modeled



Evidence For Unresolved Point Sources?

- It is difficult to tell whether these clustered gamma-rays result from unresolved sources, or from backgrounds that are less smooth than are being modeled
- These clusters consist of only a few photons each, on top of large and imperfectly known backgrounds

Lee, Lisanti, Safdi, Slatyer, Xue, arXiv:1506.05124
(see also Bartels, Krishnamurthy, Weniger, arXiv:1506.05104)

Evidence For Unresolved Point Sources?

- It is difficult to tell whether these clustered gamma-rays result from unresolved sources, or from backgrounds that are less smooth than are being modeled
- These clusters consist of only a few photons each, on top of large and imperfectly known backgrounds
- Gamma-ray point source identification is difficult in the Galactic Center region – even for bright sources – and the contents of source catalogs depend strongly on how one treats diffuse backgrounds
(try comparing the membership of Fermi's 3FGL, 1FIG, and 2FIG catalogs)

Lee, Lisanti, Safdi, Slatyer, Xue, arXiv:1506.05124
(see also Bartels, Krishnamurthy, Weniger, arXiv:1506.05104)


Millisecond Pulsars and The Galactic Center Gamma-Ray Excess

The Two Main Arguments in Favor of Pulsars:

- The gamma-ray spectrum of observed pulsars
- Small-scale power in the gamma-ray emission from the Inner Galaxy


Arguments Against Pulsars:

- The measured luminosity function of gamma-ray pulsars
- The lack of low-mass X-ray binaries in the Inner Galaxy




Comparison With The Measured MSP Luminosity Function

- The MSP populations observed in globular clusters and in the galactic disk exhibit a gamma-ray luminosity function that is very different from that indicated by the analysis of Lee et al.
- The measured MSP luminosity function is very broad and extends over several orders of magnitude and up to at least $\sim 10^{35}$ erg/s
- If the small scale power identified by these analyses does in fact originate from MSPs, this is a very different population than those observed elsewhere



Millisecond Pulsars and Low-Mass X-Ray Binaries

- While a dead pulsar is being “spun-up” by a stellar companion to become a millisecond pulsar, it exists for a time as a low-mass X-ray binary (LMXB)
- We should expect the ratio of MSPs to LMXBs to be similar in the Inner Galaxy as in the Milky Way’s globular cluster population
- We can therefore use the number of low-mass X-ray binaries in the Inner Galaxy to estimate the population of MSPs that is present there



Millisecond Pulsars and Low-Mass X-Ray Binaries in Globular Clusters and in the Inner Galaxy

We compare the following data:

- 1) The gamma-ray emission from a sample of globular clusters selected for their high stellar encounter rates
- 2) The list of bright LMXBs found within the same Globular Clusters
- 3) The list of LMXB candidates observed in the Inner Galaxy

From this comparison, we conclude that an MSP population that is capable of generating the observed GeV excess should be accompanied by ~10 times more LMXBs than are observed

Globular Cluster	Flux (erg/cm ² /s)	Distance (kpc)	Stellar Encounter Rate	TS
NGC 104	$2.51_{-0.06}^{+0.05} \times 10^{-11}$	4.46	1.00	3995.9
NGC 362	$6.74_{-2.46}^{+2.63} \times 10^{-13}$	8.61	0.74	9.69
Palomar 2	$< 2.69 \times 10^{-13}$	27.11	0.93	0.0
NGC 6624	$1.14_{-0.10}^{+0.10} \times 10^{-11}$	7.91	1.15	455.8
NGC 1851	$9.05_{-2.67}^{+2.92} \times 10^{-13}$	12.1	1.53	14.4
NGC 5824	$< 4.78 \times 10^{-13}$	32.17	0.98	0.0

LMXB	Notes	Globular Cluster	References
4U 1820-30	P	NGC 6624	[69–71]
4U 0513-40	P	NGC 1851	[72–74]
4U 1746-37	P	NGC 6441	[69, 75, 76]
XB 1832-330	P	NGC 6652	[75, 77, 78]
M15 X-2	P	NGC 7078/M15	[79–81]
AC 211	P	NGC 7078/M15	[69, 80, 82]
SAX J1748.9-2021	T, XP	NGC 6440	[75, 83, 84]
GRS 1747-312	T	Terzan 6	[85–87]
Terzan 6 X-2	T	Terzan 6	[88]
IGR J17361-4441	T	NGC 6388	[89, 90]
IGR J18245-2542	T, XP	NGC 6626/M28	[91, 92]
EXO 1745-248	T	Terzan 5	[93, 94]
IGR J17480-2446	T	Terzan 5	[95–97]
Terzan 5 X-3	T	Terzan 5	[98]
MAXI J0911-635	T	NGC 2808	[99]

Gamma-Ray Bright MSPs in The Inner Galaxy?

- The most direct way to prove that MSPs generate the GeV excess would be to detect a significant number of individual pulsars in the Inner Galaxy

Gamma-Ray Bright MSPs in The Inner Galaxy?

- The most direct way to prove that MSPs generate the GeV excess would be to detect a significant number of individual pulsars in the Inner Galaxy
- Last year, the Fermi Collaboration posted a paper which purported to present strong evidence ($\sim 7\sigma$) for a large centrally located pulsar population


CHARACTERIZING THE POPULATION OF PULSARS IN THE GALACTIC BULGE WITH THE *FERMI* LARGE AREA TELESCOPE.

ABSTRACT

An excess of γ -ray emission from the Galactic Center (GC) region with respect to predictions based on a variety of interstellar emission models and γ -ray source catalogs has been found by many groups using data from the *Fermi* Large Area Telescope (LAT). Several interpretations of this excess have been invoked. In this paper we test the interpretation that the excess is caused by an unresolved population of γ -ray pulsars located in the Galactic bulge. We use cataloged LAT sources to derive criteria that efficiently select pulsars with very small contamination from blazars. We search for point sources in the inner $40^\circ \times 40^\circ$ region of the Galaxy, derive a list of approximately 400 sources, and apply pulsar selection criteria to extract pulsar candidates among our source list. We also derive the efficiency of these selection criteria for γ -ray pulsars as a function of source energy flux and location. We demonstrate that given the observed spatial and flux distribution of pulsar candidates, a model that includes a population with about 2.7 γ -ray pulsars in the Galactic disk (in our $40^\circ \times 40^\circ$ analysis region) for each pulsar in the Galactic bulge is preferred at the level of 7 standard deviations with respect to a disk-only model. The properties of these disk and bulge pulsar populations are consistent with the population of known γ -ray pulsars as well as with the spatial profile and energy spectrum of the GC excess. Finally, we show that the dark matter interpretation of the GC excess is strongly disfavored since a distribution of dark matter is not able to mimic the observed properties of the population of sources detected in our analysis.


Evidence of a Central Pulsar Population?

- In examining this paper, my collaborators and I found that we were unable to reproduce these results; our fit favored only a $\sim 2\sigma$ preference for a central source component
- As a result of the ensuing discussions with the Fermi Collaboration, an error was identified in their code, and a replacement (v2) of their paper was posted in conjunction with our paper



Evidence of a Central Pulsar Population?

- In our paper, we also note that masking the pulsar candidate sources contained in the 2FIG Fermi catalog does *not* impact the characteristics of the excess; **a negligible fraction of the excess emission originates from these sources**



A Series of Cosmic Ray Outbursts?

- Although the observe spectrum and morphology of the excess cannot be explained by ordinary stead-state diffuse emission models, non-steady state cosmic ray scenarios are more difficult to rule out – perhaps a recent series of burst-like events might be responsible?
- Hadronic scenarios predict a signal that is more diskly than spherical; highly incompatible with the data
- Leptonic scenarios, however, are more difficult to rule out




A Series of Cosmic Ray Outbursts?

After exploring a wide range of leptonic outburst scenarios, there appear to be two main challenges:

A Series of Cosmic Ray Outbursts?

After exploring a wide range of leptonic outburst scenarios, there appear to be two main challenges:


1) The morphology from a given outburst is “convex”, whereas the data is “concave” – to fit the data, we need several outbursts, with highly tuned parameters



A Series of Cosmic Ray Outbursts?

After exploring a wide range of leptonic outburst scenarios, there appear to be two main challenges:


1) The morphology from a given outburst is “convex”, whereas the data is “concave” – to fit the data, we need several outbursts, with highly tuned parameters



A Series of Cosmic Ray Outbursts?

After exploring a wide range of leptonic outburst scenarios, there appear to be two main challenges:

1) The morphology from a given outburst is “convex”, whereas the data is “concave” – to fit the data, we need several outbursts, with highly tuned parameters




A Series of Cosmic Ray Outbursts?

After exploring a wide range of leptonic outburst scenarios, there appear to be two main challenges:


1) The morphology from a given outburst is “convex”, whereas the data is “concave” – to fit the data, we need several outbursts, with highly tuned parameters

2) The gamma-ray spectrum is approximately uniform across the Inner Galaxy, but energy losses should lead to softer emission from the outer regions – to fit the data, we need the older outbursts to inject electrons with higher energies than more recent outbursts




Testing Dark Matter Interpretations

- Searches for gamma rays from dwarf galaxies with Fermi and measurements of the cosmic-ray antiproton spectrum by AMS are each potentially sensitive to dark matter with the characteristics needed to account for the observed gamma-ray excess




Cuoco, et al., arXiv:1610.03071
 Cui, et al. arXiv:1610.03840



Fermi Collaboration, arXiv:1611.03184

Fermi Observations of Dwarf Galaxies


- Current Fermi dwarf constraints are based on stacks of dwarf galaxy candidates, aided by recent discoveries by DES and other surveys
- At this time, these constraints are compatible with dark matter interpretations of the Galactic Center excess
- That being said, if the Galactic Center signal is coming from annihilating dark matter, we should expect gamma rays to be detected from dwarfs soon




Fermi Collaboration, arXiv:1611.03184
(see also 1503.02641)

Fermi's Observations of Dwarf Galaxies

Intriguing (but inclusive) excesses have been observed from the recently discovered and nearby dwarf galaxies Reticulum II and Tucana III (Geringer-Sameth et al., Drlica-Wagner, et al., DH & Linden)




The plot I see in my dreams...



Annihilating Dark Matter And Cosmic-Ray Antiprotons

- In the AMS antiproton spectrum, there is a small excess (relative to standard secondary production) at $R \sim 10\text{-}20$ GV
- The excess is well fit by a $\sim 50\text{-}90$ GeV dark matter particle with an annihilation cross section of $\sim 10^{-26}$ cm³/s (for bb), in good agreement with the Galactic Center excess
- Although statistically significant at face value ($\sim 4.5\sigma$), the systematics associated with the antiproton production cross section and the effects of solar modulation are difficult to quantify



Cuoco, et al., arXiv:1610.03071

Cui, et al., arXiv:1610.03840

Reinert, Winkler, arXiv:1712.00002

Cui, et al., arXiv:1803.02163

Summary

- The Galactic Center's GeV excess remains compelling: highly statistically significant, robust, extended, and difficult to explain with known or proposed astrophysics
- Although millisecond pulsars could be responsible for this gamma-ray excess, the Inner Galaxy population would have to be quite different from those observed in the disk of the Milky Way and in the Milky Way's globular cluster population (strongly peaked luminosity function, accompanied by fewer LMXBs)
- Gamma-ray (and radio) searches for millisecond pulsars in the Inner Galaxy have not yet found any evidence for such sources; sub-threshold searches have yielded results that are open to multiple interpretations
- The modest excesses observed from dwarf galaxies and in the cosmic-ray antiproton spectrum are suggestive

