



Neutrinos and Gamma Rays from Extragalactic Sources

Markus Ahlers, Niels Bohr Institute

TeVPA 2018, August 28, Berlin


Multi-Messenger Astronomy

- **Cosmic ray (CR)** acceleration in the aftermath of cataclysmic events, sometimes seen in **gravitational waves**.
- Inelastic collisions with radiation or gas produce **γ -rays** and **neutrinos**, e.g.

$$\pi^0 \rightarrow \gamma + \gamma$$

$$\pi^+ \rightarrow \mu^+ + \nu_\mu \rightarrow e^+ + \nu_e + \bar{\nu}_\mu + \nu_\mu$$


- **Unique aspects of neutrino messengers:**
 - *identify* cosmic ray sources
 - *qualifies* γ -ray emission
 - *covers blind spot* of astronomy to the very-high-energy Universe




2013: A Milestone for Neutrino Astronomy

First observation of high-energy astrophysical neutrinos by IceCube!

“track event” (from ν_μ scattering)




“cascade event” (from all flavours)



[“Breakthrough of the Year” (Physics World), Science 2013]
(time-dependent neutrino signal: **early** to **late** light detection)


Cosmic TeV-PeV Neutrinos

- **High-Energy Starting Events (HESE) (6yrs):** [Science 342 (2013); update ICRC 2017]
 - bright events ($E_{\text{th}} \gtrsim 30\text{TeV}$) starting inside IceCube
 - efficient removal of atmospheric backgrounds by veto layer
- **Up-going muon-neutrino tracks (8yrs):** [Astrophys.J. 833 (2016); update ICRC 2017]
 - large effective volume due to ranging in tracks
 - efficient removal of atmospheric muon backgrounds by Earth-absorption



Multi-Messenger Interfaces

- **High-Energy Starting Events (HESE) (6yrs):** [Science 342 (2013); update ICRC 2017]
 - bright events ($E_{\text{th}} \gtrsim 30\text{TeV}$) starting inside IceCube
 - efficient removal of atmospheric backgrounds by veto layer
- **Up-going muon-neutrino tracks (8yrs):** [Astrophys.J. 833 (2016); update ICRC 2017]
 - large effective volume due to ranging in tracks
 - efficient removal of atmospheric muon backgrounds by Earth-absorption



Power-Law Fits

- power-law fit (per flavour):

$$\phi(E) = \frac{\phi_{\text{astro}} \times 10^{-8}}{\text{GeV cm}^2 \text{s sr}} \left[\frac{E}{100 \text{TeV}} \right]^{-\gamma_{\text{astro}}}$$


- HESE (6yr) fit range:

$$60 \text{TeV} \leq E \leq 3 \text{PeV}$$

- up-going $\nu_\mu + \bar{\nu}_\mu$ (8yr) fit range:


$$119 \text{TeV} \leq E \leq 4.8 \text{PeV}$$

- Hard spectrum of 2-component HESE fit consistent with $\nu_\mu + \bar{\nu}_\mu$ spectrum within 68% C.L.!**




The Cosmic “Beam”

1 PeV neutrino \leftrightarrow 20-30 PeV cosmic ray nucleon



Arrival Directions of Cosmic Neutrinos

Arrival directions of most energetic neutrino events (HESE 6yr (magenta) & $\nu_\mu + \bar{\nu}_\mu$ 8yr (red))




No significant correlation of diffuse flux with known Galactic or extragalactic sources.

Extragalactic Source Candidates


- association with sources of UHE CRs [Kistler, Stanev & Yuksel'13]
[Katz, Waxman, Thompson & Loeb'13; Fang, Fujii, Linden & Olinto'14; Moharana & Razzaque'15]
- association with diffuse γ -ray background [Murase, MA & Lacki'13]
[Chang & Wang'14; Ando, Tamborra & Zandanel'15]
- active galactic nuclei (AGN) [Stecker'13; Kalashev, Kusenko & Essey'13]
[Murase, Inoue & Dermer'14; Kimura, Murase & Toma'14; Kalashev, Semikoz & Tkachev'14]
[Padovani & Resconi'14; Petropoulou *et al.*'15; Padovani *et al.*'16; Kadler *et al.*'16; Wang & Loeb'16]
- gamma-ray bursts (GRB) [Murase & Ioka'13; Dado & Dar'14; Tamborra & Ando'15]
[Senno, Murase & Meszaros'16; Denton & Tamborra'18; Boncioli, Biehl & Winter'18]
- galaxies with intense star-formation (e.g. starbursts)
[He, Wang, Fan, Liu & Wei'13; Yoast-Hull, Gallagher, Zweibel & Everett'13; Murase, MA & Lacki'13]
[Anchordoqui, Paul, da Silva, Torres & Vlcek'14; Tamborra, Ando & Murase'14; Chang & Wang'14]
[Liu, Wang, Inoue, Crocker & Aharonian'14; Senno, Meszaros, Murase, Baerwald & Rees'15]
[Chakraborty & Izaguirre'15; Emig, Lunardini & Windhorst'15; Bechtol *et al.*'15]
- galaxy clusters/groups [Murase, MA & Lacki'13; Zandanel, Tamborra, Gabici & Ando'14]
- tidal disruption events (TDE) [Wang, Liu, Dai & Cheng'11; Senno, Murase & Més'aros'17]
[Guépin, Kotera, Barausse, Fang & Murase'17; Biehl, Boncioli, Lunardini & Winter'17]

Neutrino Point-Source Limits



[IceCube, *Astrophys.J.* 835 (2017) no.2, 151]

Extragalactic Sources



- Low neutrino absorption in the Universe allows to observe distant sources.
- Quasi-diffuse flux observed by IceCube is composed of **many individual sources**.
- Can they be identified?


lower density (ρ)
↓
higher luminosity (L)
↓
brighter sources (ϕ)

Constraints from Point-Source Limits

→ **Populations with lower density require higher luminosity and predict brighter individual sources.**

- **non-observation** of individual neutrino sources constrain source populations, e.g.


- ✗ gamma-ray bursts
 $(\rho_{\text{eff}} \simeq \text{Gpc}^{-3} \text{yr}^{-1})$
- ✗ BL Lacs / FSRQ
 $(\rho_{\text{eff}} \simeq 0.1 - 10 \text{ Gpc}^{-3})$



[MA & Halzen'18]

Gamma-Ray Bursts


- Neutrino production at various stages of a gamma-ray burst (GRB).
 - **precursor** $p\bar{p}$ and $p\gamma$ interactions in stellar envelope;
also possible for “failed” GRBs [Razzaque, Meszaros&Waxman'03]
 - **burst** $p\gamma$ interactions in internal shocks [Waxman&Bahcall'97]
 - **afterglow** $p\gamma$ interactions in reverse external shocks [Waxman&Bahcall'00; Murase&Nagataki'06; Murase'07]



[Meszaros'01]


Gamma-Ray Bursts

- strong limits on neutrino emission associated with “fireball” model [Abbasi et al.’12]
- PeV neutrino flux exceeds GRB limit by one order of magnitude.



Low-Luminosity Gamma-ray Bursts


- *loop-hole*: undetected low-luminosity γ -ray bursts (GRB)
[Murase & Ioka'13; Senno, Murase & Mészáros'16; Boncioli, Biehl & Winter'18]
- *claim*: distinct population of LL-GRB more abundant in the local ($z \ll 1$) Universe




[Liang, Zhang, Virgili & Dai'06]

Active Galactic Nuclei / Blazars


- neutrinos from $p\gamma$ interactions in AGN [Steckeret al.'91; Mannheim'96; Halzen & Zas'97]
- complex spectra due to various photon backgrounds
- typically, deficit of sub-PeV and excess of EeV neutrinos



[Murase, Inoue & Dermer'14]




TeV Gamma-Ray Blazars




Blazar stacking limits derived from Fermi-LAT AGN catalogue (2LAC) [IceCube'16]


TeV Gamma-Ray Blazars




(a) generic blazars



(b) BL Lacs




(c) FSRQs - 1



(d) FSRQs - 2


Blazar stacking limits derived from Fermi-LAT AGN catalogue (2LAC) [IceCube'16]

Candidate Neutrino Source: TXS 0506+056



EM flare


[Science 361 (2018) no.6398, eaat1378]




neutrino flare

[Science 361 (2018) no.6398, 147-151]

Fits of Multi-Messenger SED



[Keivani et al., arXiv:1807.04537]



[Gao et al., arXiv:1807.04275]

- Photon SED can be modelled with lepto-hadronic and proton-synchrotron models.
 [see also Cerruti et al. arXiv:1807.04335; Zhang, Fang & Li, arXiv:1807.11069]
 [Gokus et al. arXiv:1808.05540; Sahakyan, arXiv:1807.05651]
- Neutrino flux limited by theoretically feasible proton luminosity and X-ray data.
 [see, however, Righi, Tavecchio & Inoue, arXiv:1807.11069]
 [Murase, Oikonomou & Petropoulou, arXiv:1807.04748; Liu et al., arXiv:1807.05113]

Hadronic Gamma-Ray Emission

- Inelastic collisions of **cosmic rays (CR)** with radiation or gas produce **γ -rays** and **neutrinos** via pion decay:

$$\pi^0 \rightarrow \gamma + \gamma$$

$$\pi^+ \rightarrow \mu^+ + \nu_\mu \rightarrow e^+ + \nu_e + \bar{\nu}_\mu + \nu_\mu$$


- relative production rates:

$$\frac{1}{3} \sum_{\alpha} E_{\nu}^2 Q_{\nu_{\alpha}}(E_{\nu}) \simeq \frac{K_{\pi}}{4} \left[E_{\gamma}^2 Q_{\gamma}(E_{\gamma}) \right]_{E_{\gamma}=2E_{\nu}}$$

- ✗ TeV γ -rays scatter in cosmic microwave background (CMB) and initiate electromagnetic cascades:

$$\gamma + \gamma_{\text{CMB}} \rightarrow e^+ + e^-$$

$$e^{\pm} + \gamma_{\text{CMB}} \rightarrow e^{\pm} + \gamma$$



Isotropic Diffuse Gamma-Ray Background (IGRB)

- Gamma-ray emission from electromagnetic cascades ends up in the sub-TeV range observed with Fermi satellite.


- ✗ Cosmic ray spectral index **strongly constrained** by the isotropic diffuse gamma-ray background (IGRB)

[Murase, MA & Lacki'13]

$$\Gamma \lesssim 2.15 - 2.2$$

- ✗ IceCube best-fit: [IceCube'15]

$$\Gamma \simeq 2.4 - 2.6$$



[Murase, MA & Lacki'14; Tamborra, Ando & Murase'14]

[Ando, Tamborra & Zandanel'15]

[Bechtol, MA, Ajello, Di Mauro & Vandenbergroucke'15]

Isotropic Diffuse Gamma-Ray Background (IGRB)

- Gamma-ray emission from electromagnetic cascades ends up in the sub-TeV range observed with Fermi satellite.


- ✗ Cosmic ray spectral index **strongly constrained** by the isotropic diffuse gamma-ray background (IGRB)

[Murase, MA & Lacki'13]

$$\Gamma \lesssim 2.15 - 2.2$$

- ✗ IceCube best-fit: [IceCube'15]

$$\Gamma \simeq 2.4 - 2.6$$



[Murase, MA & Lacki'14; Tamborra, Ando & Murase'14]

[Ando, Tamborra & Zandanel'15]

[Bechtol, MA, Ajello, Di Mauro & Vandenbergroucke'15]

Isotropic Diffuse Gamma-Ray Background (IGRB)

- Gamma-ray emission from electromagnetic cascades ends up in the sub-TeV range observed with Fermi satellite.


- ✗ Cosmic ray spectral index **strongly constrained** by the isotropic diffuse gamma-ray background (IGRB)

[Murase, MA & Lacki'13]

$$\Gamma \lesssim 2.15 - 2.2$$

- ✗ IceCube best-fit: [IceCube'15]

$$\Gamma \simeq 2.4 - 2.6$$



[Murase, MA & Lacki'14; Tamborra, Ando & Murase'14]

[Ando, Tamborra & Zandanel'15]

[Bechtol, MA, Ajello, Di Mauro & Vandenbergroucke'15]

Non-Blazar Limits on Gamma-Ray Background

- **Photon fluctuation analyses** of Fermi data allow to constrain the source count distribution of blazars **below** the source detection threshold.

- inferred blazar contribution above 50 GeV:

- Fermi Collaboration'15:


$86^{+16}_{-14}\%$ of EGB

- Lisanti *et al.*'16:

$68^{+9}_{-8}(\pm 10)_{\text{sys}}\%$ of EGB

- Zechlin *et al.*'16


$81^{+52}_{-19}\%$ of EGB



[Fermi'15]

Non-Blazar Limits on Gamma-Ray Background

- non-blazar contribution above 50 GeV:
[Fermi'15]
 $14^{+14\%}_{-14\%}$ of EGB
- ✗ **strong tension** with IceCube observation ($E_\nu \lesssim 100$ TeV)
- limits apply to generic **cosmic ray calorimeters**
- *possible loop-holes:*
 - γ -absorption in source?
[Chang & Wang'14]
 - suppression of γ -ray cascades?
[Broderick, Chang & Pfrommer'12]



Fermi Bounds for $p\gamma$ Sources

- Fermi constraints less severe for $p\gamma$ scenarios:

- 1 **no power-law extrapolation** to Fermi energy range
- 2 **high pion production efficiency** implies strong γ -absorption in sources

- source candidates:


- AGN cores [Stecker'91; '13]

[Kimura, Murase & Toma'14]

- choked GRB jets

[Mészáros & Waxman'01]

[Senno, Murase & Mészáros'16]



Corresponding Opacities

- required cosmic ray energy:

$$E_{\text{CR}} \sim 20 E_\nu$$

- required target photon energy:


$$\varepsilon_t \sim 200 \text{ keV} \left(\frac{\Gamma}{10} \right)^2 \left(\frac{E_\nu}{3 \text{ TeV}} \right)^{-1}$$

- opacity relation:

$$\tau_{\gamma\gamma}(E_\gamma) \sim 1000 f_{p\gamma}(E_p)$$

- strong internal γ -absorption:

$$E_\gamma \gtrsim 100 \text{ MeV} \left(\frac{E_\nu}{3 \text{ TeV}} \right)$$



[Murase, Guetta & MA'15]

UHE CR association?

- UHE CR proton emission rate density:

[e.g. MA & Halzen'12]

$$[E_p^2 Q_p(E_p)]_{10^{19.5}\text{eV}} \simeq 8 \times 10^{43} \text{ erg Mpc}^{-3} \text{ yr}^{-1}$$

- corresponding per flavor neutrino flux ($\xi_z \simeq 0.5 - 2.4$ and $K_\pi \simeq 1 - 2$):

$$E_\nu^2 \phi_\nu(E_\nu) \simeq f_\pi \underbrace{\frac{\xi_z K_\pi}{1 + K_\pi}}_{\mathcal{O}(1)} \underbrace{1.5 \times 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}}_{\sim \text{IceCube diffuse}}$$

- **Waxman-Bahcall bound:** $f_\pi \leq 1$

[Waxman & Bahcall'98]

- **similar** UHE nucleon emission rate density (local minimum at $\Gamma \simeq 2.04$) [Auger'16]

$$[E_N^2 Q_N(E_N)]_{10^{19.5}\text{eV}} \simeq 2.2 \times 10^{43} \text{ erg Mpc}^{-3} \text{ yr}^{-1}$$

✗ **But**, how to reach $E_{\max} \simeq 10^{20}$ eV in environments of high energy loss ($f_\pi \simeq 1$)?


UHE CR association?

→ two-zone models: CR accelerator + CR “calorimeter”?


- starburst galaxies
- galaxy clusters
- “unified” sources (UHE CRs, γ -ray & neutrinos):

[Loeb & Waxman'06]

[Berezinsky, Blasi & Ptuskin'96; Beacom & Murase'13]



[Kachelriess, Kalashev, Ostapchenko & Semikoz'17]




[Fang & Murase'17]

✗ However, $E_\nu < 100$ TeV neutrino data remains a challenge!

Correlation with UHE CRs?

Auger 2014 $E \geq 52$ EeV (\times) / TA 2014 $E \geq 57$ EeV (+) / smoothed anisotropy map ($\Delta\theta_{50\%} = 15^\circ$)




- $\theta_{\text{rms}} \simeq 1^\circ (D/\lambda_{\text{coh}})^{1/2} (E/55\text{EeV})^{-1} (\lambda_{\text{coh}}/1\text{Mpc}) (B/1\text{nG})$ [Waxman & Miralda-Escude'96]
- "hot spots" (dashed), but no significant auto-correlation in Auger and Telescope Array data

Summary

- IceCube has identified a **diffuse flux of astrophysical neutrinos** in the TeV-PeV energy range of **unknown origin**.
 - Galactic and Extragalactic Sources are candidate sources, but **absence of anisotropies** favours the latter.
 - **No compelling scenario** for the TeV-PeV energy range.
 - **High intensity** of the emission is comparable to that of ultrahigh-energy cosmic rays and γ -ray backgrounds.
- Excellent conditions for **multi-messenger studies**:
- Large neutrino flux in the 1 – 10 TeV range is challenged by constraints set by the **extragalactic γ -ray background** observed by Fermi.
 - New candidate sources **TXS 0506+056** for neutrino/ γ -ray emission.
 - Saturation of calorimetric bounds of **UHE CR sources** might indicate common origin.

Appendix

Cosmic Ray Interactions



[data from PDG (<http://pdg.lbl.gov>)]

Neutrinos from Pion Decay

- Neutrinos from pion and muon decay:

$$\pi^+ \rightarrow \mu^+ + \nu_\mu$$

$$\mu^+ \rightarrow e^+ + \nu_e + \bar{\nu}_\mu$$

- average energy fraction from relativistic pions ($r_\pi \equiv m_\mu^2/m_\pi^2 \simeq 0.57$):


$$\langle x \rangle_{\pi^+ \rightarrow \nu_\mu} = \frac{1 - r_\pi}{2} \simeq 21\%$$

$$\langle x \rangle_{\pi^+ \rightarrow \nu_\mu} = \frac{3 + 4r_\pi}{20} \simeq 26\%$$

$$\langle x \rangle_{\pi^+ \rightarrow \nu_e} = \frac{2 + r_\pi}{10} \simeq 26\%$$

- In practice, we often use the **approximation**:

$$\langle x \rangle_{\nu_x} \simeq \langle x \rangle_{\bar{\nu}_x} \simeq \frac{1}{4} \quad \& \quad \kappa_\pi \simeq \frac{1}{5} \quad \rightarrow \quad \frac{\langle E_\nu \rangle}{E_N} \simeq \frac{1}{20}$$



Galactic Source Candidates

- diffuse Galactic γ -ray emission [MA & Murase'13; Joshi J C, Winter W and Gupta'13]
[Kachelriess and Ostapchenko'14; Neronov, Semikoz & Tchernin'13; Neronov & Semikoz'14,'16]
[Guo, Hu & Tian'14; Gaggero, Grasso, Marinelli, Urbano & Valli'15; Neronov, Kachelriess & Semikoz'18]
- unidentified Galactic γ -ray emission [Fox, Kashiyama & Meszaros'13]
[Gonzalez-Garcia, Halzen & Niro'14]
- *Fermi Bubbles* [MA & Murase'13; Razzaque'13]
[Lunardini, Razzaque, Theodoreau & Yang'13; Lunardini, Razzaque & Yang'15]
- supernova remnants [Mandelartz & Tjus'14]
- pulsars [Padovani & Resconi'14]
- microquasars [Anchordoqui, Goldberg, Paul, da Silva & Vlcek'14]
- Sagittarius A* [Bai, Barger, Barger, Lu, Peterson & Salvado'14; Fujita, Kimura & Murase'15,'16]
- Galactic Halo [Taylor, Gabici & Aharonian'14]
- heavy dark matter decay [Feldstein, Kusenko, Matsumoto & Yanagida'13]
[Esmaili & Serpico '13; Bai, Lu & Salvado'13; Cherry, Friedland & Shoemaker'14]
[Murase, Laha, Ando, MA'15; Boucenna *et al.*'15 ; Chianese, Miele, Morisi & Vitagliano'16]

Pion Production Efficiency

- pion production depend on target opacity $\tau = \ell\sigma n$
- “bolometric” pion production efficiency (inelasticity κ):

$$f_\pi = 1 - \exp(-\kappa\tau)$$

- inelasticity per pion : $\kappa_\pi = \kappa / \langle N_{\text{all } \pi} \rangle \simeq 0.17 - 0.2$
- “bolometric” relation of the production rates Q :

$$E_\pi^2 Q_{\pi^\pm}(E_\pi) \simeq \frac{\langle N_{\pi^+} \rangle + \langle N_{\pi^-} \rangle}{\langle N_{\pi^0} \rangle + \langle N_{\pi^+} \rangle + \langle N_{\pi^-} \rangle} \left[f_\pi E_N^2 Q_N(E_N) \right]_{E_N=E_\pi/\kappa_\pi}$$

- charged-to-neutral pion ratio:

$$K_\pi \equiv \frac{\langle N_{\pi^+} \rangle + \langle N_{\pi^-} \rangle}{\langle N_{\pi^0} \rangle} \simeq \begin{cases} 2 & pp \\ 1 & p\gamma \end{cases}$$

- or in more compact form with K_π :

$$E_\pi^2 Q_{\pi^\pm}(E_\pi) \simeq f_\pi \frac{K_\pi}{1 + K_\pi} \left[E_N^2 Q_N(E_N) \right]_{E_N=E_\pi/\kappa_\pi}$$

Neutrino and Gamma-Ray Emission

- neutrino emission from pion decay

$$\frac{1}{3} \sum_{\alpha} E_{\nu} Q_{\nu_{\alpha}}(E_{\nu}) \simeq [E_{\pi} Q_{\pi^{\pm}}(E_{\pi})]_{E_{\pi} \simeq 4E_{\nu}} \simeq \frac{1}{4} f_{\pi} \frac{K_{\pi}}{1 + K_{\pi}} \left[E_N^2 Q_N(E_N) \right]_{E_N = 4E_{\nu}/\kappa_{\pi}}$$

- neutrino and γ -ray emission are related as


$$\frac{1}{3} \sum_{\alpha} E_{\nu} Q_{\nu_{\alpha}}(E_{\nu}) \simeq \frac{1}{2} \frac{\langle N_{\pi^+} \rangle + \langle N_{\pi^-} \rangle}{\langle N_{\pi^0} \rangle} [E_{\gamma} Q_{\gamma}(E_{\gamma})]_{E_{\gamma} = 2E_{\nu}}$$

- again, a more compact form with K_{π} :

$$\frac{1}{3} \sum_{\alpha} E_{\nu}^2 Q_{\nu_{\alpha}}(E_{\nu}) \simeq \frac{K_{\pi}}{4} \left[E_{\gamma}^2 Q_{\gamma}(E_{\gamma}) \right]_{E_{\gamma} = 2E_{\nu}}$$

- γ -ray emission is attenuated in sources and, in particular, in the extragalactic radiation background

Methods of Neutrino Detection I




→ Selecting **up-going muon tracks** reduces atmospheric muon background:

$$\underbrace{10,000,000,000}_{\text{atmospheric muons (from above)}} : \underbrace{100,000}_{\text{atmospheric neutrinos}} : \underbrace{10}_{\text{cosmic neutrinos}}$$

Methods of Neutrino Detection II

- Outer layer of optical modules can be used as a **veto region** (gray area):
 - ✗ **Atmospheric muons** pass through veto from above.
 - ✗ **Atmospheric neutrinos** are produced in coincidence with atmospheric muons.
 - ✓ **Cosmic neutrino** events can start inside the fiducial volume.
 - **High-Energy Starting Event** (HESE) analysis



[IceCube Collaboration'13]

Ultra-Long Baseline Oscillations


- Energy resolution of detectors is limited and size of cosmic neutrino source is large.

$$P_{\nu_\alpha \rightarrow \nu_\beta} = \delta_{\alpha\beta} - 4 \sum_{i>j} \Re(U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^*) \underbrace{\sin^2 \Delta_{ij}}_{\rightarrow 1/2} + 2 \sum_{i>j} \Im(U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^*) \underbrace{\sin 2\Delta_{ij}}_{\rightarrow 0}$$

→ oscillation-averaged probability:

$$P_{\nu_\alpha \rightarrow \nu_\beta} \simeq \sum_i |U_{\alpha i}|^2 |U_{\beta i}|^2$$

- initial composition: $\nu_e : \nu_\mu : \nu_\tau$
pion & muon decay: $1 : 2 : 0$
muon-damped decay: $0 : 1 : 0$
neutron decay: $1 : 0 : 0$



Ultra-Long Baseline Oscillations


- Energy resolution of detectors is limited and size of cosmic neutrino source is large.

$$P_{\nu_\alpha \rightarrow \nu_\beta} = \delta_{\alpha\beta} - 4 \sum_{i>j} \Re(U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^*) \underbrace{\sin^2 \Delta_{ij}}_{\rightarrow 1/2} + 2 \sum_{i>j} \Im(U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^*) \underbrace{\sin 2\Delta_{ij}}_{\rightarrow 0}$$

→ oscillation-averaged probability:


$$P_{\nu_\alpha \rightarrow \nu_\beta} \simeq \sum_i |U_{\alpha i}|^2 |U_{\beta i}|^2$$

- initial composition: $\nu_e : \nu_\mu : \nu_\tau$
pion & muon decay: $1 : 2 : 0$
muon-damped decay: $0 : 1 : 0$
neutron decay: $1 : 0 : 0$




[Astrophys.J. 809 (2015) no.1, 98]

Non-Anthropogenic Neutrino Fluxes




Non-Anthropogenic Neutrino Fluxes



Cosmogenic (“GZK”) Neutrinos


- Observation of UHE CRs and extragalactic radiation backgrounds “guarantee” a flux of high-energy neutrinos, in particular via resonant production in CMB.
[Berezinsky & Zatsepin'69]
- “Guaranteed”, but with many model uncertainties and constraints:
 - (**low cross-over**) **proton models + CMB (+ EBL)**
[Berezinsky & Zatsepin'69; Yoshida & Teshima'93; Protheroe & Johnson'96; Engel, Seckel & Stanev'01; Fodor, Katz, Ringwald & Tu'03; Barger, Huber & Marfatia'06; Yuksel & Kistler'07; Takami, Murase, Nagataki & Sato'09, MA, Anchordoqui & Sarkar'09, Heinz, Boncioli, Bustamante & Winter'15]
 - **+ mixed compositions**
[Hooper, Taylor & Sarkar'05; Ave, Busca, Olinto, Watson & Yamamoto'05; Allard, Ave, Busca, Malkan, Olinto, Parizot, Stecker & Yamamoto'06; Anchordoqui, Goldberg, Hooper, Sarkar & Taylor'07; Kotera, Allard & Olinto'10; Decerprit & Allard'11; MA & Halzen'12]
 - **+ extragalactic γ -ray background limits**
[Berezinsky & Smirnov'75; Mannheim, Protheroe & Rachen'01; Keshet, Waxman, & Loeb'03; Berezinsky, Gazizov, Kachelriess & Ostapchenko'10; MA, Anchordoqui, Gonzalez-Garcia, Halzen & Sarkar'10; MA & Salvado'11; Gelmini, Kalashev & Semikoz'12]

Limits on Cosmogenic Neutrinos




[Phys.Rev.Lett. 117 (2016) 241101]

- Upper limits on cosmogenic (top left) and astrophysical (bottom left) neutrino emission models.
- Differential upper limits (right) in comparison with Auger and ANITA.
- **Proton-dominated cosmogenic neutrino models are disfavoured.**



Starburst Galaxies

- **Increased star formation** enhances cosmic ray production.
- **Dense environment and strong magnetic fields** enhance CR containment and interaction.
- Expect spectral break at (0.1 – 1) PeV from CR leakage (“CR knee”).
- Plot shows muon neutrinos on production (3/2 of total neutrino flux).




TeV Starburst Galaxies

Messier 82 ($\delta \simeq 69^\circ$)



NGC 253 ($\delta \simeq -25^\circ$)



$$E^2 \phi_\gamma(E) \simeq 3.3 \times 10^{-13} \left(\frac{E}{\text{TeV}} \right)^{-0.5} \frac{\text{TeV}}{\text{cm}^2\text{s}}$$

$$E^2 \phi_\gamma(E) \simeq 9.6 \times 10^{-13} \left(\frac{E}{\text{TeV}} \right)^{-0.14} \frac{\text{TeV}}{\text{cm}^2\text{s}}$$

$$E^2 \phi_\nu(E) \lesssim 1.09 \times 10^{-12} \frac{\text{TeV}}{\text{cm}^2\text{s}}$$



[IceCube 7yr $\nu_\mu + \bar{\nu}_\mu$]

no neutrino limit

expected from CR-gas interactions: $E_\nu^2 \phi_{\nu_\mu}(E_\nu) \simeq \frac{1}{2} E_\gamma^2 \phi_\gamma(E_\gamma)$

Tidal Disruption Events

- Stars torn apart by tidal forces in the vicinity of a supermassive black hole can launch jet-like outflows.
- good candidate sources of UHE CRs [Farrar & Gruzinov'09; Farrar & Piran'14]
- associate neutrino production via $p\gamma$ interactions:
[Wang, Liu, Dai & Cheng'11; Senno, Murase & Més'aros'17]
[Guépin, Kotera, Barausse, Fang & Murase'17; Biehl, Boncioli, Lunardini & Winter'17]



[e.g. Biehl, Boncioli, Lunardini & Winter'17]