

3D Topological Reconstruction for the JUNO Detector

The precise reconstruction of charged particle tracks in homogeneous liquid scintillator (LSc) neutrino detectors, e.g., from muons, is an important prerequisite for the efficient rejection of cosmogenic background events or the analysis of multi-GeV neutrino interactions. Topological information on such events, i.e., the reconstructed 3D density distribution of isotropically emitted scintillation photons, opens up new ways to accomplish these tasks. Especially future multi-kiloton LSc detectors like JUNO will profit from improved (muon) track reconstruction possibilities, both regarding their low- and high-energy neutrino physics programs.

Furthermore, the method presented here can also give valuable information on events traditionally thought as point-like (MeV). This offers the potential for particle discrimination at energies relevant for solar- or reactor-neutrino programs. First results towards this goal using the JUNO-MC environment are presented here.

Authorship annotation

B. Wonsak, C. Hagner for the JUNO collaboration

Session and Location

Monday Session, Poster Wall #177 (Ballroom)

Poster included in proceedings:

yes

Primary author: WONSAK, Björn (UniHH Opera)

Co-author: Prof. HAGNER, Caren (Universität Hamburg)

Presenter: WONSAK, Björn (UniHH Opera)

Track Classification: Poster (participating in poster prize competition)