

Mass Degeneracies in Extended Higgs Sectors

Howard E. Haber
22 May 2018



Outline

1. Questions for extended Higgs sectors with mass-degenerate scalars
2. A warmup: Natural mass degeneracies in the 2HDM
 - Neutral scalar mass degeneracies in the inert doublet model (IDM)
3. New features of mass degenerate scalars in the 3HDM
 - The replicated IDM
 - The Ivanov-Silva model (and the significance of CP4)
4. Final comments

based on: H.E. Haber, O.M. Ogreid, P. Osland and M.N. Rebelo, arXiv:180x.yyyyyy.

Mass Degeneracies in Extended Higgs Sectors

We would like to explore the possibility of mass-degenerate neutral scalars and/or mass-degenerate charged Higgs pairs that could arise in extended Higgs sectors. In each case, the critical questions to ask are:

- Is the origin of the mass degeneracy natural? (Yes, if due to a symmetry. No, if accidental)
- Can mass degenerate scalars be distinguished experimentally on an event by event basis?
- Is the only signal of the mass degeneracy a measurable multiplicity factor that arises when averaging over initial state degeneracies and summing over final state degeneracies?

Natural scalar mass degeneracies in the 2HDM

Consider the 2HDM with two hypercharge-one, doublet scalar fields. It is convenient to work in the Higgs basis in which the two Higgs doublet fields, denoted by H_1 and H_2 , satisfy $\langle H_1^0 \rangle = v/\sqrt{2}$ and $\langle H_2^0 \rangle = 0$ (i.e., the vacuum expectation value, $v = 246$ GeV, resides entirely in the neutral component of the Higgs basis field H_1 .)

We can immediately identify the physical charged Higgs field, $H^+ \equiv H_2^+$, and the neutral and charged Goldstone fields, $G^0 = \sqrt{2} \text{Im } H_1^0$ and $G^+ \equiv H_1^+$. In the Higgs basis, the scalar potential is given by:

$$\begin{aligned} \mathcal{V} = & Y_1 H_1^\dagger H_1 + Y_2 H_2^\dagger H_2 + [Y_3 H_1^\dagger H_2 + \text{h.c.}] + \frac{1}{2} Z_1 (H_1^\dagger H_1)^2 \\ & + \frac{1}{2} Z_2 (H_2^\dagger H_2)^2 + Z_3 (H_1^\dagger H_1) (H_2^\dagger H_2) + Z_4 (H_1^\dagger H_2) (H_2^\dagger H_1) \\ & + \left\{ \frac{1}{2} Z_5 (H_1^\dagger H_2)^2 + [Z_6 (H_1^\dagger H_1) + Z_7 (H_2^\dagger H_2)] H_1^\dagger H_2 + \text{h.c.} \right\}, \end{aligned}$$

where Y_1 , Y_2 and $Z_{1,2,3,4}$ are real, whereas Y_3 , $Z_{5,6,7}$ are potentially complex. After minimizing the scalar potential, $Y_1 = -\frac{1}{2} Z_1 v^2$ and $Y_3 = -\frac{1}{2} Z_6 v^2$.

Specializing to the Inert doublet model (IDM)

Suppose that the Higgs basis of the 2HDM exhibits an exact \mathbb{Z}_2 symmetry, $H_1 \rightarrow +H_1$ and $H_2 \rightarrow -H_2$. This symmetry is also preserved by the vacuum. It then follows that $Y_3 = Z_6 = Z_7 = 0$. The one remaining complex parameter, Z_5 can be chosen real by rephasing the Higgs basis field H_2 . Thus, the IDM scalar potential is CP-conserving.

The Higgs basis doublet fields are also mass eigenstate fields,

$$H_1 = \begin{pmatrix} G^+ \\ \frac{1}{\sqrt{2}}[v + h + iG^0] \end{pmatrix}, \quad H_2 = \begin{pmatrix} H^+ \\ \frac{1}{\sqrt{2}}[H + iA] \end{pmatrix},$$

where G^\pm and G^0 are the Goldstone bosons that provide the longitudinal degrees of freedom of the massive W^\pm and Z^0 gauge bosons. The tree-level properties of the scalar h are precisely those of the SM Higgs boson. The physical scalar mass spectrum is,

$$\begin{aligned} m_h^2 &= Z_1 v^2, & m_{H^\pm}^2 &= Y_2 + \frac{1}{2}Z_3 v^2, \\ m_A^2 &= m_{H^\pm}^2 + \frac{1}{2}(Z_4 - Z_5)v^2, & m_H^2 &= m_A^2 + Z_5 v^2. \end{aligned}$$

Scalar/vector Couplings of the IDM

$$\mathcal{L}_{VVH} = \left(gm_W W_\mu^+ W^{\mu-} + \frac{g}{2c_W} m_Z Z_\mu Z^\mu \right) h ,$$

$$\begin{aligned} \mathcal{L}_{VVHH} = & \left[\frac{1}{4} g^2 W_\mu^+ W^{\mu-} + \frac{g^2}{8c_W^2} Z_\mu Z^\mu \right] (h^2 + H^2 + A^2) \\ & + \left[\frac{1}{2} g^2 W_\mu^+ W^{\mu-} + e^2 A_\mu A^\mu + \frac{g^2}{c_W^2} \left(\frac{1}{2} - s_W^2 \right)^2 Z_\mu Z^\mu + \frac{2ge}{c_W} \left(\frac{1}{2} - s_W^2 \right) A_\mu Z^\mu \right] H^+ H^- \\ & + \left\{ \left(\frac{1}{2} eg A^\mu W_\mu^+ - \frac{g^2 s_W^2}{2c_W} Z^\mu W_\mu^+ \right) H^- (H + iA) + \text{h.c.} \right\} , \\ \mathcal{L}_{VHH} = & \frac{g}{2c_W} Z^\mu A^\mu \overleftrightarrow{\partial}_\mu H - \frac{1}{2} g \left[i W_\mu^+ H^- \overleftrightarrow{\partial}^\mu (H + iA) + \text{h.c.} \right] \\ & + \left[ie A^\mu + \frac{ig}{c_W} \left(\frac{1}{2} - s_W^2 \right) Z^\mu \right] H^+ \overleftrightarrow{\partial}_\mu H^- , \end{aligned}$$

where $s_W \equiv \sin \theta_W$, $c_W \equiv \cos \theta_W$.

The cubic and quartic Higgs self-interactions are governed by

$$\begin{aligned} \mathcal{L}_{3h} = & -\frac{1}{2} v [Z_1 h^3 + (Z_3 + Z_4) h (H^2 + A^2) + Z_5 h (H^2 - A^2)] - v Z_3 h H^+ H^- . \\ \mathcal{L}_{4h} = & -\frac{1}{8} [Z_1 h^4 + Z_2 (H^2 + A^2)^2 + 2(Z_3 + Z_4) h^2 (H^2 + A^2) + 2Z_5 h^2 (H^2 - A^2)] \\ & -\frac{1}{2} H^+ H^- [Z_2 (H^2 + A^2 + H^+ H^-) + Z_3 h^2] . \end{aligned}$$

A natural mass degeneracy of the IDM

$m_H = m_A$, due to $Z_5 = 0$.

This mass degeneracy is due to an exact continuous U(1) symmetry, $H_1 \rightarrow H_1$ and $H_2 \rightarrow e^{i\theta} H_2$, which is preserved by the vacuum. One can now define eigenstates of U(1) charge (not to be confused with electric charge),

$$\phi^\pm = \frac{1}{\sqrt{2}} [H \pm iA] .$$

The relevant interaction terms of ϕ^\pm are

$$\begin{aligned} \mathcal{L}_{\text{int}} = & \left[\frac{1}{2} g^2 W_\mu^+ W^{\mu-} + \frac{g^2}{4c_W^2} Z_\mu Z^\mu \right] \phi^+ \phi^- + \frac{ig}{2c_W} Z^\mu \phi^- \overleftrightarrow{\partial}_\mu \phi^+ - \frac{g}{\sqrt{2}} \left[i W_\mu^+ H^- \overleftrightarrow{\partial}^\mu \phi^+ + \text{h.c.} \right] \\ & + \frac{eg}{\sqrt{2}} \left(A^\mu W_\mu^+ H^- \phi^+ + A^\mu W_\mu^- H^+ \phi^- \right) - \frac{g^2 s_W^2}{\sqrt{2} c_W} \left(Z^\mu W_\mu^+ H^- \phi^+ + Z^\mu W_\mu^- H^+ \phi^- \right) \\ & - v(Z_3 + Z_4) h \phi^+ \phi^- - \frac{1}{2} [Z_2 (\phi^+ \phi^-)^2 + (Z_3 + Z_4) h^2 \phi^+ \phi^-] - Z_2 H^+ H^- \phi^+ \phi^- . \end{aligned}$$

Although ϕ^\pm are mass degenerate states, they can be physically distinguished.

For example, Drell-Yan production via a virtual s -channel W^+ exchange can produce H^+ in association with ϕ^- , whereas virtual s -channel W^- exchange can produce H^- in association with ϕ^+ . Thus, the sign of the charged Higgs boson reveals the U(1)-charge of the produced neutral scalar. The origin of this correlation lies in the fact that, by construction, H^+ and ϕ^+ both reside in H_2 , whereas H^- and ϕ^- both reside in H_2^\dagger .

Mass degeneracies in the most general 2HDM

It is also possible to construct examples of accidental mass degeneracies in the most general 2HDM. However, the *only* natural neutral scalar mass degeneracy in the 2HDM is precisely the case of the IDM with $Z_5 = 0$.

New features of mass degenerate scalars in the 3HDM

In the 3HDM, one can now consider mass-degenerate charged Higgs pairs, as well as mass-degenerate neutral scalars. I will focus on two special 3HDMs where mass degeneracies occur.

The replicated IDM (RIDM)

We begin with a replicated IDM, in which the inert doublets are mass-degenerate. Consider the following 3HDM scalar potential in the Higgs basis,

$$\begin{aligned}\mathcal{V}_{\text{RIDM}} = & Y_1 H_1^\dagger H_1 + Y_2 \left(H_2^\dagger H_2 + H_3^\dagger H_3 \right) + \frac{1}{2} Z_1 (H_1^\dagger H_1)^2 + \frac{1}{2} Z_2 (H_2^\dagger H_2 + H_3^\dagger H_3)^2 \\ & + Z_3 (H_1^\dagger H_1) \left(H_2^\dagger H_2 + H_3^\dagger H_3 \right) + Z_4 \left[(H_1^\dagger H_2)(H_2^\dagger H_1) + (H_1^\dagger H_3)(H_3^\dagger H_1) \right] \\ & + \frac{1}{2} Z_5 \left\{ (H_1^\dagger H_2)^2 + (H_2^\dagger H_1)^2 + (H_1^\dagger H_3)^2 + (H_3^\dagger H_1)^2 \right\} .\end{aligned}$$

Without loss of generality, we have chosen Z_5 real, so that $\mathcal{V}_{\text{RIDM}}$ is CP-conserving. There is a continuous symmetry that is responsible for the mass-degeneracy of the inert Higgs doublets H_2 and H_3 .

Consider the $U(2)$ family symmetry, where the neutral complex field H_1^0 is a singlet and the neutral complex fields H_2^0 and H_3^0 transform as,

$$\begin{pmatrix} H_2^0 \\ H_3^0 \end{pmatrix} \longrightarrow U \begin{pmatrix} H_2^0 \\ H_3^0 \end{pmatrix}, \quad \text{with } U \in U(2).$$

If $Z_5 = 0$, then $\mathcal{V}_{\text{RIDM}}$ depends only on the combination of neutral fields, $H_2^{0\dagger}H_2^0 + H_3^{0\dagger}H_3^0$, and hence is invariant under $U(2)$.

If $Z_5 \neq 0$, then $\mathcal{V}_{\text{RIDM}}$ also depends on the combination of neutral fields, $(H_2^0)^2 + (H_2^{0\dagger})^2 + (H_3^0)^2 + (H_3^{0\dagger})^2$. Hence, $\mathcal{V}_{\text{RIDM}}$ is invariant under an $O(2)$ subgroup of the $U(2)$ transformations (corresponding to real unitary matrices).

The latter guarantees that real and imaginary parts of H_2^0 and H_3^0 are separately mass degenerate. In the case of $Z_5 = 0$ (and the full $U(2)$ family symmetry), one has in addition a mass-degeneracy between the real and imaginary parts of each inert neutral scalar.

There is another continuous symmetry at play here, which takes the form of a generalized CP transformation (GCP),

$$\begin{pmatrix} H_2^0 \\ H_3^0 \end{pmatrix} \longrightarrow U \begin{pmatrix} H_2^{0\dagger} \\ H_3^{0\dagger} \end{pmatrix}, \quad \text{with } U \in \text{U}(2)_{\text{GCP}}.$$

Again, if $Z_5 = 0$, then $\mathcal{V}_{\text{RIDM}}$ is invariant under the $\text{U}(2)_{\text{GCP}}$. If $Z_5 \neq 0$, then $\mathcal{V}_{\text{RIDM}}$ is invariant under an $\text{O}(2)_{\text{GCP}}$ subgroup of $\text{U}(2)_{\text{GCP}}$.

The mass degeneracies of the neutral inert scalars can also be seen as a consequence of the above symmetries.

Remark: The mass degeneracies of the inert charged Higgs scalars are also a consequence of either the $\text{U}(2)$ family symmetry or the $\text{U}(2)_{\text{GCP}}$ generalized CP symmetry. (Note that Z_5 does not contribute to the masses of the inert charged Higgs scalars.)

In the replicated IDM, the Higgs basis doublet fields are mass eigenstate fields,

$$H_1 = \begin{pmatrix} G^+ \\ \frac{1}{\sqrt{2}}[v + h_{\text{SM}} + iG^0] \end{pmatrix}, \quad H_2 = \begin{pmatrix} H^+ \\ \frac{1}{\sqrt{2}}[H + iA] \end{pmatrix}, \quad H_3 = \begin{pmatrix} h^+ \\ \frac{1}{\sqrt{2}}[h + ia] \end{pmatrix},$$

with a minor change of notation from the IDM. The corresponding masses are,

$$m_{H^\pm}^2 = m_{h^\pm}^2 = Y_2 + \frac{1}{2}Z_3 v^2, \quad m_H^2 = m_h^2 = Y_2 + \frac{1}{2}(Z_3 + Z_4 + Z_5)v^2, \\ m_A^2 = m_a^2 = Y_2 + \frac{1}{2}(Z_3 + Z_4 - Z_5)v^2.$$

The corresponding couplings simply replicate the IDM couplings. For example,

$$\mathcal{L}_{VVH} = \left(gm_W W_\mu^+ W^{\mu-} + \frac{g}{2c_W} m_Z Z_\mu Z^\mu \right) h_{\text{SM}}, \\ \mathcal{L}_{VHH} = \frac{g}{2c_W} Z^\mu (A \overleftrightarrow{\partial}_\mu H + a \overleftrightarrow{\partial}_\mu h) - \frac{1}{2}g \left[iW_\mu^+ H^- \overleftrightarrow{\partial}^\mu (H + iA) + iW_\mu^+ h^- \overleftrightarrow{\partial}^\mu (h + ia) + \text{h.c.} \right] \\ + \left[ieA^\mu + \frac{ig}{c_W} \left(\frac{1}{2} - s_W^2 \right) Z^\mu \right] (H^+ \overleftrightarrow{\partial}_\mu H^- + h^+ \overleftrightarrow{\partial}_\mu h^-), \\ \mathcal{L}_{3h} = -\frac{1}{2}v [Z_1 h_{\text{SM}}^3 + (Z_3 + Z_4)h_{\text{SM}}(H^2 + A^2 + h^2 + a^2) + Z_5 h_{\text{SM}}(H^2 - A^2 + h^2 - a^2)] \\ - vZ_3 h_{\text{SM}}(H^+ H^- + h^+ h^-).$$

It is convenient to introduce,

$$P \equiv \frac{H + ih}{\sqrt{2}}, \quad P^\dagger \equiv \frac{H - ih}{\sqrt{2}}, \quad Q \equiv \frac{A - ia}{\sqrt{2}}, \quad Q^\dagger \equiv \frac{A + ia}{\sqrt{2}},$$

where the extra minus sign is for later use. Then, we can rewrite the RIDM couplings in terms of the complex fields P , Q (and their adjoints). For example,

$$\begin{aligned} \mathcal{L}_{VHH} = & \frac{g}{2c_W} Z^\mu (Q \overleftrightarrow{\partial}_\mu P + Q^\dagger \overleftrightarrow{\partial}_\mu P^\dagger) - \frac{g}{2\sqrt{2}} \left[(iW_\mu^+ H^- - W_\mu^- h^+) \overleftrightarrow{\partial}^\mu (P + iQ) \right. \\ & \left. - (iW_\mu^- H^+ - W_\mu^+ h^-) \overleftrightarrow{\partial}^\mu (P - iQ) + \text{h.c.} \right] \\ & + \left[ieA^\mu + \frac{ig}{c_W} \left(\frac{1}{2} - s_W^2 \right) Z^\mu \right] (H^+ \overleftrightarrow{\partial}_\mu H^- + h^+ \overleftrightarrow{\partial}_\mu h^-), \\ \mathcal{L}_{3h} = & -v \left[\frac{1}{2} Z_1 h_{\text{SM}}^3 + (Z_3 + Z_4) h_{\text{SM}} (|P|^2 + |Q|^2) + Z_5 h_{\text{SM}} (|P|^2 - |Q|^2) \right] - v Z_3 h_{\text{SM}} (H^+ H^- + h^+ h^-). \end{aligned}$$

In the RIDM, there is no experimental measurement that can physically distinguish the degenerate scalars, (H^\pm, h^\pm) , (H, h) and (A, a) . However, the multiplicity factor will appear after summing over final mass-degenerate states, e.g., $Z \rightarrow HA$, ha (or equivalently, $Z \rightarrow PQ$, $P^\dagger Q^\dagger$), doubles the rate into a pair of neutral scalars.

The Ivanov-Silva Model

Ivanov and Silva (IS) introduced a particular 3HDM model with some curious properties.* In the Higgs basis of the 3HDM, we are free to make an arbitrary $U(2)$ rotation to define the Higgs basis fields, H_2 and H_3 . We have made use of this freedom to make a minor alteration of the IS scalar potential,

$$\begin{aligned}\mathcal{V}_{\text{IS}} = \mathcal{V}_{\text{RIDM}} &+ Z'_3(H_2^\dagger H_2)(H_3^\dagger H_3) + Z'_4(H_2^\dagger H_3)(H_3^\dagger H_2) \\ &+ [Z_8(H_2^\dagger H_3)^2 + Z_9(H_2^\dagger H_3)(H_2^\dagger H_2 - H_3^\dagger H_3) + \text{h.c.}] ,\end{aligned}$$

where $\mathcal{V}_{\text{RIDM}}$ is the replicated IDM scalar potential, and Z_8 and Z_9 are potentially complex.

The IS model still yields mass-degenerate inert doublets, since none of the extra terms involve the Higgs basis field H_1 . Hence, these terms do not contribute to the tree-level scalar squared-mass matrices.

*I.P. Ivanov and J.P. Silva, Phys. Rev. D **93**, 095014 (2016) [arXiv:1512.09276],

Symmetries governing the mass degeneracies of the IS model

Note that after the extra terms above are included, there is no remaining unbroken continuous subgroup of the $U(2)$ family symmetry or the $U(2)_{\text{GCP}}$ generalized CP symmetry (either of which was responsible for the two mass-degenerate inert doublets of the RIDM).

Case 1: Z_8 and Z_9 are real.

\mathcal{V}_{IS} is invariant under a discrete \mathbb{Z}_4 subgroup of the $U(2)$ family symmetry group. The elements of this subgroup are,

$$\mathbb{Z}_4 = \{I, -I, Z, -Z\}, \quad \text{where } Z \equiv \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

where the 2×2 matrices above act on the Higgs basis fields H_2 and H_3 . Note that $Z^2 = -I$, where I is the 2×2 identity matrix.

The fields H_2 and H_3 are odd under $-I$, which simply identifies the two inert doublets. The elements Z (and $-Z$) act non-trivially on the inert doublets.

As before, we are free to combine mass-degenerate neutral fields and define,

$$P \equiv (H + ih)/\sqrt{2} \quad \text{and} \quad Q \equiv (A - ia)/\sqrt{2},$$

which are eigenstates of Z (and $-Z$). Indeed, P and Q^\dagger have eigenvalue i under Z , and P^\dagger and Q have eigenvalue $-i$ under Z . For example, this is consistent with the couplings of neutral scalars to the Z , namely

$$\mathcal{L}_{ZHH} = \frac{g}{2c_W} Z^\mu (P \overleftrightarrow{\partial}_\mu Q + P^\dagger \overleftrightarrow{\partial}_\mu Q^\dagger).$$

Likewise, \mathcal{V}_{IS} is invariant under a discrete \mathbb{Z}_4 subgroup of the $\text{U}(2)_{\text{GCP}}$ generalized CP symmetry. The element Z involved in the transformation,

$$\begin{pmatrix} H_2 \\ H_3 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} H_2^\dagger \\ H_3^\dagger \end{pmatrix},$$

is called a CP4 transformation by Ivanov and Silva.[†] Due to the extra dagger, P and Q have eigenvalue i and P^\dagger and Q^\dagger have eigenvalue $-i$ under Z . This is again consistent with the form of \mathcal{L}_{ZHH} above since the Z is CP-even and parity introduces an extra minus sign in the space derivative.

[†]Note that $(\text{CP}4)^2 = -I$ and $(\text{CP}4)^4 = I$. Hence the nomenclature.

Either discrete symmetries (family or GCP) can be invoked to explain the observed mass degeneracies of the IS model with real Z_8 and Z_9 . Moreover, the ordinary CP symmetry, called CP2 by Ivanov and Silva, corresponding to $H_i \rightarrow H_i^\dagger$, is respected since all scalar potential parameters are real.

Case 2: Z_8 and/or Z_9 are complex.

In this case, the symmetry,

$$\begin{pmatrix} H_2 \\ H_3 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} H_2 \\ H_3 \end{pmatrix},$$

is no longer respected by \mathcal{V}_{IS} . The remaining unbroken family symmetry is $\mathbb{Z}_2 = \{I, -I\}$, which protects the inertness of H_2 and H_3 but cannot enforce the mass degeneracies of the IS model.

Nevertheless, the CP4 symmetry remains intact and is ultimately responsible for the IS model mass degeneracies. Note that there is no CP2 symmetry in this case, since there is no possible change of basis in which all scalar potential parameters are real.

A physical distinction between the CP2 and CP4 symmetry


Ivanov and Silva asked: is there an experiment that can determine the order of the CP symmetry of the scalar sector? The answer is affirmative. It relies on the existence of a particular four scalar coupling of the IS model,

$$\delta\mathcal{L}_{4h} \ni \frac{1}{2} \text{Im } Z_8 [(PQ - P^\dagger Q^\dagger)(P^2 - Q^2 - P^{\dagger 2} + Q^{\dagger 2})] \\ + \frac{1}{2} i \text{Im } Z_9 [(PQ - P^\dagger Q^\dagger)(P^2 + Q^2 + P^{\dagger 2} + Q^{\dagger 2})] .$$

Self-interaction terms of this type are absent if Z_8 and Z_9 are both real. As an example, consider the case where $M_Q \ll m_Z$ and $M_P \gg m_Z$. In this case, the four-scalar interactions above mediate the four body Z decay,

$$Z \rightarrow QQQQ^\dagger, \quad QQ^\dagger Q^\dagger Q^\dagger .$$

These two final states are experimentally indistinguishable, so we must sum incoherently the squared amplitudes of both channels. Observation of such decays would be consistent with the presence of a CP4 symmetry and would force us to conclude that it is impossible to define CP as a CP2 symmetry.



We have obtained (for $M_P \gg m_Z$ and $M_Q = 0$),

$$\frac{\Gamma(Z \rightarrow QQQQ^\dagger, QQ^\dagger Q^\dagger Q^\dagger)}{\Gamma(Z \rightarrow \nu\bar{\nu})} = \frac{2[(\text{Im } Z_8)^2 + (\text{Im } Z_9)^2]}{3 \cdot 5 \cdot 2^9 \pi^4} \left(\frac{m_Z}{M_P} \right)^4,$$

where the factor of **2** accounts for the multiplicity of mass-degenerate states.

Final comments

1. It is straightforward to show that for an N -Higgs doublet model, a CP4-symmetric scalar potential *and* CP4-invariant vacuum implies the existence of mass degenerate scalar states (similar to that of the IS model).
2. There is an observable distinction between CP4-invariant models and conventionally CP-invariant (CP2) models.
3. Do CP4 invariant scalar sectors that violate CP2 yield any observable T-violating phenomena? (Ivanov says no!)

We had some hope that one could find evidence for T-violating form factors arising in the ZZZ and ZW^+W^- vertex, that would be generated at two loops due to the CP2-violating, CP4-conserving PQ^3 and P^3Q interactions. However, it seems that such contributions vanish exactly (due to the absence of diagrams or diagrams canceling in pairs).

Backup slides

Flashback to 2012: Can Mass-degenerate scalars explain the $h \rightarrow \gamma\gamma$ anomaly?



After the initial discovery of the Higgs boson in 2012, it appeared that the signal strength for $h \rightarrow \gamma\gamma$ was significantly enhanced above Standard Model (SM) expectations.

My collaborators and I proposed to explain this anomaly under the assumption that the the observed Higgs state at 125 GeV was in fact a pair of mass degenerate scalars.[‡]

We considered the Type-I and Type-II two-Higgs doublet model (2HDM), and explored various possibilities for mass degeneracy and their phenomenological consequences.

[‡]P.M. Ferreira, R. Santos, H.E. Haber and J.P. Silva, Phys. Rev. D **87**, 055009 (2013) [arXiv:1211.3131].

An enhanced $\gamma\gamma$ signal due to mass-degenerate h^0 and A^0 :



Left panel: $R_{\gamma\gamma}$ as a function of $\tan\beta$ for h (blue), A (green), and the total observable rate (cyan), obtained by summing the rates with intermediate h and A , for the unconstrained scenario (i.e., the effects of virtual charged Higgs exchange in B physics is neglected).

Right panel: Total rate for $R_{\gamma\gamma}$ as a function of $\tan\beta$ for the constrained (red) and unconstrained (green) scenarios.

Above, $R_f^H = \frac{\sigma(pp \rightarrow H)_{2\text{HDM}} \times \text{BR}(H \rightarrow f)_{2\text{HDM}}}{\sigma(pp \rightarrow h_{\text{SM}}) \times \text{BR}(h_{\text{SM}} \rightarrow f)}$, where f is the final state of interest, and H is one of the two 125 GeV mass-degenerate scalars. The observed ratio of f production relative to the SM expectation is $R_f \equiv \sum_H R_f^H$. In our analysis, we assumed that $R_{WW} \simeq R_{ZZ} \simeq 1 \pm 0.2$.

The corresponding results in the Type-II 2HDM were similar. Other degenerate-mass scalar pairs were also considered. By the end of Run I of the LHC, the $\gamma\gamma$ excess was gone, and the Higgs data appears to be consistent with SM expectations.

A simple model of scalar mass degeneracy: H^\pm

Any doublet extended Higgs model has a mass degenerate state—the charged Higgs boson, H^\pm . Indeed, H^+ and H^- are degenerate due to the $U(1)_{\text{EM}}$ gauge symmetry. Moreover, the H^+ and H^- are distinguishable by their electric charge, which we can probe using photons.

Suppose that this probe was unavailable (or equivalently, suppose one could turn off electromagnetism). Can experiment reveal the existence of a mass-degenerate scalar?

- Given a charged Higgs state, one could not physically distinguish between the two degenerate states.
- However, there would in principle be observables that are sensitive to the number of degenerate states present. Examples: $H \rightarrow H^+H^-$ (but not $Z \rightarrow H^+H^-$ due to the off-diagonal nature of this coupling).

Mass degeneracies in the most general 2HDM

To analyze the most general 2HDM, we note a remarkable tree-level relation

$$\text{Im}(Z_5^* Z_6^2) = \frac{2s_{13}c_{13}^2 s_{12}c_{12}}{v^6} (m_2^2 - m_1^2)(m_3^2 - m_1^2)(m_3^2 - m_2^2),$$

where the m_i ($i = 1, 2, 3$) are the masses of the three neutral Higgs bosons of the 2HDM, $s_{12} \equiv \sin \theta_{12}$, $c_{12} \equiv \cos \theta_{12}$, etc., and θ_{12} and θ_{13} are invariant mixing angles that are associated with the diagonalization of the neutral Higgs squared-mass matrix in the Higgs basis.

Thus, if any mass degeneracy is present, then one can find a Higgs basis in which Y_3 , Z_5 and Z_6 are simultaneously real. Any CP-violating effects arise due to a potentially complex Z_7 , which enters in the Higgs self-couplings but *not* the diagonalization of the tree-level neutral scalar squared-mass matrix.

Hence, without loss of generality, we can simply take Z_5 and Z_6 real and identify the neutral Higgs scalars as h , H and A . These are states of definite CP in their interactions with gauge bosons (and fermions).

The resulting Higgs mass relations are then,

$$m_{H^\pm}^2 = Y_2 + \frac{1}{2}Z_3v^2, \quad m_A^2 = m_{H^\pm}^2 + \frac{1}{2}(Z_4 - Z_5)v^2, \\ m_{H,h}^2 = \frac{1}{2} \left\{ m_A^2 + (Z_1 + Z_5)v^2 \pm \sqrt{[m_A^2 - (Z_1 - Z_5)v^2]^2 + 4Z_6^2v^4} \right\}.$$

Mass degenerate states arise if one of the following two quantities is zero,

$$Z_5(m_A^2 - Z_1v^2) + Z_6^2v^2 = 0 \quad \text{or} \quad [m_A^2 - (Z_1 - Z_5)v^2]^2 + 4Z_6^2v^4 = 0.$$

Case 1: $m_h = m_H$

It follows that $m_A^2 = (Z_1 - Z_5)v^2$ and $Z_6 = 0$. Thus, we recover the IDM mass spectrum for this degenerate case, although Z_7 can be nonzero. Thus, the IDM scalar self couplings are modified by the addition of the following terms,

$$\delta\mathcal{L}_{3h} = -\frac{1}{4}v[Z_7(H + iA) + Z_7^*(H - iA)](HH + AA + 2H^+H^-), \\ \delta\mathcal{L}_{4h} = -\frac{1}{4}[Z_7(H + iA) + Z_7^*(H - iA)](HH + AA + 2H^+H^-)h,$$

which provide new sources of CP violation if $\text{Im}Z_7 \neq 0$. The mass degeneracy is unnatural (moreover, $Z_6 = 0$ is also unnatural when $Z_7 \neq 0$). Nevertheless, the mass-degenerate Higgs bosons are distinguishable as in the IDM.

Cases 2 and 3: $m_h = m_A$ or $m_H = m_A$

Both these possibilities arise when $Z_5(m_A^2 - Z_1 v^2) + Z_6^2 v^2 = 0$, which is an unnatural condition (unless $Z_5 = Z_6 = Z_7 = 0$). The physical distinction of the mass degenerate states is due to the CP quantum numbers of the neutral scalar states (which are preserved by the Higgs interactions with gauge bosons and fermions). One can therefore distinguish between the corresponding production mechanisms of the degenerate scalars that are mediated by gauge boson fusion or Drell-Yan production via s -channel gauge boson exchange.

Case 4: $m_h = m_H = m_A$

This requires $Z_5 = Z_6 = 0$ and $m_A^2 = Z_1 v^2$. This leaves Z_7 as the only potentially complex parameter of the scalar potential in the Higgs basis, which can be chosen real by rephasing the Higgs basis field H_2 . Hence, the Higgs scalar potential and vacuum must be CP-conserving. However, as long as $Z_7 \neq 0$, the triply mass-degenerate case is unnatural, since the \mathbb{Z}_2 symmetry of the IDM is not present.