

Germanium-detector based study of the $2\text{H}(p,\gamma)3\text{He}$ cross section at LUNA

Monday 11 June 2018 17:20 (10 minutes)

Recent, precise measurements of the primordial 2H abundance [1] have opened the possibility to precisely determine of the primordial baryon-to-photon ratio, independent from the cosmic microwave background. For their interpretation, the 2H abundance data require equally precise nuclear data, in particular on the $2\text{H}(p,\gamma)3\text{He}$ reaction.

Deep underground in the Gran Sasso laboratory, Italy, the LUNA collaboration is undertaking a dedicated effort to measure the $2\text{H}(p,\gamma)3\text{He}$ cross section directly in the Big

Bang energy window of interest. The campaign is divided in two phases based on a BGO and a high-purity germanium (HPGe) detector, respectively.

The present poster will report on the second, HPGe-based phase of the experiment.

Due to the Doppler shift of the emitted γ -rays, in addition to the absolute yield also information on the γ -ray angular distribution, thus reducing the systematic uncertainty.

The characterization and calibration of the setup and detectors, background conditions, and potential sources of uncertainty will be discussed.

References

[1] R. J. Cooke, M. Pettini, R. A. Jorgenson, M. T. Murphy, and C. C. Steidel, *Astrophys. J.* 781, 31 (2014).

Primary author: Mr STÖCKEL, Klaus (HZDR)

Presenter: Mr STÖCKEL, Klaus (HZDR)

Session Classification: Poster Session