


# Drell-Yan production of heavy vectors in Higgs-less models

Jernej F. Kamenik



Institut "Jožef Stefan"





# Introduction

- Experiments provide unambiguous indications that the SM gauge group is spontaneously broken [ $SU(2)_L \times U(1)_Y \rightarrow U(1)_Q$ ]
- One elementary  $SU(2)_L$  scalar doublet with  $\varphi^4$  potential is the **most economical & simple choice**
- $L_{\text{Higgs}}(\varphi, A^i, \psi^i) = D_\mu \varphi^\dagger D_\mu \varphi + \mu^2 \varphi^\dagger \varphi - \lambda (\varphi^\dagger \varphi)^2 + Y^{ij} \psi_L^i \psi_R^j$ 
  - **not the only allowed possibility**
- So far only the ground state of this Lagrangian has been tested with good accuracy
  - $\langle \varphi \rangle = 246 \text{ GeV} \leftrightarrow m_W, m_Z$
- Some dynamical sensitivity to the Higgs mechanism is obtained from EWPO
  - **Indirect indication of a light  $m_H$  (in the SM)**

c.f. talk by Pomarol

Peskin & Takeuchi  
[PRL65:964,1990]

Altarelli & Barbieri  
[PLB253:161,1991]

...

Barbieri et al.  
[hep-ph/0405040]



# Do we need a fundamental Higgs field?

- EWPO indicate:

- a spontaneous breaking of  $SU(2)_L \times U(1)_Y$
- the breaking mechanism must respect, to a good accuracy, the **custodial symmetry** [  $m_Z^2/m_W^2 \approx 1 + (g'/g)^2$  ]

- General formulation of the symmetry breaking mechanism in absence of a fundamental Higgs (or for large Higgs masses) in terms of a Chiral Lagrangian:

$$\mathcal{L}_X^{(2)} = \frac{v^2}{4} \text{Tr}(D_\mu U^\dagger D^\mu U)$$

$$U \rightarrow g_R U g_L^\dagger = e^{i\pi/v}$$

3 Goldstones of the SM

- Global:  $SU(2)_L \times SU(2)_R \times U(1)_{B-L} \rightarrow SU(2)_{L+R} \times U(1)_{B-L}$

- Local:  $SU(2)_L \times U(1)_Y \rightarrow U(1)_Q$



$$D_\mu U = -ig' B_\mu U + ig U W_\mu$$



# EW Chiral Lagrangian

$$\mathcal{L}_{eff} = \mathcal{L}_{gauge}(A^i, \psi^i) + \mathcal{L}_{Yukawa}(U, \psi^i) + \frac{v^2}{4} \text{Tr}(D_\mu U^\dagger D^\mu U)$$

- contains all the degrees of freedom we have directly probed in experiments
- naive cut-off dictated by the convergence of EW loops:  $\Lambda_{NDA} = 4\pi v \approx 3 \text{ TeV}$
- perfectly describes particle physics up to 3 TeV, beyond the tree level, with only two drawbacks
  - (point toward the existence of new degrees of freedom below the naive cut-off):
  - Violation of unitarity in  $W_L W_L \rightarrow W_L W_L$  scattering (tree-level amplitude violates unitarity for  $s \approx 1 \text{ TeV}$ )
  - Bad fit to S and T





# Introducing heavy vectors

- A natural alternative to Higgs-type mechanisms in curing the problem of unitarity in  $WW \rightarrow WW$  scattering is represented by heavy vector fields
- Expected in many non-SUSY scenarios:
  - techni-rho in technicolor,
  - massive gauge bosons in 5-dimensional theories, hidden gauge-models
- Difficult task is to cure at the same time unitarity and EWPO
  - can be analysed in general terms constructing an appropriate effective chiral Lagrangian with the heavy vectors as new explicit d.o.f.

$$\mathcal{L}_\chi = \frac{v^2}{4} \text{Tr}(D_\mu U^\dagger D^\mu U) + \mathcal{L}_{kin}(R, U, A_i; m_R) + \mathcal{L}_{int}(R, U, A_i; G_R)$$



# Heavy vectors in the EW Chiral Lagrangian

Barbieri et al.  
[0806.1624]

- Consider an effective theory based on the following two main assumptions:
  - The (new) dynamics that breaks the SM EW symmetry is invariant under the global symmetry  $SU(2)_L \times SU(2)_R$  and under the discrete parity  $P$ :  $SU(2)_L \leftrightarrow SU(2)_R$
  - One vector ( $V$ ), or one vector + one axial-vector ( $V+A$ ), both belonging to the adjoint representation of  $SU(2)_{L+R}$  (triplets), are the only light fields below a cut-off  $\Lambda = 2-3$  TeV
- Effective Lagrangian analysis based on ordering of operators according to the standard derivative (momentum) expansion



# Unitarizing $W_L W_L$ scattering

Barbieri et al.  
[0806.1624]


- No tree-level violation of unitarity for

- $G_V^2 = v^2/3$

- The unitarity constraint is almost insensitive to the value  $m_V$



$$\mathcal{M} = \frac{s}{v^2} - \frac{G_V^2}{v^4} \left[ 3s + m_V^2 \left( \frac{s-u}{t-m_V^2} + \frac{s-t}{u-m_V^2} \right) \right]$$





# EWPO

Barbieri et al.  
[0806.1624]


Cata & J.F.K  
[1006.xxxx]

- The leading contributions to S & T generated by the exchange of single heavy fields

$$\Delta\hat{S} = g^2 \left( \frac{F_V^2}{4m_V^2} - \frac{F_A^2}{4m_A^2} \right) + \frac{g^2}{6} \frac{\Lambda^2}{(4\pi v)^2} \left[ - \left( \frac{F_V - 2G_V}{2m_V} \right)^2 + \left( \frac{F_A}{2m_A} \right)^2 - \frac{v^2}{m_V^2} - \frac{v^2}{m_A^2} \right] + \dots$$


$$\Delta\hat{T} = \frac{3\pi\alpha}{c_W^2} \left[ \frac{F_A^2}{4m_A^2} + \left( \frac{F_V - 2G_V}{2m_V} \right)^2 \right] \frac{\Lambda^2}{16\pi^2 v^2} + \dots$$

O(1) factors [ $\Lambda$  replaced by some heavy mass]



- Two natural ways to accommodate the bounds:
  - Both V and A light, almost degenerate
  - Only V light, with small  $F_V$
- EWPO & unitarity can be accommodated for specific choices of the free parameters

**Main conclusion:**  
We need at least one relatively light vector field





# Producing the heavy vectors at the LHC

- Main properties of vector fields

- Leading decay mode: 2 longitudinal SM gauge bosons

- $\Gamma_{V^+} \approx \Gamma_{WZ}^V = \frac{G_V^2 m_V^3}{48\pi v^4} [1 + \mathcal{O}(g^2 \epsilon^2)]$  ,     5 GeV [  $m_V = 0.5$  TeV ]  
40 GeV [  $m_V = 1.0$  TeV ]

- $\Gamma_{V^0} \approx \Gamma_{WW}^V = \Gamma_{WZ}^V [1 + \mathcal{O}(g^2 \epsilon^2)]$

- Narrow widths!

- ZZ channel forbidden

- Coupling to SM fermions highly suppressed

- $Br(V^0 \rightarrow q\bar{q}) \approx 3Br(V^0 \rightarrow \ell^+\ell^-) \approx \frac{6F_V^2 m_W^4}{G_V^2 m_V^4}$      1.6% [  $m_V = 0.5$  TeV,  $F_V = 2G_V$  ]  
0.1% [  $m_V = 1.0$  TeV,  $F_V = 2G_V$  ]


- Leading decay modes of axial fields can be to a vector and SM g. b.




# Producing the heavy vectors at the LHC

- The most general signature of Higgsless models is the appearance of the vector state in WW scattering [  $pp \rightarrow \nu + jj$  (WW fusion)  $\rightarrow WW(WZ) + jj$  ]

- Model-independent link with the unitarity problem



Belyaev  
[0711.1919]






# Producing the heavy vectors at the LHC

- The most general signature of Higgsless models is the appearance of the vector state in WW scattering [  $pp \rightarrow V + jj$  (WW fusion)  $\rightarrow WW(WZ) + jj$  ]

- A difficult analysis, which requires high statistics.



Belyaev  
[0711.1919]


- Resonant cross section including
- leptonic BR's ( $l=e,\mu$ ) [  $\epsilon_{\text{lept}} = 21\% \times 6.7\% = 1.5\%$  ]
  - $p_T(\text{jets}) > 30$  GeV
  - standard VBF jet cuts [  $\Delta\eta > 4, M_{jj} > 1\text{TeV}$   $\epsilon_{\text{VBF}} < 30\%$  ]



# Producing the heavy vectors at the LHC

- A potentially cleaner signal (if the resonances are not too heavy) is the Drell-Yan production of the resonances and subsequent decay into  $l^+l^-$ , 2 and 3 SM heavy gauge bosons

- [Link to the contribution of the heavy vectors to EPWO](#)





# Producing the heavy vectors at the LHC

- Given the narrow widths, for low masses the signals are quite large

	$M = 500 \text{ GeV}$	$M = 750 \text{ GeV}$	$M = 1000 \text{ GeV}$
$\sigma(pp \rightarrow V^+ \rightarrow X)_{\sqrt{s}=14 \text{ TeV}}$	11 pb	1.2 pb	0.23 pb
$\sigma(pp \rightarrow V^+ \rightarrow X)_{\sqrt{s}=10 \text{ TeV}}$	6.7 pb	0.71 pb	0.13 pb
$\sigma(pp \rightarrow V^+ \rightarrow X)_{\sqrt{s}=7 \text{ TeV}}$	4.2 pb	0.32 pb	0.06 pb


- However...
  - The leading decay modes (2W, 3W) have low efficiencies
  - The  $l^+l^-$  case is suppressed by the small  $\text{Br}(R \rightarrow l^+l^-)$

update of  
Cata, Isidori & J.F.K  
[0905.0490]  
using  
 $F_V^2 = 4G_V^2 = 4v^2/3$



# Signal of heavy vectors at the Tevatron?


- The  $l^+l^-$  state of the art are the analyses of the  $e^+e^-$  and  $\mu^+\mu^-$  final states in  $p\text{-}\bar{p}$  collisions published by CDF & D0
- Using data as normalization for the SM events (takes into account all the relevant exp. efficiencies!), one can produce an exclusion plot in the  $F_V\text{-}m_V$  plane - complementary to EWPT bounds
- Two main assumptions:
  - $G_V$  fixed by unitarity
  - $m_A \gg m_V$



CDF [0810.2059]  
[0811.0053]  
Publ. 10165

D0 Conf. Note 5923

See talk by Wittich



Cata, Isidori & J.F.K  
[0905.0490]




# Signal of heavy vectors at the Tevatron?

- The  $l^+l^-$  state of the art are the analyses of the  $e^+e^-$  and  $\mu^+\mu^-$  final states in  $p\text{-}\bar{p}$  collisions published by CDF & D0

- If, on the other hand, the excess at higher mass will become significant, we can hope to see a clear signal at the LHC (even with  $1\text{-}2 \text{ fb}^{-1}$  @  $14\text{TeV}$ )


- Not huge peaks as with a sequential  $Z'$ , but they should be clearly visible.



CDF [0810.2059]  
[0811.0053]  
Publ. 10165

D0 Conf. Note 5923

See talk by Wittich



Cata, Isidori & J.F.K  
[0905.0490]


Analysis of prospects  
for  $1\text{fb}^{-1}$  @  $7\text{TeV}$   
in progress



# Signals for heavy vectors at the LHC

- Two SM gauge boson final states
- Some illustrative examples
  - $[WZ] \text{ Br}Z_{\text{lept}} \times \text{Br}W_{\text{lept}} = 1.5 \%$
  - $F_V = 2G_V$
  - $F_A = F_V$
  - $G_V$  fixed by unitarity

- [Warning: the configurations of free params. are realistic, but maximize the signal...]



See also  
He et al.  
[0708.2588]



Accomando et al.  
[0807.5051, 0807.2951]

Belyaev et al.  
[0809.0793]



# Signals for heavy vectors at the LHC

- Three SM gauge boson final states
- Some illustrative examples
  - [WWZ]  $\text{BrZ}_{\text{lept}} \times \text{BrW}_{\text{lept}} \times \text{BrW}_{\text{had}} = 0.9\%$
  - $F_V = 2G_V$
  - $F_A = F_V$
  - $G_V$  fixed by unitarity
  - $g_A = 1/2$
- In the WWZ final state it is also worth to look at the WZ invariant-mass distribution





# Conclusions

- Heavy vector fields, which replace the Higgs boson in maintaining perturbative unitarity up to LHC energies, are naturally expected in a wide class of Higgs-less models.
- The most general signature of these models is the appearance of the lightest vector state in  $WW$  scattering (model-independent link with the unitarity problem).
- The Drell-Yan production of the new states is subject to larger uncertainties.
- For light  $m_{V(A)}$  we could expect visible signals (even with low statistics), and the information could help to clarify the role of the heavy vectors in EWPO.
- The results in the  $e^+e^-$  and  $\mu^-\mu^+$  channels from Tevatron are already providing significant information.
- The 2 and 3 SM gauge boson final states seems to be quite promising and a more realistic study is in progress.

See also  
Barbieri et al.  
[0911.1942]



Backup Slides




# EWPO in the SM

Peskin & Takeuchi  
[PRL65:964,1990]  
Altarelli & Barbieri  
[PLB253:161,1991]  
...  
Barbieri et al.  
[hep-ph/0405040]

- Some dynamical sensitivity to the Higgs mechanism is obtained from EWPO

- Indirect indication of a light  $m_H$  under the hypothesis of a heavy cut-off for the SM as effective theory

- ( $\leftrightarrow$  fine tuning in the Higgs mass term)



$$T = \frac{\Pi_{33}(0) - \Pi_{WW}(0)}{m_W^2}$$

$$S = \frac{g}{g'} \left. \frac{d\Pi_{30}(q^2)}{dq^2} \right|_{q^2=0}$$



# Heavy vectors in the EW Chiral Lagrangian

- With heavy spin-1 fields, there is a peculiar problem related to the possible mixing of the heavy states and the Goldstone bosons.
- Describing the heavy states in terms of Lorentz vectors ( $V_\mu$  &  $A_\mu$ ), we have a possible mass-mixing of  $O(p)$  [  $\rightarrow$  tedious redefinition of the fields ]

- $V_\mu \rightarrow V_\mu + \beta [ \pi, \partial_\mu \pi ], \quad A_\mu \rightarrow A_\mu + \alpha \partial_\mu \pi$

- This problem can be avoided describing the heavy spin-1 states by means of **antisymmetric tensors** ( $R_{\mu\nu} = V_{\mu\nu}, A_{\mu\nu}$ ):

$$\mathcal{L}_{kin}(R^{\mu\nu}) = -\frac{1}{2} \text{Tr}(\nabla_\mu R^{\mu\nu} \nabla^\sigma R_{\sigma\nu}) + \frac{1}{4} m_R^2 \text{Tr}(R^{\mu\nu} R_{\mu\nu})$$


$$\langle 0 | R^{\mu\nu} | R(p, \epsilon) \rangle = \frac{i}{m_R} [p_\mu \epsilon_\nu - p_\nu \epsilon_\mu]$$

$$\nabla_\mu R = \partial_\mu R + [\Gamma_\mu, R] \quad \Gamma_\mu = \frac{1}{2} [u^\dagger D_\mu u + u D_\mu u^\dagger], \quad u^2 = U$$



# Heavy vectors in the EW Chiral Lagrangian

- In the antisymmetric formulation the couplings between heavy fields and Goldstone bosons start at  $O(p^2)$   $\Rightarrow$  integrating out the heavy fields we are automatically projected into the basis of the  $O(p^4)$  chiral operators with light fields only.
  - In QCD case this procedure leads to a successful description of all the leading  $O(p^4)$  light-field couplings
- $1 \Leftrightarrow 1$  correspondence between lowest-order vector couplings [  $O(p^2)$  ] and next-to-leading order Goldstone-boson couplings [  $O(p^4)$  ]





# Heavy vectors in the EW Chiral Lagrangian

- The dynamics of the system below the cut-off is described by 3 + 2 parameters:  $(M_V, G_V, F_V) + (M_A, F_A)$ .


- Naive dimensional analysis implies  $F_{V(A)}, G_V = O(v)$

$$[u_\mu = iu^\dagger D_\mu U u^\dagger]$$

$$\mathcal{L}_{int} = \frac{i}{2\sqrt{2}} G_V \text{Tr}(V^{\mu\nu} [u_\mu, u_\nu])$$

$$+ \frac{1}{2\sqrt{2}} F_V \text{Tr}(V^{\mu\nu} (u \hat{W}^{\mu\nu} u^\dagger + u^\dagger \hat{B}^{\mu\nu} u))$$

$$+ \frac{1}{2\sqrt{2}} F_A \text{Tr}(A^{\mu\nu} (u \hat{W}^{\mu\nu} u^\dagger - u^\dagger \hat{B}^{\mu\nu} u))$$



- Specific UV completions of this effective theory correspond to specific choices of the free parameters.



# Producing the heavy vectors at the LHC

- Main properties of axial fields

- $O(m_A^3)$  widths only from  $A \rightarrow VW$

- [mediated by effective ops. with two heavy fields  $A[\partial V, \partial U]$ , not included in  $L_{int}$ ]

- potentially suppressed if  $m_A \approx m_V$

- $\Gamma_{V+W^-}^A = \Gamma_{V^-W^+}^A = \Gamma_{V^0W^+}^A = \Gamma_{V+Z}^A \doteq \Gamma_{VW}^A,$

- $\Gamma_{VW}^A = \frac{m_A^3}{48\pi v^2} (1-r^2)^3 \left[ g_A^2 (1+2r^2) + g_V^2 \left( 1 + \frac{2}{r^2} \right) + 6g_A g_V \right]$

- $O(m_A)$  widths of the type  $A \rightarrow$  longitudinal + transverse SM gauge bosons,

- $\Gamma_{WW}^A = \frac{g^2 F_A^2 m_A}{192\pi v^2}, \quad \Gamma_{WZ}^A = \frac{1}{2} \Gamma_{WW}^A \left[ 1 + \frac{(1-2s_W^2)^2}{c_W^2} \right], \quad \Gamma_{W\gamma}^A = 2s_W^2 \Gamma_{WW}^A$



- leading decay modes if  $m_A \approx m_V$

- Decay widths to SM fermions identical to the vector case, with corresponding BR enhanced by the suppression of the total rate



# Producing the heavy vectors at the LHC

- A potentially clean signal (if the resonances are not too heavy) is the Drell-Yan production of the resonances and subsequent decay into  $l^+l^-$ , 2 and 3 SM heavy gauge bosons
- easy to estimate (and simulate) normalizing the non-standard rate to SM Drell-Yan processes at the partonic level





# Producing the heavy vectors at the LHC

Cata, Isidori & J.F.K  
[0905.0490]

- E.g. for charged final states we define the form factor

- $$F_f^{R^+}(q^2) = \frac{\sigma(u\bar{d} \rightarrow R^+ \rightarrow f)}{\sigma(u\bar{d} \rightarrow \mu^+\nu)_{\text{SM}}}$$

- $$\frac{d}{dq^2}\sigma(pp \rightarrow R^+ \rightarrow f) = F_f^{R^+}(q^2) \frac{d}{dq^2}\sigma(pp \rightarrow \mu^+\nu)_{\text{SM}}$$

- As long as we can neglect interference effects (with SM or among different resonant contributions), the partonic resonant width is simply given by

- $$\sigma(q_i\bar{q}_j \rightarrow R \rightarrow f) = \frac{12\pi\Gamma_R^2 Br_{\text{in}}^R Br_f^R}{(q^2 - m_R^2)^2 + m_R^2\Gamma_R^2} \left[ 1 + \mathcal{O}\left(\frac{q^2 - m_R^2}{m_R^2}\right) \right]$$



# Signal of heavy vectors at the Tevatron?

- The  $l^+l^-$  state of the art is the analysis of the  $e^+e^-$  final state in  $p\bar{p}$  collisions published by CDF
- The “ $2\sigma$  excess” can be fitted nicely by a light vector resonance:

- $m_V \approx 246 \text{ GeV}$


- $F_V \approx 50 \text{ GeV}$

- Predictions derived within the effective theory:


- similar peak also in the  $\mu^+\mu^-$  final state

- axial state with  $m_A \approx 1.3 \text{ TeV}$  to obtain a good EWPO fit

excluded by  
CDF [0811.0053]



CDF [0810.2059]



Cata, Isidori & J.F.K  
[0905.0490]



# Signals for heavy vectors at the LHC

- Two & three SM gauge boson final states
- A detailed estimate of the realistic efficiency for the detection of the heavy vectors in these final states  $[WZ, WW] + [WWW, WWZ, WZZ]$  has not been performed yet. So far we have analysed only the signal against the irreducible SM background = same e.w. final state
- Selecting leptonic decay is a high price to pay (in terms of efficiencies), but it should ensure a good rejection against non-irreducible backgrounds.
  - Some reference theoretical efficiencies:
    - $[WZ] \text{ Br}_{Z_{\text{lept}}} \times \text{Br}_{W_{\text{lept}}} = 1.5 \%$
    - $[WWZ] \text{ Br}_{Z_{\text{lept}}} \times \text{Br}_{W_{\text{lept}}} \times \text{Br}_{W_{\text{had}}} = 0.9 \%$
    - $[WZZ] \text{ Br}_{Z_{\text{lept}}} \times \text{Br}_{W_{\text{lept}}} \times \text{Br}_{Z_{\text{had}}} = 1 \%$
    - $[WZZ] (\text{Br}_{Z_{\text{lept}}})^3 \times \text{Br}_{Z_{\text{had}}} = 0.4 \%$
    - $[WWW] (\text{Br}_{W_{\text{lept}}})^3 = 1\%$



# Signals for heavy vectors at the LHC

- In the  $WWZ$  final state it is also worth to look at the  $WZ$  invariant-mass distribution
- With high statistics ( $100 \text{ fb}^{-1}$ ), here we can hope to see a signal even without a light axial vector

