



Probing of XYZ meson structure with near threshold pp and pA collisions

Thursday 27 September 2018 14:37 (20 minutes)

The spectroscopy of charmonium-like mesons with masses above the $2m_D$ open charm threshold has been full of surprises and remains poorly understood [1]. The currently most compelling theoretical descriptions of the mysterious XYZ mesons attribute them to hybrid structure with a tightly bound $c\bar{c}$ diquark [2] or $cq(c\bar{q})$ tetraquark [3 - 5] core that strongly couples to S-wave $D\bar{D}$ molecular-like structures. In this picture, the production of a XYZ particle in high energy hadron collisions and its decays into light hadron plus charmonium final states proceed via the core component of the meson, while decays to pairs of open charmed mesons proceed via the $D\bar{D}$ component.

These ideas have been applied with some success to the $X(3872)$ [2], where a detailed calculation finds a $c\bar{c}$ core component that is only above 5% of the time with the $D\bar{D}$ component (mostly $D^0\bar{D}^0$) accounting for the rest. In this picture, the $X(3872)$ is composed of three rather disparate components: a small charmonium-like $c\bar{c}$ core with $r_{\text{rms}} < 1$ fm, a larger D^+D^- component with $r_{\text{rms}} = \hbar/\sqrt{2\mu(B^+)} \approx 1.5$ fm and a dominant component $D^0\bar{D}^0$ with a huge, $r_{\text{rms}} = \hbar/\sqrt{2\mu(B^0)} > 9$ fm spatial extent. Here $\mu(B^+)$ and $B^+(B^0)$ denote the reduced mass for the D^+D^- ($D^0\bar{D}^0$) system and the relevant binding energy $|m_D + m_{\bar{D}} - M_{X(3872)}|$ ($B^+ = 8.2$ MeV, $B^0 < 0.3$ MeV). The different amplitudes and spatial distributions of the D^+D^- and $D^0\bar{D}^0$ components ensure that the $X(3872)$ is not an isospin eigenstate. Instead it is mostly $I = 0$, but has a significant ($\sim 25\%$) $I = 1$ component.

In the hybrid scheme, an $X(3872)$ is produced in high energy pA collisions via its compact ($r_{\text{rms}} < 1$ fm) charmonium-like structure and this rapidly mixes in a time ($t \sim \hbar/\delta M$) into a huge and fragile, mostly $D^0\bar{D}^0$ molecular-like structure. δM is the difference between the $X(3872)$ mass and that of the nearest $c\bar{c}$ mass pole core state, which we take to be that of the $\chi_{c1}(2P)$ pure charmonium state which is expected to lie about 20 ~ 30 MeV above $M_{X(3872)}$ [6, 7]. In this case, the mixing time, $\tau_{\text{mix}} \sim 5 \sim 10$ fm, is much shorter than the lifetime of $X(3872)$ which is $\tau_{X(3872)} > 150$ fm [8].

The experiments with proton-proton (pp) and proton-nuclear (pA) collisions with momentum up to 26 GeV/c and luminosity up to $10^{32} \text{ cm}^{-2}\text{s}^{-1}$ are well suited to test this picture for the $X(3872)$ and, possibly, other XYZ mesons. In near threshold production experiments in the $\sqrt{s_{NN}} \approx 8$ GeV energy range, $X(3872)$ mesons can be produced with typical kinetic energies of a few hundred MeV (i.e. with $\gamma\beta \approx 0.3$). In the case of $X(3872)$, its decay length will be greater than 50 fm while the distance scale for the $c\bar{c} \rightarrow D^0\bar{D}^0$ transition would be 2 ~ 3 fm. Since the survival probability of an $r_{\text{rms}} \sim 9$ fm “molecular” inside nuclear matter should be very small, $X(3872)$ meson production on a nuclear target with $r_{\text{rms}} \sim 5$ fm or more ($A \sim 60$ or larger) should be strongly quenched. Thus, if the hybrid picture is correct, the atomic number dependence of $X(3872)$ production at fixed $\sqrt{s_{NN}}$ should have a dramatically different behavior than that of the ψ' , which is long lived compact charmonium state.

The current experimental status of XYZ mesons together with hidden charm tetraquark candidates and present simulations what we might expect from A-dependence of $X(3872)$ mesons in pp and pA collisions are summarized.

References

- [1] S. Olsen, Front. Phys. 10 101401 (2015)
- [2] S. Takeuchi, K. Shimizu, M. Takizawa, Progr. Theor. Exp. Phys. 2015, 079203 (2015)
- [3] A. Esposito, A. Pilloni, A.D. Poloza, arXiv:1603.07667[hep-ph]
- [4] M.Y. Barabanov, A.S. Vodopyanov, S.L. Olsen, A.I. Zinchenko, Phys. Atom. Nuc. 79, 1, 126 (2016)
- [5] M.Yu. Barabanov, A.S. Vodopyanov, S.L. Olsen, Phys. Scripta 166 014019 (2015)
- [6] N. Isgur, Phys. Rev. D 32, 189 (1985)

[7] K. Olive et al. (PDG), Chin. Phys. C 38, 090001 (2014)

[8] The width of $X(3872)$ is experimentally constrained to be $\Gamma X(3872) < 1.2$ (90% CL) in S.-K. Choi et al (Belle Collaboration), Phys. Rev. D 84, 052004 (2011)

Primary author: Prof. BARABANOV, Mikhail (JINR)

Co-authors: Prof. VODOPYANOV, Alexander (JINR); Prof. STEPHEN, Olsen (UCAS)

Presenter: Prof. BARABANOV, Mikhail (JINR)

Session Classification: Parallel Session: Pheno 4

Track Classification: Particle Phenomenology