












Atomic diffusion investigation by XPCS

B. Sepiol and M. Leitner
Faculty of Physics, Universität Wien

Atomic diffusion mechanisms







H. Mehrer,
 Diffusion in
 Solids,
 Springer 2007

Our primary goal now and in future:

overcome limitations of atomistic methods (Mössbauer, QNS, NRS) to a few elements (^{57}Fe , H, Ni, Co, Ti) and fast diffusion using


X-ray Photon Correlation Spectroscopy

Single atom diffusion




M. Leitner et al., *Nature Mater.* **8**, 717 (2009)

reciprocal (001) plane




slice through (100) MC-cell:



B. Schönfeld, M.J. Portmann, S.Y. Yu,
G. Kostorz, *Acta Mat.* **47**, 1413 (1999)


1) calculating

$$g^{(2)}(\mathbf{q}, \Delta t) = \frac{\langle I(\mathbf{q}, \cdot) I(\mathbf{q}, \cdot + \Delta t) \rangle}{\langle I(\mathbf{q}, \cdot) \rangle^2}$$





2) fitting



$$g^{(2)}(\mathbf{q}, \Delta t) = 1 + \beta e^{-2\Delta t/\tau(\mathbf{q})}$$




3) verifying hypotheses $\tau(\mathbf{q}) = \tau_0 \frac{I_{\text{SRO}}(\mathbf{q})}{1 - \sum_i p_i \cos(\mathbf{s}_i \cdot \mathbf{q})}$














simple exponential decay!







only one parameter fitted !!



Local dynamics in metallic glass

Systems far from equilibrium characterized by spatial and/or temporal **heterogeneity**
e.g. dilute colloidal gels

Zr-based amorphous alloy $\text{Zr}_{65}\text{Al}_{7.5}\text{Ni}_{10}\text{Cu}_{17.5}$


metallic glasses - the paradigm of dense random packing of spheres

Calorimetric glass transition $T_g=624\text{K}$ at $2\text{K}/\text{min}$


Extrapolated quasi-stationary $T_g=605\text{K}$





T. Zhang, A. Inoue and T. Masumoto, Mater. Trans. JIM **32** (1991) 1005

Very long relaxation times





600K (327°C)





Brownian dynamics ?



Key parameters from your point of view the instrument should have like:

Source parameters:

- energy tunable energy
- pulse pattern uniform distribution preferable
- pulse length as long as possible
- polarization irrelevant

Beamline optics:

- monochromatizity as high as possible (preferred 10^{-6}) due to the increased pulse length and the lower peak intensity
- spot size about 10 μm
- degree of coherence as high as possible (lead to lower peak intensity)
- diagnostics not relevant

Detector:

- pixel size 10-20 μm
- number of pixels at least 10^6 , the more the better
- framerate 10 Hz
- accessible q-range 40° scattering angle (about 3 \AA^{-1})

Sample environment:

- temperature self-made furnaces
- external fields

Thanks to:

Lorenz M. Stadler
Bastian Pfau
Friedrich Gröstlinger
Gero Vogl

Faculty of Physics
Universität Wien

A. Fluerasu, A. Madsen at ESRF