


Weak Interactions


The Theory of GLASHOW, SALAM and WEINBERG

~ 1959-1968




(Nobel 1979)

Theory of the unified weak and electromagnetic interaction, transmitted by exchange of "intermediate vector bosons"




(a) Electromagnetic scattering



(b) Weak scattering (charged current)

heavy!

mass generated by Higgs field



(c) Weak scattering (neutral current)

Discovery of the W and Z (1983)


- To produce the heavy W and Z bosons ($m \sim 80\text{-}90\text{ GeV}$) need high energy collider!
- 1978-80: conversion of SPS proton accelerator at CERN into proton-antiproton collider
challenge: make antiproton beam!
- success!
→ first W and Z produced 1982/83

(Nobel 1984)



Carlo
Rubbia



Simon
van der
Meer



Z production at LHC





Now millions of events ...

yesterday's signal is today's background and tomorrow's calibration




Three Boson Coupling @ LEP

W/Z bosons carry electroweak charge (like colour for gluons)
→ measure rate of W pair production at LEP II






Electroweak Physics at HERA

Neutral Current (NC) interactions




Charged Current (CC) interactions




Weak interactions are "left-handed"

- lefthanded electrons interact (CC)




- righthanded electrons do not!



- cross section linearly proportional to polarization


$$\sigma_{polCC}^{e^{\pm}p} = (1 \pm P_e) \cdot \sigma_{unpolCC}^{e^{\pm}p}$$



It works!

Electroweak Unification

H1 and ZEUS




Strength of weak and electromagnetic forces become similar at scale $Q^2 \sim M_W^2$


$$\frac{d^2\sigma_{NC}}{dQ^2 dx} \sim \alpha^2 \frac{1}{Q^4} \frac{1}{x} \Phi_{NC}(x, Q^2)$$

$$\frac{d^2\sigma_{CC}}{dQ^2 dx} \sim G_F^2 \left(\frac{M_W^2}{M_W^2 + Q^2} \right)^2 \frac{1}{x} \Phi_{CC}(x, Q^2)$$

The Quest for Unification of Forces




α_s running and Grand Unification



?

with SUSY (see later):



hep-ph/0407067 B.Allanach ... P.Zerwas

Antimatter

relativistic Schrödinger equation
(Dirac equation)

two solutions:

one with positive, one with negative energy

Dirac: interpret negative solution as **antiparticle**

1932 antielectrons (positrons) found in conversion
of energy into matter

1995 antihydrogen consisting of antiprotons and
positrons produced at CERN



P.A.M.
Dirac
(Nobel 1933)

C.D. Anderson
(Nobel 1936)




In principle: antiworld can be built from antimatter

In practice: produced only in accelerators and
in cosmic rays



Pair Production



e.g. $\gamma \rightarrow e^+ + e^-$



Equal amounts of matter and antimatter are produced
when radiation is converted to matter

Annihilation

$$e^{+} + e^{-} \rightarrow 2hf$$





Antimatter can be produced.
It annihilates with matter to produce radiation

The Matter Antimatter Puzzle

Why does the Universe look like
this

not

that?




As far as we can see in universe, no large-scale antimatter.
-> need CP violation!

The Matter Antimatter Puzzle

Early Universe


- > particles, anti-particles and photons in thermal equilibrium
- colliding, annihilating, being re-created etc.



Slight difference in fundamental interactions between matter and antimatter ("CP violation") ?


-> matter slightly more likely to survive

Ratio of baryons (e.g. p, n) to photons today tells us about this asymmetry - it is about $1:10^9$




CP symmetry

graphics: M.C. Escher




Parity
(reflection)



Charge
Conjugation ↓ (black → white)





P
→



Like weak interaction, symmetric under CP (at first sight!)
Can there be small deviations from this symmetry?

CP violation in B meson decays



$$Asymmetry(t) = \frac{\bar{B}_d^0 - B_d^0}{\bar{B}_d^0 + B_d^0}$$

Simply count decays as
function of t !

CP violation in B meson decays

Example: measurement from BaBar at SLAC

(also Belle
at KEK)

B and anti-B
are indeed
different


(also found
earlier for
K decays:)



Val L.
Fitch

James W. Cronin (Nobel 1980)

18.-20.7.18



data taking stopped

Belle/Super-Belle continuing (DESY!)


A. Geiser, Particle Physics

Weak
Interactions
violate CP!



M. Kobayashi T. Maskawa
(Nobel 2008)

The Mystery of Mass




The Mass (BEH) Mechanism

P. Higgs et al. (1964-66,71)


Brout, Englert, Guralnik, Hagen, Kibble, ...

many subvariants

which is right?




Peter Higgs



François Englert

(Nobel 2013)



■ source: viXra blog

Fermion Mass from Higgs field?


very brilliant scientist (fermion)
works with speed of light!

-> "massless"



room = vacuum

people = Higgs vacuum expectation value



Fermion Mass from Higgs field?


scientist becomes famous!
enters room with people






Fermion Mass from Higgs field?

people cluster around him
hamper his movement/working speed
→ he becomes "massive"!


**Why are some
fermions so much
more famous
than others?**



Neutrino oscillations: neutrinos are massive!




1998



Takaaki Kajita Arthur McDonald
(Nobel 2015)

What do we know about Neutrino mass?



Why???

- are the masses of Dirac type (generated by Higgs)?
or of Majorana type (ν 's are their own antiparticles,
masses have non-Standard Model origin)?
- CP violation?

possibly first evidence
for physics
beyond Standard Model