


LUXE Background Study in Simulation

Oleksandr Borysov



LUXE Meeting
July 15, 2019

LUXE geometry in Geant4



- Check background in OPPP detectors: trackers and calorimeters;
- Optimize detectors position, shielding, beam pipes and windows;
- Establish a benchmark in a simple geometry for comparison with more detailed implementation.




Sketch and Geant4



Beam Dump with Hole for Photons




Fraction of photons inside the circle as a function of its radius for different distances from the target




Front view of the beam dump through the beam pipe




Tracking Planes



No Target, 1.4 T



Performance with test settings






50k e-


Simulated 50k events recording any track that enters detectors volume
1 was registered;

```
*****
*      Row      *      pdg      *      E      *      vtxx      *      vtxy      *      vtxz      *      px      *      py      *      pz      *
*****
*      0      *      -11      *      1.1105072      *      0.6417944      *      -1.773730      *      -3752.636      *      0.4198077      *      0.0007512      *      1.0286506      *
*****
```

Bremsstrahlung photons 22.5 m from the collimator (beam dump)



Test with 17.5 GeV e- and 6 T



```


root [2] oppphits->Scan("*")
*****
*      Row      * Instance *   eventid *   detid *   layerid *   cellx *   celly *   edep *   hitid * track lis *
*****
*      0 *      0 *      0 *      1 *      2 *      830 *   1003 * 0.0001291 *      2 *      1 *
*      1 *      0 *      0 *      1 *      1 *     1301 *   1003 * 9.361e-05 *      1 *      1 *
*      2 *      0 *      0 *      1 *      0 *     1771 *   1003 * 0.0001062 *      0 *      1 *
*****

```


Track in detector volume

10.72M e-

All tracks entering detector
container volume






e+e- tracks entering detector
container volume




Hits with origin in collimator


10.72 M electrons of
17.5 GeV
with 35 μm tungsten target
were simulated




Hits with origin in spectrometer magnet




Cleaning beam after the dump




e-, 17.5 GeV





Zoom





Z distribution of charged vertexes





13.54 M e^-



X vs Z Distribution of all and charged





Beam Pipe D = 10 cm




Electron energy (Ev): 1.75×10^{10}
Magnetic field (T): 2
Curvature radius (m): 29.1869
Displacement at the exit from the magnet (m): 0.017136
Angle (rad): 0.0342687
Displacement at 2.6 m from the magnet (m): 0.10627

Beam Pipe D = 10 cm. Tracks in detector volume.


3M e-;
1m < Z < 2.5m



Hits xz distribution e-,e+







hhvtxxy



Beam Pipe D = 10 cm. Vertexes of the e⁺, e⁻ tracks in detector volume.






30M e⁻;

1m < Z < 2.5m; (Spectrometer magnet)



Beam Pipe D = 10 cm. Vertexes of the e⁺, e⁻ tracks in detector volume.




30M e⁻;
2.5m < Z < 6m; (detectors and beam pipe)



Beam Pipe D = 10 cm. Vertexes of the e⁺, e⁻ tracks in detector volume.

30M e⁻;




-1.2m < Z < -1m; (Shielding)



Beam Pipe $D = 5$ cm, Geometry with a magnet between beam dump (collimator) and shielding. Vertexes of the e^+ , e^- tracks in detector volume.



$\sim 3M$ e^- ;

Statistically can be compared with the simulation on slide 18



Backup

Tungsten Target, 1.4 T, ~ 30 e-



Sketches of Bremsstrahlung Area

