

Searches for an exotic decay of the Higgs boson to a pair of pseudoscalars in CMS

Danyer Pérez Adán

On behalf of the CMS collaboration



- Motivation
- $h(125) \rightarrow a_1 a_1$ searches in CMS at 13 TeV
 - Resolved Topology
 - $h(125) \rightarrow a_1 a_1 \rightarrow \tau\tau\mu\mu$
 - $h(125) \rightarrow a_1 a_1 \rightarrow bb\tau\tau$
 - $h(125) \rightarrow a_1 a_1 \rightarrow bb\mu\mu$
 - Boosted Topology
 - $h(125) \rightarrow a_1 a_1 \rightarrow \mu\mu\mu\mu$
 - $h(125) \rightarrow a_1 a_1 \rightarrow \tau\tau\tau\tau$
- Summary of the results
- Conclusion

Motivation

The 125 GeV Higgs boson

$$BR_{BSM} < 34\%$$

Combined ATLAS and CMS coupling analysis for the Run1 Data

[arXiv:1606.02266]

2HDM

- One of the simplest possible extensions of the SM
- They play an important role in:
 - **Supersymmetry:** *holomorphy and cancellation of anomalies*
 - **Axion models:** *imposing Peccei and Quinn symmetry only possible if there are two Higgs doublets*
 - **Baryon asymmetry:** *it could contain additional sources of CP violation*

However, 2HDMs are by now strongly constrained from existing data

2HDM+S

- The current constraints can be avoided by:
 - assuming that the 2HDM is in the decoupling limit
the couplings of $h(125)$ become SM-like or very close to SM-like
 - adding one complex scalar singlet
it only couples to $H_{1,2}$ and it is allowed to have small mixing with these

$$\alpha \rightarrow \beta - \frac{\pi}{2}$$

$$S = \frac{1}{\sqrt{2}}(S_R + iS_I)$$

Motivation

Light Pseudoscalar (a_1)

- The 2HDM+S contains 7 physical states:

2 charged (H^+, H^-), 3 CP-even (h_1, h_2, h_3) and 2 CP-odd (a_1, a_2)

a_1 is the mostly-singlet-like pseudoscalar $a_1 = \cos(\theta_{a_1})S_1 + \sin(\theta_{a_1})A$ $\theta_{a_1} \ll 1$

- There exist scenarios in which a_1 is lighter than the SM-like Higgs, namely:

$$m_{a_1} < \frac{m_{h(125)}}{2} \approx 63 \text{ GeV}$$


in this case, there are exotic Higgs decays of the form: $h(125) \rightarrow a_1 a_1 \rightarrow X\bar{X}Y\bar{Y}$

Table: Types of fermion couplings (w/o FCNC)

2HDM	u-type	d-type	lepton
Type-I	H_2	H_2	H_2
Type-II	H_2	H_1	H_1
Type-III	H_2	H_2	H_1
Type-IV	H_2	H_1	H_2

$BR(a_1 \rightarrow X\bar{X})$ for Type II

[arXiv:1312.4992]



CMS Searches

Resolved Topology




$h(125) \rightarrow a_1 a_1 \rightarrow \tau\tau\mu\mu$

[arXiv:1805.04865]

- Mass range probed: $15.0 < m_{a_1} < 62.5$ GeV
- Production modes of $h(125)$ considered: gluon fusion (**ggH**) and vector boson fusion (**VBF**)
- $h(125) \rightarrow a_1 a_1 \rightarrow \tau\tau\tau\tau$ events are also treated as part of the signal by means of the relation:

$$\frac{B(a_1 \rightarrow \mu\mu)}{B(a_1 \rightarrow \tau\tau)} = \frac{m_\mu^2 \sqrt{1 - (2m_\mu/m_{a_1})^2}}{m_\tau^2 \sqrt{1 - (2m_\tau/m_{a_1})^2}} \quad (1)$$




- Four different final states were covered: $\mu\mu + e\mu$, $\mu\mu + e\tau_h$, $\mu\mu + \mu\tau_h$ and $\mu\mu + \tau_h\tau_h$
- Signal extraction: unbinned maximum-likelihood fit based on the $m_{\mu\mu}$ distribution
- The shape and yield of the backgrounds are estimated from data



- Mass range probed: $15 < m_{a_1} < 60$ GeV
- Production modes of $h(125)$ considered: **ggH**, **VBF** and associated vector boson (**VH**)
- Three different final states were covered: $b_{\text{tagged} \geq 1} + e\mu$, $b_{\text{tagged} \geq 1} + e\tau_h$ and $b_{\text{tagged} \geq 1} + \mu\tau_h$
- Categorization (4 cats.) according to the variables:

$$m_{b\tau\tau}^{\text{vis}} \quad m_T(e(\mu), \vec{p}_T^{\text{miss}}) \quad m_T(\mu(\tau_h), \vec{p}_T^{\text{miss}}) \quad D_\zeta [^*]$$

- Signal extraction: binned maximum-likelihood fit based on the $m_{\tau\tau}^{\text{vis}}$ distribution
- The backgrounds are estimated from a combination of simulation and data



$$[^*] D_\zeta \equiv \vec{p}_T^{\text{miss}} \cdot \frac{\vec{\zeta}}{|\vec{\zeta}|} - 0.85 (\vec{p}_{T,1} + \vec{p}_{T,2}) \cdot \frac{\vec{\zeta}}{|\vec{\zeta}|}$$

$\vec{\zeta} \rightarrow$ bisector of the transverse momenta of the two τ candidates

- Mass range probed: $20.0 < m_{a_1} < 62.5$ GeV
- Production modes of $h(125)$ considered: **ggH** and **VBF**
- Final state: $b_{tagged} b_{tagged} + \mu\mu$

- Events are selected if they have $\chi^2 < 5$:

$$\chi^2 = \chi_{bb}^2 + \chi_h^2 \quad \chi_{bb} = \frac{m_{bb} - m_{\mu\mu}}{\sigma_{bb}} \quad \chi_h = \frac{m_{\mu\mu} - m_h}{\sigma_h}$$




- Categorization (3 cats.) according to the b tagging discriminator value of one of the jets:

Tight-Tight

Tight-Medium

Tight-Loose



- Signal extraction: unbinned maximum-likelihood fit based on the $m_{\mu\mu}$ distribution
- The background estimation fully relies on data by using the discrete profiling method



$h(125) \rightarrow a_1 a_1 \rightarrow \mu\mu\mu\mu$

[arXiv:1812.00380]

- Mass range probed: $0.25 < m_{a_1} < 3.55$ GeV
 - Production mode of $h(125)$ considered: **ggH**
 - Final state: $\mu\mu + \mu\mu$
 - Events are selected if they fulfill the relation $m_{(\mu\mu)1} \simeq m_{(\mu\mu)2}$, as shown in the figure
 - Signal extraction: unbinned maximum-likelihood fit based on the 2D $m_{(\mu\mu)1}$ vs. $m_{(\mu\mu)2}$ distribution
 - The estimation of the main background contribution ($b\bar{b}$) is from data
 - The prompt double J/ψ and electroweak backgrounds are modelled from data and simulation respectively
- ★ **Total expected background events:** $9.90 \pm 1.24(stat) \pm 1.84(syst)$ **Observed events:** 13






CMS Searches

Boosted Topology


$h(125) \rightarrow a_1 a_1 \rightarrow \tau\tau\tau\tau$ **NEW!**

[CMS-PAS-HIG-18-006]

- Mass range probed: $4 < m_{a_1} < 15$ GeV
- Production mode of $h(125)$ considered: **ggH, VBF, VH** and top quark pair associated production (**ttH**)
- $h(125) \rightarrow a_1 a_1 \rightarrow \mu\mu\tau\tau$ events are also included as part of the signal by using Eq. 1
- Final state covered: $\mu^\pm \tau_{one-prong}^\mp + \mu^\pm \tau_{one-prong}^\mp$ (See sketch)
- Events are selected if they have exactly two isolated $\mu^\pm - trk^\mp$ pairs within a cone of size $\Delta R = 0.5$
- Signal extraction: binned maximum-likelihood fit based on the 2D $m_{(\mu-trk)_1}$ vs. $m_{(\mu-trk)_2}$ distribution
- The background modelling is based on data, although some auxiliary tests also include simulation



Summary of the results



Conclusion

- Many $h(125) \rightarrow a_1 a_1$ searches performed in different decay channels and final states
- The searches cover a large variety of 2HDM+S models
 - Almost all possible masses for the a_1 boson have been probed, having to deal with different boosted regimes
 - Scenarios comprising all types of fermion coupling have been tested
- No sign of $h(125) \rightarrow a_1 a_1$ decay yet ...
- Limits are becoming more stringent as more data is added
- Other interesting analyses ongoing:
 - $h(125) \rightarrow a_1 a_1 \rightarrow bbbb$ (first time in CMS)
 - $h(125) \rightarrow a_1 a_1 \rightarrow \tau\tau\mu\mu$ (for light a_1 masses)






Conclusion

- Many $h(125) \rightarrow a_1 a_1$ searches performed in different decay channels and final states
- The searches cover a large variety of 2HDM+S models
 - Almost all possible masses for the a_1 boson have been probed, having to deal with different boosted regimes
 - Scenarios comprising all types of fermion coupling have been tested
- No sign of $h(125) \rightarrow a_1 a_1$ decay yet ...
- Limits are becoming more stringent as more data is added
- Other interesting analyses ongoing:
 - $h(125) \rightarrow a_1 a_1 \rightarrow bbbb$ (first time in CMS)
 - $h(125) \rightarrow a_1 a_1 \rightarrow \tau\tau\mu\mu$ (for light a_1 masses)





Thanks for your attention!

Backup









$BR(a_1 \rightarrow X\bar{X})$ for types of 2HDM+S models






$$h(125) \rightarrow a_1 a_1 \rightarrow \tau\tau\mu\mu$$




$$h(125) \rightarrow a_1 a_1 \rightarrow bb\tau\tau$$



$$h(125) \rightarrow a_1 a_1 \rightarrow b b \mu \mu$$



- ◆ Data, 35.9 fb⁻¹ @ 13 TeV
- Top
- Z/ γ^* ($\rightarrow ll$) + jets
- Diboson
- Simulation stat. unc.
- $m_{a_1} = 20$ GeV
- $m_{a_1} = 40$ GeV
- $m_{a_1} = 60$ GeV



$$h(125) \rightarrow a_1 a_1 \rightarrow \tau\tau\tau\tau$$

Background model constructed as:

$$f_{2D}(i, j) = C(i, j) \cdot (f_{1D}(i) \cdot f_{1D}(j)) \quad (2)$$

