## Luminescence of ice as a new detection channel for neutrino telescopes

Tuesday 13 July 2021 13:18 (12 minutes)

Natural water and ice are currently used as optical detection media in large scale neutrino telescopes, such as IceCube, KM3NeT/ANTARES and GVD. When charged particles, such as those produced by high energy neutrino interactions, pass through ice or water at relativistic speeds they induce Cherenkov light emission. This is detected by the optical modules of neutrino telescopes. However, slower moving particles, including potential exotic matter such as Magnetic Monopoles or Q-balls, cannot be detected using this channel.

A new kind of signature can be detected by using light emission from luminescence in water or ice. This detection channel enables searches for exotic particles which are too slow to emit Cherenkov light and currently cannot be probed by the largest particle detectors in the world, i.e. neutrino telescopes. Luminescence light is highly dependent on the ice structure, impurities, pressure and temperature which demands a comprehensive study.

Luminescence light is induced by highly ionizing particles passing through a medium and exciting the surrounding matter. To utilise this new detection channel in neutrino telescopes, laboratory measurements using water and ice as well as an in-situ measurement in Antarctic ice were performed. The experiments as well as the measurement results will be presented covering light yields, spectra and decay times. The impact on searches for new physics with neutrino telescopes will be discussed.

## Keywords

particle detection; neutrino telescopes; luminescence; Q-balls; magnetic monopoles

## Collaboration

## other Collaboration

Subcategory

Experimental Methods & Instrumentation

Primary author: Dr POLLMANN, Anna (Universität Wuppertal)Presenter: Dr POLLMANN, Anna (Universität Wuppertal)Session Classification: Discussion

Track Classification: Scientific Field: NU | Neutrinos & Muons