

The Southern Wide-field Gamma-ray Observatory reach for Primordial Black Hole evaporation

Friday 16 July 2021 19:18 (12 minutes)

The search for Primordial Black Hole (PBH) signatures is very broad in techniques and the origin of these signatures. Searches for imprints of evaporation involve several observables such as the Extragalactic Gamma-Ray background or direct measurement of different species of cosmic rays. Using these observables, one can put very tight constraints on the PBH number density in a mass range $\sim 10^{14}$ g. To perform direct observations of the evaporation of these PBHs, one needs to perform observations in the Very High Energy gamma-ray range, either using Imaging Atmospheric Cherenkov telescopes or wide field of view gamma-ray arrays. The Southern Wide-field Gamma-ray Observatory is a projected ground-based gamma-ray detector that will be located in the Southern Hemisphere and it is now in its design phase. In this contribution, we will show the anticipated sensitivity for PBH evaporation achievable by SWGO.

Keywords

Collaboration

SWGO

other Collaboration

Subcategory

Future projects

Primary author: LOPEZ-COTO, Ruben (INFN Padova)

Co-authors: DORO, Michele (University and INFN Padova); DE ANGELIS, Alessandro; Prof. MARIOTTI, Mose (University of Padova); HARDING, Pat (Los Alamos National Laboratory); FOR THE SWGO COLLABORATION

Presenter: LOPEZ-COTO, Ruben (INFN Padova)

Session Classification: Discussion

Track Classification: Scientific Field: GAI | Gamma Ray Indirect