



QCD jet production at a high energy muon collider

Yang Ma

Pittsburgh Particle-physics, Astrophysics, and Cosmology Center,
Department of Physics and Astronomy,
University of Pittsburgh, PA 15260, USA

July 26, 2021

In collaboration with **Tao Han** and **Keping Xie**

[T. Han, Y. Ma, K.Xie 2007.14300]

[T. Han, Y. Ma, K.Xie 2103.09844]

What is the PDF of a lepton?


“Equivalent photon approximation (EPA)”

[C. F. von Weizsacker, Z. Phys. 88, 612 (1934)]

Treat photon as a parton constituent in the electron [E. J. Williams, Phys. Rev. 45, 729 (1934)]

$$\sigma(\ell^- + a \rightarrow \ell^- + X) = \int dx f_{\gamma/\ell} \hat{\sigma}(\gamma a \rightarrow X)$$

$$f_{\gamma/\ell, \text{EPA}}(x_\gamma, Q^2) = \frac{\alpha}{2\pi} \frac{1 + (1 - x_\gamma)^2}{x_\gamma} \ln \frac{Q^2}{m_\ell^2}$$



Extra terms:

[Frixione, Mangano, Nason, Ridolfi 2103.09844]

[Budnev, Ginzburg, Meledin, Serbo, Phys. Rept.(1975)]

Applications at muon collider


- Production cross sections

$$\sigma(\ell^+ \ell^- \rightarrow F + X) = \int_{\tau_0}^1 d\tau \sum_{ij} \frac{d\mathcal{L}_{ij}}{d\tau} \hat{\sigma}(ij \rightarrow F), \quad \tau = \hat{s}/s$$


- Partonic luminosities

$$\frac{d\mathcal{L}_{ij}}{d\tau} = \frac{1}{1 + \delta_{ij}} \int_{\tau}^1 \frac{d\xi}{\xi} \left[f_i(\xi, Q^2) f_j\left(\frac{\tau}{\xi}, Q^2\right) + (i \leftrightarrow j) \right]$$

$\gamma\gamma$ collision



ISR



A possible high-energy lepton collider: Why?

Why lepton colliders?

- **Leptons** are the ideal probes of short-distance physics
 - Cleaner background comparing to hadron colliders
 - High-energy physics probed with much smaller collider energy
- **ee colliders**
 - A glorious past: discovery of charm, τ , and gluon
 - Important future: Precision EW constraints on BSM physics, Higgs physics
- **Muon colliders**
 - A *s*-channel Higgs factory: Higgs production enhanced by $m_\mu^2/m_e^2 \sim 40000$
 - Direct measurements on y_μ and Γ_H
 - **Multi-TeV muon colliders**: Less radiations than electron
 - Center of mass energy 3–15 TeV and the more speculative $E_{\text{cm}} = 30$ TeV
 - New particle mass coverage $M \sim (0.5 - 1)E_{\text{cm}}$
 - Great accuracies for WWH , $WWHH$, H^3 , H^4
 - ...

Muon Collider Physics Potential Pillars

Direct search of heavy particles

SUSY-inspired, WIMP, VBF production, $2 \rightarrow 1$

High rate indirect probes

Higgs single and self-couplings, rare Higgs decays, exotic decays

High energy probes

difermion, diboson, EFT, Higgs compositeness

A high-energy muon collider at first glance

What are the dominant processes at a high-energy muon collider?



- Leading-order: $\mu^+\mu^- \rightarrow \mu^+\mu^-$, $\tau^+\tau^-$, $q\bar{q}$, W^+W^- , and $\gamma\mu \rightarrow \gamma\mu$
- $\gamma\gamma$ scatterings: $\gamma\gamma \rightarrow \tau^+\tau^-$, $q\bar{q}$, W^+W^-

Need some cuts:

- Detector angle: $\theta > 5^\circ$ (10°) $\iff |\eta| < 3.13$ (2.44)
- Threshold: $m_{ij} > 20$ GeV
- Need a p_T cut to separate from the nonperturbative hadronic production

[Chen, Barklow, and Peskin, hep-ph/9305247; Drees and Godbole, PRL 67, 1189, T. Barklow, et al, LCD-2011-020]

$$p_T > (4 + \sqrt{s}/3 \text{ TeV}) \text{ GeV}$$




Go beyond the EPA at a high-energy muon collider

We have been doing:

- $\ell^+\ell^-$ annihilation




- EPA and ISR



- “Effective W Approx.” (EWA)

[G. Kane, W. Repko, and W. Rolnick, PLB 148 (1984) 367]


[S. Dawson, NPB 249 (1985) 42]




We will add:

[T. Han, Y. Ma, K.Xie 2007.14300, 2103.09844]

- Above μ_{QCD} : $\text{QED} \otimes \text{QCD}$
 q/g emerge



- Above $\mu_{\text{EW}} = M_Z$: $\text{EW} \otimes \text{QCD}$
EW partons emerge



In the end, everything is parton, i.e. **the full SM PDFs.**

The PDFs for a muon collider

■ QED \otimes QCD PDFs:

$$f_{\mu_{\text{val}}}, f_{\gamma}, f_{\ell_{\text{sea}}}, f_q, f_g$$

- Scale uncertainty: 20% for $f_{g/\mu}$

- The averaged momentum fractions

$$\langle x_i \rangle = \int x f_i(x) dx$$

$Q(\mu^\pm)$	μ_{val}	γ	ℓ_{sea}	q	g
30 GeV	98.2	1.72	0.019	0.024	0.0043
50 GeV	98.0	1.87	0.023	0.029	0.0051
M_Z	97.9	2.06	0.028	0.035	0.0062



■ EW PDFs: All SM particles

Q	μ	$\gamma, Z, \gamma/Z$	W^\pm	ν	ℓ_{sea}	q	g
M_Z	97.9	2.06	0	0	0.028	0.035	0.0062
3 TeV	91.5	3.61	1.10	3.59	0.069	0.13	0.019
5 TeV	89.9	3.82	1.24	4.82	0.077	0.16	0.022

- Scale uncertainty: $\sim 20\%$ between $Q = 3$ TeV and $Q = 5$ TeV

- The EW correction is not small: $\sim 100\%$ for $f_{d/\mu}$ due to **relatively large SU(2) gauge coupling**.

- W_L does not evolve





Parton luminosities at a possible muon collider

Consider a 3 TeV and a 10 TeV machine

■ Partonic luminosities for

$\mu^+\mu^-$, $\gamma\mu$, $\gamma\gamma$, qq , γq , γg , gq , and gg



- The partonic luminosity of $\gamma g + \gamma q$ is $\sim 20\%$ of the $\gamma\gamma$ one
- The partonic luminosities of qq , gq , and gg are $\sim 0.5\%$ of the $\gamma\gamma$ one
- Given the stronger QCD coupling, **sizable QCD cross sections are expected.**
- Scale uncertainty is $\sim 20\%$ ($\sim 50\%$) for photon (gluon) initiated processes.

Jet production of possible lepton colliders (I)

- Large photon induced non-perturbative hadronic production


[Drees and Godbole, PRL 67 1189, hep-ph/9203219]

[Chen, Barklow, and Peskin, hep-ph/9305247; Godbole, Grau, Mohan, Pancheri, SrivastavaNuovo Cim. C 034S1]


- $\sigma_{\gamma\gamma}$ may reach micro-barns level at TeV c.m. energies
- $\sigma_{\ell\ell}$ may reach nano-barns, after folding in the $\gamma\gamma$ luminosity

- The events populate at low p_T regime

So we can separate from this non-perturbative range via a p_T cut.



(a) Pythia sample



(b) SLAC sample

[T. Barklow, D. Dannheim, M. O. Sahin, and D. Schulte, LCD-2011-020]

Jet production at a possible lepton collider (II)

- Low- p_T range: photon induced non-perturbative hadronic production



[Chen, Barklow, and Peskin, hep-ph/9305247; Drees and Godbole, PRL 67, 1189, T. Barklow, et al, LCD-2011-020]

- High- p_T range [$p_T > (4 + \sqrt{s}/3 \text{ TeV}) \text{ GeV}$]: perturbatively computable

$$\gamma\gamma \rightarrow q\bar{q}, \gamma g \rightarrow q\bar{q}, \gamma q \rightarrow gq,$$

$$qq \rightarrow qq (gg), gq \rightarrow gq \text{ and } gg \rightarrow gg (q\bar{q}).$$

- $Q = \sqrt{\hat{s}}/2$, due to large $\alpha_s \ln(Q^2)$, a 30 ~ 40% enhancement if $Q = \sqrt{\hat{s}}$




- Including the QCD contribution leads to much larger total cross section.
- gg initiated cross sections are large for its large multiplicity;
- gq initiated cross sections are large for its large luminosity.
- $\gamma\gamma$ initiated cross sections here are smaller than the EPA results.

Refresh the picture of high-energy muon colliders




What is the dominant process at a high-energy muon collider?

- Quark/gluon initiated jet production dominates

Before:



After:





Di-jet distributions at a muon collider

Rather a conservative set up: $\theta = 10^\circ$




■ Some physics:

Two different mechanisms: $\mu^+\mu^-$ **annihilation** VS **Fusion processes**

- Annihilation is more than 2 orders of magnitude smaller than fusion process.
- Annihilation peaks at $m_{ij} \sim \sqrt{s}$;
- Fusion processes peak near m_{ij} threshold.
- Annihilation is very central, spread out due to ISR;
- Fusion processes spread out, especially for γq and γg initiated ones.



Inclusive jet distributions at a muon collider




- Jet production dominates over WW production until $p_T > 60$ GeV;
- WW production takes over around energy ~ 200 GeV.
- QCD contributions are mostly forward-backward; $\gamma\gamma$, γq , and γg initiated processes are more isotropic.

An EW version of HE LHC




- All SM particles are partons when the machine energy is high
- We are able to determine the partons with their different polarizations

The EW parton luminosities of a 30 TeV muon collider







[T. Han, Y. Ma, K.Xie 2007.14300]

One example: $t\bar{t}$ production at a muon collider




Other processes: W^+W^- , ZH , HH , $t\bar{t}H$



The full picture: Semi-inclusive processes

Just like in hadronic collisions: $\mu^+\mu^- \rightarrow$ exclusive particles + remnants



[T. Han, Y. Ma, K.Xie 2007.14300]

Some observations:

- The annihilations decrease as $1/s$.
- ISR needs to be considered, which can give over 10% enhancement.
- The fusions increase as $\ln^p(s)$, which take over at high energies.
- The large collinear logarithm $\ln(s/m_\mu^2)$ needs to be resummed, set $Q = \sqrt{\hat{s}}/2$.

Summary and prospects

EWPDF is important and necessary:

- At very high energies, the collinear splittings dominate. **All SM particles should be treated as partons that described by proper PDFs.**
 - The large collinear logarithm needs to be resummed via solving the DGLAP equations, so the **QCD partons (quarks and gluons) emerge.**
 - When $Q > M_Z$, the EW splittings are activated: the EW partons appear, and the existing $\text{QED} \otimes \text{QCD}$ PDFs may receive big corrections.

A high-energy muon collider is an EW version of HE LHC

- There are many things to work on: SUSY, DM, Higgs, etc.
- Two classes of processes: $\mu^+ \mu^-$ annihilation VS fusions

[T. Han, Y. Ma, K.Xie 2007.14300]

■ The main background of is the jet production:

- Low p_T range: non-perturbative $\gamma\gamma$ initiated hadronic production dominates

[Chen, Barklow, and Peskin, hep-ph/9305247; Drees and Godbole, PRL 67, 1189, T. Barklow, etal, LCD-2011-020]

- High p_T range, q and g initiated jet production dominates

[T. Han, Y. Ma, K.Xie 2103.09844]

- EWPDF allows to determine the contributions from different partons and their different polarizations.