



Study of e^+e^- annihilation to hadrons with SND at VEPP-2000


Vladimir Druzhinin for SND Collaboration

Budker INP, Novosibirsk, Russia

July 29, 2021

EPS-HEP2021

SND detector




1 – beam pipe, 2 – tracking system, 3 – aerogel Cherenkov counter, 4 – NaI(Tl) crystals, 5 – phototriodes, 6 – iron muon absorber, 7–9 – muon detector, 10 – focusing solenoids.

SND collected data at the VEPP-2M (1996-2000) and VEPP-2000 (2010-2013, 2016-...)

Main physics task of SND is study of all possible processes of e^+e^- annihilation into hadrons below 2 GeV.

- ✓ The total hadronic cross section, which is calculated as a sum of exclusive cross sections.
- ✓ Study of hadronization (dynamics of exclusive processes).
 - Properties of excited vector mesons of the ρ , ω , ϕ families
 - Development of MC event generator for $e^+e^- \rightarrow$ hadrons below 2 GeV.

VEPP-2000 e^+e^- collider




VEPP-2000 parameters:

- c.m. energy $E=0.3-2.0$ GeV
- circumference – 24.4 m
- round beam optics
- Luminosity at $E=1.8$ ГэВ
 $1 \times 10^{32} \text{ cm}^{-2} \text{ sec}^{-1}$ (project)
 $4 \times 10^{31} \text{ cm}^{-2} \text{ sec}^{-1}$ (achieved)
- Two detectors: SND and CMD-3

2010-2013 – experiments, 70 pb^{-1}
2013-2016 – upgrade, new injector
2016- ... – experiments, 300 pb^{-1}

SND data

~20 hadronic processes are currently under analysis




	Below ϕ	Near ϕ	Above ϕ
$IL, \text{ pb}^{-1}$	77	31	259.0
$E_{\text{cm}}, \text{ GeV}$	0.30- 0.97	0.98- 1.05	1.05- 2.00

- $e^+e^- \rightarrow \pi^+\pi^-$
- $e^+e^- \rightarrow K^+K^-\pi^0$
- $e^+e^- \rightarrow n\bar{n}$
- $e^+e^- \rightarrow \pi^0\eta\gamma$
- $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
- $e^+e^- \rightarrow \eta\eta\gamma$


e/π separation parameter

$$e^+ e^- \rightarrow \pi^+ \pi^-$$

JHEP 2021,113 (2021)



The analysis is based on 4.7 pb⁻¹ data recorded in 2013 (1/10 full SND data set)






Systematic uncertainty on the cross section (%)

Source	< 0.6 GeV	0.6 - 0.9 GeV
Trigger	0.5	0.5
Selection criteria	0.6	0.6
e/π separation	0.5	0.1
Nucl. interaction	0.2	0.2
Theory	0.2	0.2
Total	0.9	0.8

	SND @ VEPP-2000	SND @ VEPP-2M	PDG
M_ρ , MeV	$775.4 \pm 0.5 \pm 0.4$	$775.6 \pm 0.4 \pm 0.5$	775.3 ± 0.3
Γ_ρ , MeV	$145.7 \pm 0.7 \pm 1.0$	$146.1 \pm 0.8 \pm 1.5$	147.8 ± 0.9
$B_{\rho ee} \times 10^5$	$4.89 \pm 0.2 \pm 0.4$	$4.88 \pm 0.2 \pm 0.6$	4.72 ± 0.5
$B_{\omega\pi\pi}$ %	$1.77 \pm 0.08 \pm 0.02$	$1.66 \pm 0.08 \pm 0.05$	1.53 ± 0.06

$$e^+e^- \rightarrow \pi^+\pi^-$$






$$0.53 < \sqrt{s} < 0.88 \text{ GeV}$$


	$a_\mu(\pi^+\pi^-) \times 10^{10}$
SND & VEPP-2000	$411.8 \pm 1.0 \pm 3.7$
SND & VEPP-2M	$408.9 \pm 1.3 \pm 5.3$
BABAR	$414.9 \pm 0.3 \pm 2.1$

$$e^+ e^- \rightarrow n \bar{n}$$

- This process near threshold was previously measured by FENICE and SND using the 2011-2012 dataset.
- The new measurement is based on the 2017, 2019 data and uses time measurement in the calorimeter.
- The time distribution is fitted by a sum of distributions for signal, cosmic background, and beam + e^+e^- annihilation background.
- Our new result is lower than the previous. The reasons are underestimated beam background and incorrect MC simulation.



The systematic uncertainty is 10%






Neutron electromagnetic form factors

From the measured cross section, we determine effective form factor

$$|F|^2 = \frac{|G_M|^2 + \frac{2m_n^2}{s}|G_E|^2}{1 + \frac{2m_n^2}{s}}$$


$$\sigma(e^+e^- \rightarrow n\bar{n}) = \frac{\alpha^2\beta}{4s} \left[|G_M|^2(1 + \cos^2\theta) + \frac{4m_n^2}{s}|G_E|^2\sin^2\theta \right]$$

From analysis of the antineutron polar-angle distribution we determine the ratio of the form factors



$e^+e^- \rightarrow \pi^+\pi^-\pi^0$ cross section




EPJ C **80** (2020) 993



- ✓ Both SND measurements are consistent with each other and with the the BABAR measurement.
- ✓ Two peaks in the cross section corresponds to the) $\omega' \equiv \omega(1420)$ and $\omega'' \equiv \omega(1650)$ resonances.
- ✓ The systematic uncertainty on the cross section is 4.4%.


The previous SND measurement [J. Exp. Theor. Phys. 121, 27 (2015)] is based on 2011 data set. The 2012 data set has been added.

$e^+e^- \rightarrow \pi^+\pi^-\pi^0$ dynamics



- We analyze the two-dimensional distribution of the charged-pion momenta and the $\pi^+\pi^-$ mass spectrum.
- These distributions are fitted with a model including the $\rho\pi$, $\rho'\pi \equiv \rho(1450)\pi$, and $\omega\pi^0$ states.
- A significant fraction of the $\rho'\pi$ intermediate state is observed in the energy region 1.55-1.75 GeV.

$e^+e^- \rightarrow \pi^+\pi^-\pi^0$ dynamics





- ✓ The isovector intermediate state $\omega\pi^0$ gives sizable (up to 20%) contribution to the $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ cross section.
- ✓ The cross section for the intermediate state $\rho'\pi$ differs significantly from zero in the range 1.55 - 1.75 GeV, where the resonance ω'' is located.
- ✓ In the $\rho\pi$ cross section the resonance structure near 1650 MeV is not seen.

We conclude that the $\rho'\pi$ intermediate state gives a significant contribution to the decay $\omega'' \rightarrow \pi^+\pi^-\pi^0$, and that the $\omega' \rightarrow \pi^+\pi^-\pi^0$ decay is dominated by the $\rho\pi$ intermediate state.

$$e^+e^- \rightarrow K^+K^-\pi^0$$

EPJ C **80** (2020) 1139





- The analysis is based on 26 pb⁻¹ data recorded in the c.m. energy range 1.27 – 2 GeV.
- The cross sections for the $K^{*\pm}K^{\mp}$ and $\phi\pi^0$ intermediate states are measured separately.
- The $e^+e^- \rightarrow K^{*\pm}K^{\mp}$ cross section is dominated by the $\phi' \equiv \phi(1680)$ resonance.

The isovector process $e^+e^- \rightarrow \phi\pi^0$ is suppressed by the Okubo-Zweig-Iizuka (OZI) rule. Three measurements of the cross section are fitted simultaneously. The fit by a sum of the ρ' and ρ'' contributions cannot describe data near 1.6 GeV. The inclusion of an unknown resonance with $m=1585\pm 15$ MeV and $\Gamma=75\pm 30$ MeV improves fit. The significance of the structure is about 3σ .

$$e^+ e^- \rightarrow \eta \pi^0 \gamma$$


2E= 1.320 - 2.000 GeV




- ✓ There is a significant contribution of the $\omega\eta$ intermediate state, which is seen as a peak in the $\pi^0\gamma$ mass distribution.
- ✓ The non- $\omega\eta$ signal is observed with significance of 5.6σ . It has a wide $\eta\pi^0$ mass distribution and may arise from the processes $e^+ e^- \rightarrow a_0(1450)\gamma$ and $a_2(1320)\gamma$.

- The process $e^+ e^- \rightarrow \eta\pi^0\gamma$ above 1.05 GeV is studied for the first time.
- Data set with $IL \approx 100 \text{ pb}^{-1}$ recorded in 2010-2012 and 2017
- The five-photon final state is used.

$$e^+e^- \rightarrow \eta\pi^0\gamma$$




The measured $e^+e^- \rightarrow \omega\eta$ cross section is in good agreement with the SND and CMD-3 measurements in the $\omega \rightarrow \pi^+\pi^-\pi^0$ decay mode.



The non-VP $e^+e^- \rightarrow \eta\pi^0\gamma$ process is observed with significance of 5.6σ .
This is the first measurement of this cross section.

$$e^+e^- \rightarrow \eta\eta\gamma$$

Cross section $e^+e^- \rightarrow \eta\eta\gamma$



- The $e^+e^- \rightarrow \eta\eta\gamma$ cross section is measured for the first time in the energy range 1.17 – 2.0 GeV.
- The main intermediate state is $\phi\eta$.
- The measured cross section is consistent with CMD-3 result on $e^+e^- \rightarrow \phi\eta$, $\phi \rightarrow K^+K^-$.
- The contribution from intermediate states other than $\phi\eta$ is not seen.

Upper limits on possible contribution of radiative intermediate states ($f_0(1500)\gamma$, $f'_2(1525)\gamma$) is set.

preliminary

2E, GeV	95% CL Upper limit, pb
1.17-1.32	9
1.32-1.57	5
1.57-1.80	11
1.80-2.00	4

Summary

- ✓ The SND detector accumulated 370 pb^{-1} of integrated luminosity in the energy range $0.3 - 2 \text{ GeV}$.
- ✓ The $e^+e^- \rightarrow \pi^+\pi^-$ cross section has been measured in the energy range $0.53-0.88 \text{ GeV}$ with a systematic uncertainty better than 1%.
- ✓ The accuracy of the $e^+e^- \rightarrow n\bar{n}$ measurement has been significantly improved.
- ✓ The dynamics of the process $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ has been studied in the energy range $1.15-2.0 \text{ GeV}$.
- ✓ The process $e^+e^- \rightarrow K^+K^-\pi^0$ has been studied in the $K^{*\pm}K^{\mp}$ and $\phi\pi^0$ intermediate states.
- ✓ Rare radiative processes $e^+e^- \rightarrow \eta\pi^0\gamma$ and $\eta\eta\gamma$ have been measured in the energy range $1.05-2 \text{ GeV}$.