



# Probing the CP structure of the top quark Yukawa coupling: Loop sensitivity vs. on-shell sensitivity

Ren-Qi Pan  
Zhejiang University  
[renqi.pan@cern.ch](mailto:renqi.pan@cern.ch)

in collaboration with Till Martini, Markus Schulze, Meng Xiao

Based on arXiv: 2104.04277

# Motivation

---

## Why CP property matters?

- ◆ Baryon asymmetry of the universe requires CP violation
- ◆ CKM matrix is the only source of CP violation in SM ,  
but can't explain the baryon asymmetry
- ◆ Additional CP violation needed to explain baryon asymmetry
- ◆ CP violation might exist in Higgs sector in some new physics models: 2HDM, SUSY

## Why Higgs-top interaction is interesting?

- ◆ Heaviest mass among fermions
- ◆ Interacts with the Higgs boson with the largest Yukawa coupling
- ◆ Lots of top quarks produced at LHC

# Htt Effective Lagrangian


$$\mathcal{L}(Htt) = -\frac{m_t}{v} \bar{\psi}_t (\kappa + i\tilde{\kappa}\gamma_5) \psi_t H$$

- ✿  $\kappa$  term: CP-even,  $\tilde{\kappa}$  term: CP-odd; In SM:  $\kappa = 1, \tilde{\kappa} = 0$
- ✿ If  $\kappa, \tilde{\kappa}$  both are non-zero, implies CP violation
- ✿  $\kappa, \tilde{\kappa}$  correspond to  $C_{33}^{u\varphi}$  in SMEFT:



$$\kappa = 1 - \frac{v}{\sqrt{2}m_t} \frac{v^2}{\Lambda^2} \operatorname{Re} [C_{tt}^{u\varphi}], \tilde{\kappa} = - \frac{v}{\sqrt{2}m_t} \frac{v^2}{\Lambda^2} \operatorname{Im} [C_{tt}^{u\varphi}]$$

- ✿ In SMEFT Warsaw basis:  $Q_{33}^{u\varphi} = (\varphi^\dagger \varphi) (\bar{q}'_{3L} t'_R \tilde{\varphi}) \Leftrightarrow C_{33}^{u\varphi}$

# Current Constraints from LHC



ttH+ $t\bar{t}q$  in diphoton: Phys. Rev. Lett. 125.061801



ttH+ $t\bar{t}q$  in 4 leptons and diphoton: arXiv:2104.12152

- ♦ Fractional cross section of CP-odd component:  $f_{CP} = \frac{|\tilde{\kappa}|^2}{|\kappa|^2 + |\tilde{\kappa}|^2} \text{ sign} \left( \frac{\tilde{\kappa}}{\kappa} \right)$
- ♦ Pure CP-odd scenario is excluded at more than  $3\sigma$
- ♦ Leaving large room of parameter space

# $t\bar{t}$ And tHW Production

**New approaches to access CP property:**




- $t\bar{t}$  production with electroweak loops
- tHW production

# Top Quark Paris Production


Theoretical predictions of  $t\bar{t}$  production reach an accuracy at a few percent level:

- QCD calculation up to NNLO+NNLL: [Czakon,Fiedler,Heymes,Mitov],[Brucherseifer,Caola,Melnikov],[Berger,Gao,Yuan,Zhu], [Czakon,Mitov,Sterman],[Beneke,Czakon,Falgari,Mitov,Schwinn],[Beneke,Falgari,Klein,Schwinn],[Kidonakis], [Ferroglia,Pecjak,Yang],[Ferroglia,Marzani,Pecjak,Yang],[Czakon,Ferroglia,Heymes,Mitov,Pecjak,Scott]
- High ambitions of the theory community towards N3LO calculations: [Duhr, Mistlberger], [Piclum, Schwinn], [Duhr, Mistlberger]

LHC measured differential  $t\bar{t}$  production cross sections and unfolded to particle level




# Top Quark Pairs Production



$t\bar{t}$  production at leading order

Calculate NLO EW corrections of  $t\bar{t}$  production with arbitrary CP mixing:

$$\mathcal{L}(Htt) = -\frac{m_t}{v} \bar{\psi}_t (\kappa + i\tilde{\kappa}\gamma_5) \psi_t H$$



Typical Feynman diagrams of NLO EW Corrections



# NLO EW Corrections

---

- ◆ Interference of diagrams with Higgs loops and Born level either  $\propto \kappa^2$  or  $\tilde{\kappa}^2$
- ◆ Loop diagrams involving the Higgs boson are IR finite but contain UV poles ; renormalization necessary within SMEFT
- ◆ One-loop amplitude is UV finite after renormalization
- ◆ EW correction factor:  $\delta_{\text{wk}} = \frac{d\sigma_{\text{wk}}^{\text{NLO}} - d\sigma^{\text{LO}}}{d\sigma^{\text{LO}}}$ ,
- ◆  $\delta_{\text{wk}}$  can be used to reweight distributions to include EW effects


# Loop sensitivity on CP Property in $t\bar{t}$ production

distributions of  $\Delta y_{t\bar{t}}$  and  $M_{t\bar{t}}$  sensitive to CP structure of top Yukawa coupling




# Loop sensitivity on CP Property in $t\bar{t}$ production

- Extract CP structure in  $(\Delta y_{t\bar{t}}, M_{t\bar{t}})$  2-D distribution
- Events simulated at MadGraph and interfaced to Pythia8
- simulate detector effects through Delphes3
- NLO EW effects included through reweighting
- semileptonic channel; Main backgrounds: single top, V+jets, QCD multijets




Distribution of reconstructed events




Relative EW corrections when  $\kappa = 1$  and  $\tilde{\kappa} = 0$

# Sensitivity of $t\bar{t}$ Production



Sensitivity of  $t\bar{t}$  production at  $300 \text{ fb}^{-1}$

# tHW Production






Two typical Feynman diagrams of the tHW

- **Htt induced diagram interferes with HWW induced diagram**
- **Destructive interference in the SM, leading to small cross section**
- **If the relative sign of Htt and HWW flips, cross section would increase obviously**
- $\sigma(\kappa, \tilde{\kappa})_{tHW} = \sigma_{SM}^{tHW} (2.82 |\kappa|^2 + 2.08 |\tilde{\kappa}|^2 - 3.87 \kappa + 2.05)$
- **HWW coupling is set same as the value in the SM**


# Matrix Element Approach

$$\mathcal{D}_{0^-}(\Omega) = \frac{\mathcal{P}_{0^-}(\Omega)}{\mathcal{P}_{0^-}(\Omega) + \mathcal{P}_{0^+}(\Omega)}, \quad \mathcal{D}_{\text{bkg}} = \frac{\mathcal{P}_{\text{sig}}(\vec{\Omega})}{\mathcal{P}_{\text{sig}}(\vec{\Omega}) + \mathcal{P}_{\text{bkg}}(\vec{\Omega})}, \quad \mathcal{D}_{\text{int}}(\Omega) = \frac{\mathcal{P}_{\text{int}}(\Omega)}{2\sqrt{\mathcal{P}_{\text{sig}}(\Omega)\mathcal{P}_{\text{alt}}(\Omega)}}$$


- $\mathcal{D}_{0^-}(\Omega)$ : distinguish CP-odd from CP-even
- $\mathcal{D}_{\text{bkg}}(\Omega)$ : separate signal from background
- $\mathcal{D}_{\text{int}}(\Omega)$ : sensitive to interference term



# Results: 2D Likelihood Scan of $\kappa$ and $\tilde{\kappa}$





$t\bar{t}$  production at  $300 \text{ fb}^{-1}$



tHW production at  $300 \text{ fb}^{-1}$

# Results: Likelihood Scan of $f_{CP}$

Fractional contribution of CP-odd component:  $f_{CP} = \frac{|\tilde{\kappa}|^2}{|\kappa|^2 + |\tilde{\kappa}|^2} \text{ sign}\left(\frac{\tilde{\kappa}}{\kappa}\right)$



ttH and tHq results cited from [arXiv: 1606.03107]

# Summary

---

- Probing CP structure of top Yukawa coupling through EW loops in  $t\bar{t}$  production
- Analyze tHW production using matrix element approach
- Compare constraints on CP structure from EW loops in  $t\bar{t}$  production with on-shell Higgs production(ttH, tHq, tHW)
- Loop sensitivity in  $t\bar{t}$  production and on-shell sensitivity in ttH and tH provide complementary handles over a wide range of parameter space
- $t\bar{t}$  and tHW production with anomalous couplings publicly available as add-on to MCFM or via JHU Generator respectively