



CEU

*Universidad  
Cardenal Herrera*

EPS-HEP 2021

Online conference, July 28, 2021

# Heavy states and electroweak effective approaches

Ignasi Rosell  
Universidad CEU Cardenal Herrera  
València (Spain)

In collaboration with:  
A. Pich (IFIC, UV-CSIC, València, Spain)  
J.J. Sanz-Cillero (UCM, Madrid, Spain)



---

[PRD 102 \(2020\) 035012 \[arXiv: 2004.02827\]](#)  
[JHEP 05 \(2019\) 092 \[arXiv: 1810.10544\]](#)  
[JHEP 04 \(2017\) 012 \[arXiv: 1609.06659\]](#)  
[PRD 93 \(2016\) 055041 \[arXiv: 1510.03114\]](#)  
[JHEP 01 \(2014\) 157 \[arXiv: 1310.3121\]](#)  
[PRL 110 \(2013\) 181801 \[arXiv: 1212.6769\]](#)

# OUTLINE



- 1) Motivation
  - 2) The effective Lagrangians
    - 1) Low energies: the non-linear Electroweak Effective Theory
    - 2) High energies: Resonance Electroweak Theory
    - 3) Matching low and high energies
  - 3) Phenomenology I: bosonic LECs
  - 4) Phenomenology II: and T at NLO
  - 5) Conclusions
- Bottom-up  
approach**

Also known as  
HEFT or EWChL



# 1. Motivation


- The **Standard Model** (SM) provides an extremely successful description of the **electroweak and strong** interactions.
- A **key feature** is the particular mechanism adopted to break the electroweak gauge symmetry to the electroweak subgroup,  $SU(2)_L \times U(1)_Y \rightarrow U(1)_{QED}$ , so that the **W** and **Z** bosons become **massive**. The **LHC** discovered a new particle around **125 GeV\***.
- Up to now all searches for **New Physics** have given negative results: **Higgs couplings** compatible with the SM and **no new states**. Therefore we can use **EFTs** because it seems there is a large **mass gap**.



\* [CMS](#) and [ATLAS](#) Collaborations.


[Diagram by C. Krause \[PhD thesis, 2016\]](#)

- Depending on the nature of the EWSB we have two possibilities for these EFTs\* (or something in between):
  - The decoupling (linear) EFT: SMEFT
    - SM-Higgs (forming a doublet with the EW Goldstones, as in the SM)
    - Weakly coupled
    - LO: SM
    - Expansion in canonical dimensions
  - The more general non-decoupling (non-linear) EFT: EWET, HEFT, EWChL
    - Non-SM Higgs (being a scalar singlet)
    - Strongly coupled
    - LO: Higgsless SM + scalar  $h$  + 3 GB (chiral Lagrangian)
    - Expansion in loops or chiral dimensions
    - Some composite Higgs models can be described within the EWET.



\* [LHCXSWG Yellow Report '16](#)

# What do we want to do?



## Similarities to Chiral Symmetry Breaking in QCD

- i) **Custodial symmetry**: The Lagrangian is approximately invariant under global  $SU(2)_L \times SU(2)_R$  transformations. **Electroweak Symmetry Breaking** (EWSB) turns to be  $SU(2)_L \times SU(2)_R \xrightarrow{\text{EWSB}} SU(2)_{L+R}$ .
- ii) Similar to the **Chiral Symmetry Breaking** (ChSB) occurring in **QCD**, i.e., similar to the “pion” Lagrangian of **Chiral Perturbation Theory** (ChPT)\*^, by replacing  $f_\pi$  by  $v=1/\sqrt{2G_F}=246$  GeV. **Rescaling** naively we expect resonances at the TeV scale.

\* [Weinberg '79](#)

\* Gasser and Leutwyler ['84](#) ['85](#)

\* Bijens et al. ['99](#) ['00](#)

\*\* [Ecker et al. '89](#)

\*\* [Cirigliano et al. '06](#)


^ [Dobado, Espriu and Herrero '91](#)

^ [Espriu and Herrero '92](#)

^ [Herrero and Ruiz-Morales '94](#)


# What do we want to do?

Estimation of the LECs




Estimation of the **Low Energy Constants (LECs)** of the **EWET** in terms of **resonance parameters**.

Short-distance constraints



**Short-distance constraints** are fundamental because we understand the **resonance Lagrangian** as an **interpolation between low- and high energies** and in order to reduce **the number of resonance parameters**.

Phenomenology



Following a typical **bottom-up** approach, what values for **resonance masses** from **phenomenology**?

## Similarities to Chiral Symmetry Breaking in QCD




Diagram by J. Santos [VIII CPAN days, 2016]

## 2. The effective Lagrangians

- ✓ Two electroweak Lagrangians for two energy regions:
  - ✓ Electroweak Effective Theory (EWET) at low energies (without resonances).
  - ✓ Resonance Electroweak Theory at high energies\* (with resonances).
- ✓ The aim of this work:

Estimation of the Low-Energy Constants (LECs) in terms of resonance parameters and phenomenological consequences: constraining the BSM heavy masses.
- ✓ Steps:
  1. Building the EWET and resonance Lagrangian
  2. Matching the two effective theories
  3. Phenomenology at low energies.

Bottom-up approach
- ✓ High-energy constraints
  1. From QCD we know the importance of sum-rules and form factors at large energies.
  2. Operators with a large number of derivatives tend to violate the asymptotic behaviour.
  3. The constraints reduce the number of unknown resonance parameters.
- ✓ This program works pretty well in QCD: estimation of the LECs (Chiral Perturbation Theory) by using Resonance Chiral Theory\*\* and importance of short-distance constraints\*\*\*.

\* Pich, IR, Santos and Sanz-Cillero '16 '17

\* Krause, Pich, IR, Santos and Sanz-Cillero '19

\*\* Cirigliano et al. '06  
\*\*\* Ecker et al. '89

# How do we build the Lagrangian?

✓ Custodial symmetry

✓ Degrees of freedom:

- ✓ At low energies: bosons  $X$  (EW goldstones, gauge bosons,  $h$ ), fermions  $\Psi$
- ✓ At high energies: previous dof + resonances (V,A,S,P and fermionic)

✓ Chiral power counting\*

$$\frac{\chi}{v} \sim \mathcal{O}(p^0) \quad \frac{\psi}{v} \sim \mathcal{O}(p) \quad \partial_\mu, m \sim \mathcal{O}(p) \quad \mathcal{T} \sim \mathcal{O}(p) \quad g, g' \sim \mathcal{O}(p)$$



SM fermions are assumed to couple weakly to the strong sector.

Explicit breaking of the custodial symmetry is assumed to be suppressed.

\* Weinberg '79

\* Appelquist and Bernand '80

\* Longhitano '80 '81

\* Manohar, and Georgi '84

\* Gasser and Leutwyler '84 '85

\* Hirn and Stern '05

\* Alonso et al. '12

\* Buchalla, Catá and Krause '13

\* Brivio et al. '13

\* Delgado et al. '14

\* Pich, IR, Santos and Sanz-Cillero '16 '17

\* Krause, Pich, IR, Santos and Sanz-Cillero '19

# How do we build the Lagrangian?

✓ Custodial symmetry

✓ Degrees of freedom:

- ✓ At low energies: bosons  $X$  (EW goldstones, gauge bosons,  $h$ ), fermions  $\Psi$
- ✓ At high energies: previous dof + resonances (V,A,S,P and fermionic)

✓ Chiral power counting\*

$$\frac{\chi}{v} \sim \mathcal{O}(p^0) \quad \frac{\psi}{v} \sim \mathcal{O}(p) \quad \partial_\mu, m \sim \mathcal{O}(p) \quad \mathcal{T} \sim \mathcal{O}(p) \quad g, g' \sim \mathcal{O}(p)$$

$$\mathcal{M}(2 \rightarrow 2) \approx \frac{p^2}{v^2} \left[ 1 + \left( \frac{c_k^r p^2}{v^2} - \frac{\Gamma_k p^2}{16\pi^2 v^2} \ln \frac{p^2}{\mu^2} + \dots \right) + \mathcal{O}(p^4) \right]$$

Finite pieces from loops  
(amplitude dependent)

Order-by-order renormalization

LO  
NLO (tree)  
suppression  
 $\sim 1/v^2 + \dots$

NLO (loop)  
typical loop suppression  
 $\sim 1/(16\pi^2 v^2)$

heavier states      non-linearity

\* Weinberg '79

\* Appelquist and Bernand '80

\* Longhitano '80 '81

\* Manohar, and Georgi '84

\* Gasser and Leutwyler '84 '85

\* Hirn and Stern '05

\* Alonso et al. '12

\* Buchalla, Catá and Krause '13

\* Brivio et al. '13

\* Delgado et al. '14

\* Pich, IR, Santos and Sanz-Cillero '16 '17

\* Krause, Pich, IR, Santos and Sanz-Cillero '19

Diagram by J.J. Sanz-Cillero [HEP 2017]

## 2.1. Low energies: the Electroweak Effective Theory (no resonances)\*

$$\begin{aligned}\mathcal{L}_{\text{EWET}}^{(2)} = & \sum_{\xi} \left( i \bar{\xi} \gamma^\mu d_\mu \xi - v \left( \bar{\xi}_L \mathcal{Y} \xi_R + \text{h.c.} \right) \right) \\ & - \frac{1}{2g^2} \langle \hat{W}_{\mu\nu} \hat{W}^{\mu\nu} \rangle_2 - \frac{1}{2g'^2} \langle \hat{B}_{\mu\nu} \hat{B}^{\mu\nu} \rangle_2 - \frac{1}{2g_s^2} \langle \hat{G}_{\mu\nu} \hat{G}^{\mu\nu} \rangle_3 \\ & + \frac{1}{2} \partial_\mu h \partial^\mu h - \frac{1}{2} m_h^2 h^2 - V(h/v) + \frac{v^2}{4} \mathcal{F}_u(h/v) \langle u_\mu u^\mu \rangle_2\end{aligned}$$

\* Longhitano '80 '81  
\* Buchalla et al. '12 '14  
\* Alonso et al. '13

\* Guo, Ruiz-Femenia and Sanz-Cillero '15  
\* Pich, IR, Santos and Sanz-Cillero '16 '17  
\* Krause, Pich, IR, Santos and Sanz-Cillero '19

## 2.1. Low energies: the Electroweak Effective Theory (no resonances)\*

Bosonic sector

$$\begin{aligned} \mathcal{L}_{\text{EWET}}^{(4)} = & \sum_{i=1}^{12} \mathcal{F}_i \mathcal{O}_i + \sum_{i=1}^3 \tilde{\mathcal{F}}_i \tilde{\mathcal{O}}_i + \sum_{i=1}^8 \mathcal{F}_i^{\psi^2} \mathcal{O}_i^{\psi^2} \\ & + \sum_{i=1}^3 \tilde{\mathcal{F}}_i^{\psi^2} \tilde{\mathcal{O}}_i^{\psi^2} + \sum_{i=1}^{10} \mathcal{F}_i^{\psi^4} \mathcal{O}_i^{\psi^4} + \sum_{i=1}^2 \tilde{\mathcal{F}}_i^{\psi^4} \tilde{\mathcal{O}}_i^{\psi^4} \end{aligned}$$

$F_1 \rightarrow$  oblique S parameter

$F_1, F_3 \rightarrow$  trilinear gauge coupling

$F_1, F_3, F_4, F_5 \rightarrow$  quartic gauge coupling

$F_6, F_7, F_8, F_9 \rightarrow$  vertexes involving H

$i$	$\mathcal{O}_i$	$\tilde{\mathcal{O}}_i$
1	$\frac{1}{4} \langle f_+^{\mu\nu} f_{+\mu\nu} - f_-^{\mu\nu} f_{-\mu\nu} \rangle_2$	$\frac{i}{2} \langle f_-^{\mu\nu} [u_\mu, u_\nu] \rangle_2$
2	$\frac{1}{2} \langle f_+^{\mu\nu} f_{+\mu\nu} + f_-^{\mu\nu} f_{-\mu\nu} \rangle_2$	$\langle f_+^{\mu\nu} f_{-\mu\nu} \rangle_2$
3	$\frac{i}{2} \langle f_+^{\mu\nu} [u_\mu, u_\nu] \rangle_2$	$\frac{(\partial_\mu h)}{v} \langle f_+^{\mu\nu} u_\nu \rangle_2$
4	$\langle u_\mu u_\nu \rangle_2 \langle u^\mu u^\nu \rangle_2$	—
5	$\langle u_\mu u^\mu \rangle_2^2$	—
6	$\frac{(\partial_\mu h)(\partial^\mu h)}{v^2} \langle u_\nu u^\nu \rangle_2$	—
7	$\frac{(\partial_\mu h)(\partial_\nu h)}{v^2} \langle u^\mu u^\nu \rangle_2$	—
8	$\frac{(\partial_\mu h)(\partial^\mu h)(\partial_\nu h)(\partial^\nu h)}{v^4}$	—
9	$\frac{(\partial_\mu h)}{v} \langle f_-^{\mu\nu} u_\nu \rangle_2$	—
10	$\langle \mathcal{T} u_\mu \rangle_2^2$	—
11	$\hat{X}_{\mu\nu} \hat{X}^{\mu\nu}$	—
12	$\langle \hat{G}_{\mu\nu} \hat{G}^{\mu\nu} \rangle_3$	—

\* Longhitano '80 '81  
 \* Buchalla et al. '12 '14  
 \* Alonso et al. '13

\* Guo, Ruiz-Femenia and Sanz-Cillero '15  
 \* Pich, IR, Santos and Sanz-Cillero '16 '17  
 \* Krause, Pich, IR, Santos and Sanz-Cillero '19

## 2.1. Low energies: the Electroweak Effective Theory (no resonances)\*

$$\begin{aligned}\mathcal{L}_{\text{EWET}}^{(4)} = & \sum_{i=1}^{12} \mathcal{F}_i \mathcal{O}_i + \sum_{i=1}^3 \tilde{\mathcal{F}}_i \tilde{\mathcal{O}}_i + \sum_{i=1}^8 \mathcal{F}_i^{\psi^2} \mathcal{O}_i^{\psi^2} \\ & + \sum_{i=1}^3 \tilde{\mathcal{F}}_i^{\psi^2} \tilde{\mathcal{O}}_i^{\psi^2} + \sum_{i=1}^{10} \mathcal{F}_i^{\psi^4} \mathcal{O}_i^{\psi^4} + \sum_{i=1}^2 \tilde{\mathcal{F}}_i^{\psi^4} \tilde{\mathcal{O}}_i^{\psi^4}\end{aligned}$$

## 2.2. High energies: Resonance Electroweak Theory (with resonances)\*\*

$$\mathcal{L}_{\text{RT}} = \mathcal{L}_{\text{R}}[R, \chi, \psi] + \mathcal{L}_{\text{non-R}}[\chi, \psi]$$

- Bosonic resonances:
  - V, A, S and P
  - SU(2) singlets and triplets
  - SU(3) singlets and octets
  - Spin-1 resonances with Proca or antisymmetric formalism
- Fermionic doublet resonances:
  - Including operators with one heavy fermionic resonance

Number of operators				
Field ( $\mathbf{R}^{\text{QCD}}_{\text{EW}}$ )	$\mathbf{R}^1_1$	$\mathbf{R}^1_3$	$\mathbf{R}^8_1$	$\mathbf{R}^8_3$
S	3	1	1	1
P	1	2	1	1
V with Proc	3	2	2	2
A with Proc	3	2	2	2
V with ant.	2	5	2	1
A with ant.	2	5	2	1
Fermionic				6

\*\* Pich, IR, Santos and Sanz-Cillero '16 '17

\*\* Krause, Pich, IR, Santos and Sanz-Cillero '19

## 2.1. Low energies: the Electroweak Effective Theory (no resonances)\*

$$\begin{aligned}\mathcal{L}_{\text{EWET}}^{(4)} = & \sum_{i=1}^{12} \mathcal{F}_i \mathcal{O}_i + \sum_{i=1}^3 \tilde{\mathcal{F}}_i \tilde{\mathcal{O}}_i + \sum_{i=1}^8 \mathcal{F}_i^{\psi^2} \mathcal{O}_i^{\psi^2} \\ & + \sum_{i=1}^3 \tilde{\mathcal{F}}_i^{\psi^2} \tilde{\mathcal{O}}_i^{\psi^2} + \sum_{i=1}^{10} \mathcal{F}_i^{\psi^4} \mathcal{O}_i^{\psi^4} + \sum_{i=1}^2 \tilde{\mathcal{F}}_i^{\psi^4} \tilde{\mathcal{O}}_i^{\psi^4}\end{aligned}$$

## 2.2. High energies: Resonance Electroweak Theory (with resonances)\*\*

$$\mathcal{L}_{\text{RT}} = \mathcal{L}_{\text{R}}[R, \chi, \psi] + \mathcal{L}_{\text{non-R}}[\chi, \psi]$$

- Bosonic resonances:
  - V, A, S and P
  - SU(2) singlets and triplets
  - SU(3) singlets and octets
  - Spin-1 resonances with Proca or antisymmetric formalism
- Fermionic doublet resonances:
  - Including operators with one heavy fermionic resonance

## 2.3. Matching low and high energies

$$e^{iS_{\text{eff}}[\chi, \psi]} = \int [dR] e^{iS[\chi, \psi, R]}$$

- ✓ Integration of the heavy modes
- ✓ Similar to the ChPT case\*\*\*
- ✓ EWET LECs in terms of resonance parameters\*\*
- ✓ Tracks of resonances in the EWET.

\*\* Pich, IR, Santos and Sanz-Cillero '16 '17

\*\* Krause, Pich, IR, Santos and Sanz-Cillero '19

\*\*\* Ecker et al. '89

### 3. Phenomenology I: bosonic LECs\*

- ✓ Integration of the heavy modes
- ✓ The case of P-even **bosonic operators**<sup>\*\*</sup>:
- ✓ Short-distance constraints
- ✓ Experimental constraints [95% CL]:

$i$	$\mathcal{O}_i$	$\mathcal{F}_i$
1	$\frac{1}{4} \langle f_+^{\mu\nu} f_{+\mu\nu} - f_-^{\mu\nu} f_{-\mu\nu} \rangle_2$	$-\frac{F_V^2 - \tilde{F}_V^2}{4M_{V_3^1}^2} + \frac{F_A^2 - \tilde{F}_A^2}{4M_{A_3^1}^2}$
3	$\frac{i}{2} \langle f_+^{\mu\nu} [u_\mu, u_\nu] \rangle_2$	$-\frac{F_V G_V}{2M_{V_3^1}^2} - \frac{\tilde{F}_A \tilde{G}_A}{2M_{A_3^1}^2}$
4	$\langle u_\mu u_\nu \rangle_2 \langle u^\mu u^\nu \rangle_2$	$\frac{G_V^2}{4M_{V_3^1}^2} + \frac{\tilde{G}_A^2}{4M_{A_3^1}^2}$
5	$\langle u_\mu u^\mu \rangle_2 \langle u_\nu u^\nu \rangle_2$	$\frac{c_d^2}{4M_{S_1^1}^2} - \frac{G_V^2}{4M_{V_3^1}^2} - \frac{\tilde{G}_A^2}{4M_{A_3^1}^2}$
6	$\frac{(\partial_\mu h)(\partial^\mu h)}{v^2} \langle u_\nu u^\nu \rangle_2$	$-\frac{\tilde{\lambda}_1^{hV} v^2}{M_{V_3^1}^2} - \frac{\lambda_1^{hA} v^2}{M_{A_3^1}^2}$
7	$\frac{(\partial_\mu h)(\partial_\nu h)}{v^2} \langle u^\mu u^\nu \rangle_2$	$\frac{d_P^2}{2M_{P_3^1}^2} + \frac{\lambda_1^{hA} v^2}{M_{A_3^1}^2} + \frac{\tilde{\lambda}_1^{hV} v^2}{M_{V_3^1}^2}$
8	$\frac{(\partial_\mu h)(\partial^\mu h)(\partial_\nu h)(\partial^\nu h)}{v^4}$	0
9	$\frac{(\partial_\mu h)}{v} \langle f_-^{\mu\nu} u_\nu \rangle_2$	$-\frac{F_A \lambda_1^{hA} v}{M_{A_3^1}^2} - \frac{\tilde{F}_V \tilde{\lambda}_1^{hV} v}{M_{V_3^1}^2}$

LEC	Data
$0.89 < \kappa_W < 1.13$	LHC[1]
$-1.02 < c_{2V} < 2.71$	LHC[2]
$-0.004 < \mathcal{F}_1 < 0.004$	LEP via S[3]
$-0.06 < \mathcal{F}_3 < 0.20$	LEP & LHC[4]
$-0.0006 < \mathcal{F}_4 < 0.0006$	LHC[5]
$-0.0010 < \mathcal{F}_4 + \mathcal{F}_5 < 0.0010$	LHC[5]

From one-loop considerations one would expect  $F_i \approx 1/(4\pi^2) \approx 10^{-3}$ .

The running is known\*\*\*:  
 $|F_i(\mu = M_R) - F_i(\mu = m_h)| \approx 10^{-3}$

[1] [Blas, Eberhardt and Krause '18](#)

[2] [ATLAS-CONF-2019-030](#)

[3] [PDG '18](#)

[4] [Da Silva et al. '19](#)

[5] [CMS '19](#)

\*\* [Pich, IR, Santos and Sanz-Cillero '17](#)

\*\* [Krause, Pich, IR, Santos and Sanz-Cillero '19](#)

\*\*\* [Guo, Ruiz-Femenía and Sanz-Cillero '15](#)

\* [Pich, IR, Santos and Sanz-Cillero '16](#)

\* [Pich, IR and Sanz-Cillero '20](#)

### 3. Phenomenology I: bosonic LECs\*

- ✓ Integration of the heavy modes
- ✓ The case of P-even **bosonic operators**<sup>\*\*</sup>:
- ✓ Short-distance constraints
- ✓ Experimental constraints [95% CL]:

$i$	$\mathcal{O}_i$	$\mathcal{F}_i$
1	$\frac{1}{4} \langle f_+^{\mu\nu} f_{+\mu\nu} - f_-^{\mu\nu} f_{-\mu\nu} \rangle_2$	$-\frac{F_V^2 - \tilde{F}_V^2}{4M_{V_3^1}^2} + \frac{F_A^2 - \tilde{F}_A^2}{4M_{A_3^1}^2}$
3	$\frac{i}{2} \langle f_+^{\mu\nu} [u_\mu, u_\nu] \rangle_2$	$-\frac{F_V G_V}{2M_{V_3^1}^2} - \frac{\tilde{F}_A \tilde{G}_A}{2M_{A_3^1}^2}$
4	$\langle u_\mu u_\nu \rangle_2 \langle u^\mu u^\nu \rangle_2$	$\frac{G_V^2}{4M_{V_3^1}^2} + \frac{\tilde{G}_A^2}{4M_{A_3^1}^2}$
5	$\langle u_\mu u^\mu \rangle_2 \langle u_\nu u^\nu \rangle_2$	$\frac{c_d^2}{4M_{S_1^1}^2} - \frac{G_V^2}{4M_{V_3^1}^2} - \frac{\tilde{G}_A^2}{4M_{A_3^1}^2}$
6	$\frac{(\partial_\mu h)(\partial^\mu h)}{v^2} \langle u_\nu u^\nu \rangle_2$	$-\frac{\tilde{\lambda}_1^{hV} v^2}{M_{V_3^1}^2} - \frac{\lambda_1^{hA} v^2}{M_{A_3^1}^2}$
7	$\frac{(\partial_\mu h)(\partial_\nu h)}{v^2} \langle u^\mu u^\nu \rangle_2$	$\frac{d_P^2}{2M_{P_3^1}^2} + \frac{\lambda_1^{hA} v^2}{M_{A_3^1}^2} + \frac{\tilde{\lambda}_1^{hV} v^2}{M_{V_3^1}^2}$
8	$\frac{(\partial_\mu h)(\partial^\mu h)(\partial_\nu h)(\partial^\nu h)}{v^4}$	0
9	$\frac{(\partial_\mu h)}{v} \langle f_-^{\mu\nu} u_\nu \rangle_2$	$-\frac{F_A \lambda_1^{hA} v}{M_{A_3^1}^2} - \frac{\tilde{F}_V \tilde{\lambda}_1^{hV} v}{M_{V_3^1}^2}$

LEC	coupling
$0.89 < \kappa_W < 1.13$	hWW coupling
$-1.02 < c_{2V} < 2.71$	hhWW coupling
$-0.004 < \mathcal{F}_1 < 0.004$	S parameter
$-0.06 < \mathcal{F}_3 < 0.20$	triple gauge coupling
$-0.0006 < \mathcal{F}_4 < 0.0006$	quartic gauge coupling
$-0.0010 < \mathcal{F}_4 + \mathcal{F}_5 < 0.0010$	

From one-loop considerations one would expect  $F_i \approx 1/(4\pi^2) \approx 10^{-3}$ .





The running is known\*\*\*:  
 $|F_i(\mu = M_R) - F_i(\mu = m_h)| \approx 10^{-3}$

- [1] [Blas, Eberhardt and Krause '18](#)
- [2] [ATLAS-CONF-2019-030](#)
- [3] [PDG '18](#)
- [4] [Da Silva et al. '19](#)
- [5] [CMS '19](#)

\*\* [Pich, IR, Santos and Sanz-Cillero '17](#)  
 \*\* [Krause, Pich, IR, Santos and Sanz-Cillero '19](#)  
 \*\*\* [Guo, Ruiz-Femenía and Sanz-Cillero '15](#)



\* [Pich, IR, Santos and Sanz-Cillero '16](#)

\* [Pich, IR and Sanz-Cillero '20](#)





$$\begin{aligned} M_A &= M_V & M_A &= 1.1M_V \\ M_A &= 1.2M_V & M_A &\rightarrow \infty \end{aligned}$$

\* [Pich, IR, Santos and Sanz-Cillero '16](#)  
 \* [Pich, IR and Sanz-Cillero '20](#)





$$\begin{aligned} M_A &= M_V \\ M_A &= 1.1M_V \\ M_A &\rightarrow \infty \end{aligned}$$



\* [Pich, IR, Santos and Sanz-Cillero '16](#)  
\* [Pich, IR and Sanz-Cillero '20](#)

## 4. Phenomenology II: S and T at NLO\*

- ✓ Oblique electroweak observables\*\* (S and T)
- ✓ Dispersive relations for both S\*\* and T\*
- ✓ Short-distance constraints



- ✓ Room for these scenarios ( $M_R \approx \text{TeV}$ )
- ✓ Similar preliminary results including BSM fermionic corrections.

\* Pich, IR and Sanz-Cillero '12 '13 '14 '21

\*\* Peskin and Takeuchi '92




## 4. Conclusions

- ✓ Up to now all searches for **New Physics** have given negative results: **Higgs couplings compatible with the SM and no new states**. Therefore we can use **EFTs** because we have a **mass gap**.
- ✓ As a consequence of the **mass gap**, **bottom-up EFTs** are appropriate to search for BSM. Depending on the nature of the EWSB we have two possibilities:
  - ✓ Decoupling (linear) EFT: **SMEFT**
    - ✓ **SM-Higgs** and **weakly coupled**
    - ✓ Expansion in **canonical dimensions**
  - ✓ Non-decoupling (non-linear) EFT: **EWET (HEFT or EWChL)**
    - ✓ **Non-SM Higgs** and **strongly coupled**
    - ✓ Expansion in **loops or chiral dimensions**
- ✓ Similarities to **ChSB of QCD** -> **ChPT** and **RChT**
- ✓ **Phenomenology**
  - ✓ Estimation of the LECs by using resonance Lagrangians and short-distance constraints.
  - ✓ S and T at NLO by using resonance Lagrangians and short-distance constraints.

**Experimental LHC constraints  
start to be competitive.**


**Room for these BSM scenarios  
and  $M_R \gtrsim 2$  TeV.**

# Phenomenology I: bosonic LECs\* [annex]



\* [Pich, IR, Santos and Sanz-Cillero '16](#)  
 \* [Pich, IR and Sanz-Cillero '20](#)

## Phenomenology II: S and T at NLO\* [annex]



\* Pich, IR and Sanz-Cillero '12 '13 '14

## Phenomenology III: contact four-fermion operators\*

- ✓ With light leptons and/or quarks

- ✓ From dijet production

$\Lambda \geq 21.8 \text{ TeV}$  from ATLAS

$\Lambda \geq 18.6 \text{ TeV}$  from CMS

$\Lambda \geq 16.2 \text{ TeV}$  from LEP

- ✓ From dilepton production

$\Lambda \geq 26.3 \text{ TeV}$  from ATLAS

$\Lambda \geq 19.0 \text{ TeV}$  from CMS

$\Lambda \geq 24.6 \text{ TeV}$  from LEP

- ✓ Including top and bottom quarks

- ✓ From high-energy collider studies

$\Lambda \geq 1.5 \text{ TeV}$  from multi-top production at LHC and Tevatron

$\Lambda \geq 2.3 \text{ TeV}$  from  $t$  and  $t\bar{t}$  production at LHC and Tevatron

$\Lambda \geq 4.7 \text{ TeV}$  from dilepton production at LHC

- ✓ From low-energy studies

$\Lambda \geq 14.5 \text{ TeV}$  from  $B_s - \bar{B}_s$  mixing



$\Lambda \geq 3.3 \text{ TeV}$  from semileptonic B decays

GeV

\* See references in [Krause, Pich, IR, Santos and Sanz-Cillero '19](#)

## Phenomenology IV: HVT diboson searches\*

- ✓ Our model-independent approach can be related to the popular Heavy Vector Triplet simplified model (HVT)\*\*.
- ✓ LHC diboson production experimental analysis (ATLAS and CMS).
- ✓ Exclusion in the (mass, coupling) plane and the scale  $\Lambda$



Integration of heavy modes

$$\frac{2\pi}{\Lambda^2} \equiv \mathcal{F}_7^{\psi^4} + \mathcal{F}_8^{\psi^4} + \frac{\mathcal{F}_{10}^{\psi^4}}{4} = \frac{c_{\text{eff}}^2}{4M_V^2}$$

EWET

Resonance Lagrangian

The diagram illustrates the decomposition of the loop correction  $\frac{2\pi}{\Lambda^2}$  into two parts: EWET (Electroweak Effective Theory) and Resonance Lagrangian. The EWET part is shown as a red bracket spanning from  $\Lambda \approx 400$  to  $800$  TeV, while the Resonance Lagrangian part is shown as a blue bracket spanning from  $\Lambda \approx 400$  to  $600$  TeV.

\* Krause, Pich, IR, Santos and Sanz-Cillero '19

\*\* Pappadopulo et al. '14

## Proca vs. antisymmetric formalism\*

- ✓ By using path integral and changes of variables both formalisms are proven to be equivalent:
  - ✓ A set of relations between resonance parameters emerges.
  - ✓ The couplings of the non-resonant operators are different:  $\mathcal{L}_{\text{non-R}}^{(P)} \neq \mathcal{L}_{\text{non-R}}^{(A)}$
- ✓ High-energy behaviour is fundamental:

$$\mathbb{F}_{\varphi\varphi}^V(s) = \begin{cases} 1 + \frac{F_V G_V}{v^2} \frac{s}{M_V^2 - s} + \frac{\tilde{F}_A \tilde{G}_A}{v^2} \frac{s}{M_A^2 - s} - 2 \mathcal{F}_3^{\text{SDA}} \frac{s}{v^2} & (\text{A}) \\ 1 + \frac{f_{\hat{V}} g_{\hat{V}}}{v^2} \frac{s^2}{M_V^2 - s} + \frac{\tilde{f}_{\hat{A}} \tilde{g}_{\hat{A}}}{v^2} \frac{s^2}{M_A^2 - s} - 2 \mathcal{F}_3^{\text{SDP}} \frac{s}{v^2} & (\text{P}) \end{cases}$$



\* Ecker et al. '89

\* Bijens and Pallante '96

\* Kampf, Novotny and Trnka '07

\* Pich, IR, Santos and Sanz-Cillero '16 '17

\* Krause, Pich, IR, Santos and Sanz-Cillero '19

$$\mathcal{F}_3^{\text{SDA}} = 0$$

$$\mathcal{F}_3^{\text{SDP}} = -\frac{f_{\hat{V}} g_{\hat{V}}}{2} - \frac{\tilde{f}_{\hat{A}} \tilde{g}_{\hat{A}}}{2}$$