



Contribution ID: 48

Type: not specified

## Renormalization of the radiative jet function and its zero-bin subtraction

Friday, 19 November 2021 15:40 (20 minutes)

We show how to compute directly the renormalization/evolution of the radiative jet function that appears in the factorization theorems for  $B \rightarrow \gamma \ell \nu$  and  $H \rightarrow \gamma \gamma$  through a  $b$ -quark loop. We point out that, in order to avoid double counting of soft contributions, one should use in the factorization theorems a subtracted radiative jet function, from which soft contributions have been removed. The soft-contribution subtractions are zero-bin subtractions in the terminology of soft-collinear effective theory. We show that they can be factored from the radiative jet function and that the resulting soft-subtraction function gives rise to a nonlocal renormalization of the subtracted radiative jet function. This is a novel instance in which zero-bin subtractions lead to a nonlocality in the renormalization of a subtracted quantity that is not present in the renormalization of the unsubtracted quantity. We demonstrate the use of our formalism by computing the order- $\alpha_s$  evolution kernel for the subtracted radiative jet function.

**Primary authors:** EE, June-Haak (Fudan university); Dr BODWIN, Geoffrey T. (High Energy Physics Division, Argonne National Laboratory); Prof. LEE, Jungil (Department of Physics, Korea University); Dr WANG, Xiang-Peng (High Energy Physics Division, Argonne National Laboratory)

**Presenter:** EE, June-Haak (Fudan university)

**Session Classification:** Parallel session