

CP-violation in SUSY cascades at the LHC

Jamie Tattersall

Bethe Centre for Theoretical Physics,
Universität Bonn

In collaboration with P. Bechtle, B. Godzik, G.
Moortgat-Pick, K. Rolbiecki, P. Wienemann.
(Previous work: J. Ellis, F. Moortgat, J. Smillie)

Physics at the Terascale, Dresden



Bethe Center for
Theoretical Physics

In the Standard Model, the only source of CP violation comes from the complex phase within the CKM matrix.

- The phase of the CKM in the Standard Model contains too little CP violation for Baryogenesis.

(Phys. Rept. 401, 1 (2005): Chung, Everett, Kane, King, Lykken and Wang)

- Consequently, we require new CP violating terms to explain the asymmetry we see in the universe.

MSSM (Minimal Supersymmetric Standard Model) contains several complex parameters that can all contribute.

In the Standard Model, the only source of CP violation comes from the complex phase within the CKM matrix.

- The phase of the CKM in the Standard Model contains too little CP violation for Baryogenesis.

(Phys. Rept. 401, 1 (2005): Chung, Everett, Kane, King, Lykken and Wang)

- Consequently, we require new CP violating terms to explain the asymmetry we see in the universe.

MSSM (Minimal Supersymmetric Standard Model) contains several complex parameters that can all contribute.

We explore methods for determining the CP violating effects in the electroweak part of the MSSM at the LHC.

- Most detailed phenomenological analyses have been based on a future LC.
- Precise determination of phases only expected at a LC.
- Crucial for future search strategy to use LHC data to learn as much as possible.
- Choose processes with the most promising discovery potential at LHC (coloured states).

Production	Decay	Paper
$\tilde{t}\tilde{t}^*$	$\tilde{t} \rightarrow \tilde{\chi}_2^0 t, \tilde{\chi}_2^0 \rightarrow \tilde{\ell}\ell$	<p>A. Bartl, E. Christova, K. Hohenwarter-Sodek, T. Kernreiter, hep-ph/0409060</p> <p>F. Deppisch and O. Kittel, 0905.3088</p> <p>G. Moortgat-Pick, K. Rolbiecki, J. Tattersall, 1008.2206</p>
$\tilde{t}\tilde{t}^*$	$\tilde{t} \rightarrow \tilde{\chi}_2^0 t, \tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 \ell\ell$	<p>P. Langacker, G. Paz, L.-T. Wang, I. Yavin, hep-ph/0702068</p> <p>J. Ellis, F. Moortgat, G. Moortgat-Pick, J. M. Smillie J. Tattersall, 0905.3088</p>
$\tilde{b}\tilde{b}^*$	$\tilde{b} \rightarrow \tilde{\chi}^+ t, \tilde{\chi}^+ \rightarrow \tilde{\nu}_\ell \ell$	<p>A. Bartl, E. Christova, K. Hohenwarter-Sodek, T. Kernreiter, hep-ph/0610234</p> <p>F. Deppisch and O. Kittel, 1003.5186</p>
$\tilde{q}\tilde{q}$	$\tilde{q} \rightarrow \tilde{\chi}_2^0 q, \tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 \ell\ell$	<p>G. Moortgat-Pick, K. Rolbiecki, J. Tattersall, P. Wienemann, 0908.2631</p>

We consider the MSSM with parameters defined at the weak scale.

- In this framework the gaugino and higgsino mass parameters and the trilinear couplings can have complex phases.

$$M_i = |M_i|e^{i\phi_i}, \quad \mu = |\mu|e^{i\phi_\mu}, \quad A_f = |A_f|e^{i\phi_f}$$

- CP-even observables
 - Affect sparticle masses and couplings.
 - Influence cross sections and branching ratios.
- Generate CP odd observables (unique determination of CP phases) that can in principle be large as they are already present at tree level.

Certain combinations of the CP violating phases are constrained by experimental upper bounds on various EDMs (Electric Dipole Moments).

- ϕ_μ is the most severely constrained.
 - Contributes at the one loop level to EDMs.
- ϕ_{M_1} also contributes at the one loop level to EDMs.
 - Accidental cancellations may allow it to become less constrained.
- The phases of the third-generation trilinear couplings, $\phi_{A_{t,b,\tau}}$ have weaker constraints.
 - Only contribute to EDMs at the two-loop level.

(arXiv:0710.5117, Kraml) ref therein.

Triple Product Correlations of momenta are a useful tool for studying CP-violating effects.

- Construct a T_N -odd observable:

$$\mathcal{T} = \vec{p}_1 \cdot (\vec{p}_2 \times \vec{p}_3)$$

- It is CP-odd (by CPT_N) if higher order effects and finite widths can be neglected.
- **Originates from Dirac traces in matrix element:**

$$\text{tr}(\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma \gamma^5) \longrightarrow i \epsilon_{\mu\nu\rho\sigma} p_a^\mu p_b^\nu p_c^\rho p_d^\sigma.$$

- Together with imaginary part of couplings gives rise to CP-odd asymmetries.
- The covariant product can be expanded in terms of explicit 4-momentum components:

$$E_a \vec{p}_b \cdot (\vec{p}_c \times \vec{p}_d) \pm \dots\dots\dots$$

- mSUGRA parameters:

Parameter	m_0	$m_{1/2}$	$\tan \beta$	$\text{sign}(\mu)$	A_0
Value	65	210	5	+	0

- Masses in GeV and branching ratios:

$m_{\tilde{\chi}_1^0}$	$m_{\tilde{\chi}_2^0}$	$m_{\tilde{t}_1}$	$m_{\tilde{\ell}_L}$	$m_{\tilde{\ell}_R}$
77.7	142.4	347.7	163.4	110.8


$\tilde{t}_1 \rightarrow \tilde{\chi}_2^0 t$	$\tilde{t}_1 \rightarrow \tilde{\chi}_1^0 t$	$\tilde{t}_1 \rightarrow \tilde{\chi}_2^+ b$	$\tilde{t}_1 \rightarrow \tilde{\chi}_1^+ b$	$\tilde{\chi}_2^0 \rightarrow \tilde{\ell}_R \ell$
7.5%	34.5%	8%	50%	23%

- Stop production cross section $\sigma(pp \rightarrow \tilde{t}_1 \tilde{t}_1^*) = 3.4\text{pb}$.
- Introduce the complex phase for the stop trilinear coupling:

$$A_t = |A_t| e^{i\phi_t}, \quad 0 \leq \phi_t \leq 2\pi.$$

Process studied:

$$\begin{aligned}
 p p &\implies \tilde{t} \tilde{t}^*, \\
 \tilde{t} &\implies t \tilde{\chi}_2^0, \\
 \tilde{\chi}_2^0 &\implies \tilde{l}^- l_N^+, \\
 \tilde{l}^- &\implies \tilde{\chi}_1^0 l_F^-,
 \end{aligned}$$




- All SUSY decays are 2-body, on-shell.
- Top required to decay hadronically for reconstruction.
- Opposite \tilde{t}_1^* decay required to have single charged lepton in final state.

Realising CP asymmetry

I choose an example triple product:

$$\mathcal{T}_t = \vec{p}_{\ell_N} \cdot (\vec{p}_t \times \vec{p}_b)$$

Two-body decay forces t , b and W to define a plane.




- A non-zero expectation value of \mathcal{T} , implies a non-zero average angle between the plane and p_{ℓ} .
- Define asymmetry parameter:

$$\eta = \frac{N_+ - N_-}{N_+ + N_-} = \frac{N_+ - N_-}{N_{total}}$$

where:


$$N_+ = \int_0^1 \frac{d\Gamma}{d\cos\theta} d\cos\theta, \quad N_- = \int_{-1}^0 \frac{d\Gamma}{d\cos\theta} d\cos\theta,$$

Partonic Level Asymmetry




Blue: $\mathcal{T}_t = \vec{p}_{\ell_N} \cdot (\vec{p}_t \times \vec{p}_b)$
Purple: $\mathcal{T}_b = \vec{p}_b \cdot (\vec{p}_{\ell_N} \times \vec{p}_{\ell_F})$

- Asymmetry $\approx 15\%$ maximum at the parton level in the \tilde{t}_1 rest frame.
- Maximum asymmetry for triple product $\mathcal{T}_t = \vec{p}_{\ell_N} \cdot (\vec{p}_t \times \vec{p}_b)$.
 - $\mathcal{T}_b = \vec{p}_b \cdot (\vec{p}_{\ell_N} \times \vec{p}_{\ell_F})$ not a 'true' triple product.
- T-odd observable.



Blue: $\mathcal{T}_t = \vec{p}_{\ell_N} \cdot (\vec{p}_t \times \vec{p}_b)$
 Purple: $\mathcal{T}_b = \vec{p}_b \cdot (\vec{p}_{\ell_N} \times \vec{p}_{\ell_F})$

- \tilde{t}_1 are boosted due to production process and PDFs.
- Asymmetry is maximal in rest frame of decaying particle.
 - $\epsilon_{\mu\nu\rho\sigma} p_{\tilde{t}_1}^\mu p_{\ell_N}^\nu p_t^\rho p_b^\sigma \longrightarrow m_{\tilde{t}_1} \vec{p}_{\ell_N} \cdot (\vec{p}_t \times \vec{p}_b)$.
- Dilution of asymmetry due to ℓ_N flipping orientation in comparison to plane defined by tb .



Blue: $\mathcal{T}_t = \vec{p}_{\ell_N} \cdot (\vec{p}_t \times \vec{p}_b)$
Purple: $\mathcal{T}_b = \vec{p}_b \cdot (\vec{p}_{\ell_N} \times \vec{p}_{\ell_F})$

- After including production process and folding in PDF's, asymmetry drops to $\approx 4\%$ maximum.
- Far less dilution seen for triple product $\mathcal{T}_b = \vec{p}_b \cdot (\vec{p}_{\ell_N} \times \vec{p}_{\ell_F})$.
- All results generated analytically, cross-checked with Herwig++.

Main problem with measuring asymmetries at the LHC is the dilution in the lab frame.

- In the rest frame of the decaying particle the asymmetry is maximal.
- Reconstruct LSP momentum using the set of invariant equations:

$$\begin{aligned}m_{\tilde{\chi}_1^0}^2 &= (P_{\tilde{\chi}_1^0})^2, \\m_{\tilde{\ell}^\pm}^2 &= (P_{\tilde{\chi}_1^0} + P_{\ell_F^\pm})^2, \\m_{\tilde{\chi}_2^0}^2 &= (P_{\tilde{\chi}_1^0} + P_{\ell_F^\pm} + P_{\ell_N^\mp})^2, \\m_{\tilde{t}_1}^2 &= (P_{\tilde{\chi}_1^0} + P_{\ell_F^\pm} + P_{\ell_N^\mp} + P_t)^2.\end{aligned}$$

- We reconstruct the frame of the decaying particle on an event by event basis and the asymmetry is restored.

(Phys. Rev. D71 (2005) 035008: K. Kawagoe, M. M. Nojiri, and G. Polesello)


Solution Procedure:

- Assume particle masses are previously measured.
 - Mass errors of up to 20 GeV tested.
- Solve for three linear equations first.
- Then solve final quadratic equation ($P_{\tilde{\chi}_1^0}$) \rightarrow All events have at least 2 solutions.


Experimental procedure:

- Try to reconstruct event with all combinatorial particle assignments.
- Only use real physical solutions.
- Effective discrimination between near and far leptons.
- Boost to rest frame for all solutions but only select events with the same sign triple product.

Lab Frame




\tilde{t}_1 Rest Frame



- Using events generated by `Herwig++`, improvement in rest frame can clearly be seen.
- Angle between tb plane and ℓ_N is enhanced.
 - Asymmetry becomes more resolvable.

Final state: $pp \rightarrow \tilde{t}_1 \tilde{t}_1^* \rightarrow l^+ l^- bjj + l^\pm b + E_{miss} + jets$.



Final state: $pp \rightarrow \tilde{t}_1 \tilde{t}_1^* \rightarrow \ell^+ \ell^- bjj + \ell^\pm b + E_{miss} + jets$.

We perform the study at the hadronic level using Rivet.


- Tri-lepton final state required (1 OSSF).
- At least one b-jet, 60% efficiency.
- At least two additional jets.
- A top candidate with mass $150 < M_{jjb} < 190\text{GeV}$ and a W with mass $70 < M_{jj} < 90\text{GeV}$.
- Anti- k_t jet algorithm used ($R=0.5$).
- Selection cuts:

$$\begin{aligned} p_{Tj_i} &> 20\text{GeV}, & p_{T\ell_i} &> 10\text{GeV}, \\ |\eta| &< 2.5, & M_{\ell^+\ell^-} &> 10\text{GeV}. \end{aligned}$$


- Lepton jet isolation, $\Delta R = 0.5$.

Following standard model backgrounds included:

- **Generated by Herwig++:**
 - $t\bar{t}$, Drell-Yan (Z, γ, W), W +jet, Z +jet, WW , WZ , ZZ , $W\gamma$, WZ .
- **Generated by Madgraph:**
 - $t\bar{t}l^+l^-$,
- Only $t\bar{t}l^+l^-$ events seen $\lesssim 1\%$ of signal.
- No jet mis-tagging included.
- $t\bar{t}$ background could still be an issue \rightarrow Needs full experimental study.





- After momentum reconstruction, asymmetry $\approx 15\%$ maximum recovered.
- Slight dilution ($\approx 1.5\%$) due to hadronic smearing of rest frame.
- Slight enhancement ($\approx 1.5\%$) due to removal of small triple products.




- Expected luminosity reach at the LHC.
- Includes cross sections and branching ratios that both alter with phase ϕ_{A_t} .

Luminosity Reach





- Low mass scenario requires less luminosity since production cross section higher.
- Maximum asymmetry is reduced as A_0 increases ($|A_t|$ is smaller).
- An increase in $\tan\beta$ reduces $\tilde{\chi}_2^0 \rightarrow \tilde{\ell}\ell$ branching ratio.

- *Delphes* is a new tool that allows quick detector simulation.





(arXiv:0903.2225: S. Oryn, X. Rouby, V. Lemaitre)


- We would like to complete a detector simulation to verify observables.
- First need to confirm Delphes is a good description of ATLAS detector.
- Validation and tuning has been performed by Björn Gosdzik.
- Results expected soon.



- η and ϕ reconstruction was initially too optimistic.
- Retuned to match performance of ATLAS.
- Plots courtesy of Björn Godzik.



- p_T reconstruction shows good agreement with ATLAS.
- Monte-carlo Z-production from Herwig++ matches expected ATLAS distribution.
- Plots courtesy of Björn Godzik.



- Electron efficiency was initially a step function.
- Efficiency for various ATLAS selection algorithms is now complete.
- Plots courtesy of Björn Godzik.

- New forms of CP violation are required to explain the baryon asymmetry we see in the universe.
- **MSSM can contain new phases that lead to CP violation.**
- SUSY decays of \tilde{t}_1 can produce large CP-odd asymmetries.
- Diluted by PDFs and production process.
- **Momentum reconstruction offers a way to observe these at the LHC.**
- Delphes validation is complete and an experimental simulation is expected soon.

Backup Slides

Final state:

- $\ell^+ \ell^- bjj + \ell^\pm b + E_{miss} + jets$


Most significant background are other SUSY channels.

- $pp \rightarrow \tilde{g}\tilde{g}/\tilde{g}\tilde{q}$ followed by $\tilde{g} \rightarrow \tilde{b}_i$ has far higher rate.
- **Momentum reconstruction significantly reduces background.**
- Extra cuts needed \rightarrow SUSY background has higher average p_T and more jet activity.

Cuts used:


$$\begin{aligned} p_T(\text{Hardest Jet}) &< 200 \text{ GeV}, \\ p_T(\text{2nd Jet}) &< 130 \text{ GeV}, \\ p_T(\text{3rd Jet}) &< 80 \text{ GeV (if applicable)}, \\ p_T(\text{Any } b \text{ Jet}) &< 150 \text{ GeV}, \\ p_T(\text{Any Lepton}) &< 100 \text{ GeV}, \\ \text{Number of jets} &< 7. \end{aligned}$$

	$\tilde{t}_1 \tilde{t}_1^*$	SUSY	Signal/Background
Cross Section (pb ⁻¹)	3.44	80.1	
Events with 500 fb ⁻¹	1.7×10^6	4×10^7	0.043
Events with 500 fb ⁻¹ Initial selection	32389	410735	0.079
Events with 500 fb ⁻¹ Top Reconstruction	7117	64729	0.11
Events with 500 fb ⁻¹ Kinematic Reconstruction	1213	3759	0.32
Events with 500 fb ⁻¹ Extra SUSY cuts	901	967	0.93



- After reconstruction and cuts, SUSY background \approx signal.
 - Asymmetry halved.
- Significant luminosity required to see asymmetry in this scenario.

SUSY background - Luminosity Reach



- mSUGRA parameter space reach challenging.
- Scenarios with heavy $\tilde{g}/\tilde{q}_{1st,2nd}$ will be better.
- If SUSY spectra known, background may be estimated and subtracted.
 - Asymmetry recovered.
 - Need information on cross sections and branching ratios.