## **Resummation, Evolution, Factorization 2022**



Contribution ID: 75

Type: not specified

## Measurement of the 1-jettiness event shape observable in deep-inelastic electron-proton scattering at HERA

Tuesday 1 November 2022 17:00 (15 minutes)

A first measurement of the 1-jettiness event shape observable in neutral-current deep-inelastic electron-proton scattering is presented. The 1-jettiness observable  $\tau_{1b}$  is defined such that it is equivalent to the thrust observable defined in the Breit frame. The data were taken in the years 2003 to 2007 with the H1 detector at the HERA ep collider at a center-of-mass energy of 319 GeV and correspond to an integrated luminosity of 351.6 pb<sup>-1</sup>. The triple-differential cross sections are presented as a function of the 1-jettiness  $\tau_{1b}$ , the event virtuality  $Q^2$  and the inelasticity y in the kinematic region  $Q^2 > 150 \text{ GeV}^2$ . The data have sensitivity to the parton distribution functions of the proton, the strong coupling constant and to resummation and hadronisation effects. The data are compared to selected predictions.

H1prelim-21-032

Primary author: HESSLER, Johannes (ATLAS (LHC Experiment ATLAS))
Co-authors: SCHMITT, Stefan (ATLAS (ATLAS Scientific Computing)); H1, Collaboration (DESY)
Presenter: HESSLER, Johannes (ATLAS (LHC Experiment ATLAS))
Session Classification: Parrlel Session A: TMD in experiment