

HONEST Workshops: Hot Topics in High Energy Astrophysics



Report of Contributions

Contribution ID: 1

Type: **not specified**

Welcome

Tuesday 29 November 2022 09:00 (5 minutes)

Primary author: LOPEZ-COTO, Ruben (IAA-CSIC)

Co-author: DE ONA WILHELMI, Emma Maria (Z_GA (Gammaastronomie))

Presenters: DE ONA WILHELMI, Emma Maria (Z_GA (Gammaastronomie)); LOPEZ-COTO, Ruben (IAA-CSIC)

Session Classification: PeV Accelerators: The traditional paradigm - SNRs as PeVatrons

Contribution ID: 2

Type: **not specified**

Introduction to the motivations and topics of the workshop

Tuesday 29 November 2022 09:05 (15 minutes)

Primary author: LOPEZ-COTO, Ruben (IAA-CSIC)

Co-author: DE ONA WILHELMI, Emma Maria (Z_GA (Gammaastronomie))

Presenters: DE ONA WILHELMI, Emma Maria (Z_GA (Gammaastronomie)); LOPEZ-COTO, Ruben (IAA-CSIC)

Session Classification: PeV Accelerators: The traditional paradigm - SNRs as PeVatrons

Contribution ID: 3

Type: **not specified**

CRs around the knee - composition and latest results

Tuesday 29 November 2022 09:20 (30 minutes)

Presenter: KANG, Donghwa (KIT)

Session Classification: PeV Accelerators: The traditional paradigm - SNRs as PeVatrons

Contribution ID: 4

Type: **not specified**

Supernova Remnants in the context of Cosmic PeVatrons

Tuesday 29 November 2022 11:30 (30 minutes)

Presenter: SAFI-HARB, Samar

Session Classification: PeV Accelerators: The traditional paradigm - SNRs as PeVatrons

Contribution ID: 5

Type: **not specified**

The plasma physics of acceleration by shocks

Tuesday 29 November 2022 09:50 (30 minutes)

Presenter: BELL, Tony (University of Oxford)

Session Classification: PeV Accelerators: The traditional paradigm - SNRs as PeVatrons

Contribution ID: 6

Type: **not specified**

Interacting SNR blast waves in star forming region

Tuesday 29 November 2022 11:10 (20 minutes)

Presenter: GABICI, Stefano

Session Classification: PeV Accelerators: The traditional paradigm - SNRs as PeVatrons

Contribution ID: 7

Type: **not specified**

Maximum energy in SNR shocks

Tuesday 29 November 2022 10:50 (20 minutes)

Presenter: REVILLE, Brian (Max Planck Institute for Nuclear Physics)

Session Classification: PeV Accelerators: The traditional paradigm - SNRs as PeVatrons

Contribution ID: 8

Type: **not specified**

Stellar Cluster as PeVatrons: Experimental results with wide FoV experiments

Wednesday 30 November 2022 15:00 (20 minutes)

Presenter: YANG, Ruizhi (Max-Planck-Institut für Kernphysik)

Session Classification: PeV Accelerators: Alternative source types and large-scale diffuse emission

Contribution ID: 9

Type: **not specified**

Stellar Cluster as PeVatrons: Morphological studies of stellar clusters using IACTs

Wednesday 30 November 2022 15:20 (20 minutes)

Presenter: MOHRMANN, Lars (MPIK)

Session Classification: PeV Accelerators: Alternative source types and large-scale diffuse emission

Contribution ID: **10**

Type: **not specified**

Stellar Clusters as PeVatrons: Acceleration of CRs in Stellar Clusters

Wednesday 30 November 2022 15:40 (20 minutes)

Presenter: MORLINO, Giovanni (Gran Sasso Science Institute)

Session Classification: PeV Accelerators: Alternative source types and large-scale diffuse emission

Contribution ID: 11

Type: **not specified**

VHE view of the Galactic Centre

Wednesday 30 November 2022 16:00 (20 minutes)

Presenter: LEMIERE, ANNE (APC)

Session Classification: PeV Accelerators: Alternative source types and large-scale diffuse emission

Contribution ID: 12

Type: **not specified**

Propagation of PeV particles: Size of PeV sources

Wednesday 30 November 2022 17:50 (20 minutes)

Presenter: MERTSCH, Philipp (RWTH Aachen University, TTK)

Session Classification: PeV Accelerators: Alternative source types and large-scale diffuse emission

Contribution ID: 13

Type: **not specified**

Propagation of PeV particles: Outflows in Fermi bubbles

Wednesday 30 November 2022 17:10 (20 minutes)

Presenter: TAYLOR, Andrew (Z_THAT (Theoretische Astroteilchenphysik))

Session Classification: PeV Accelerators: Alternative source types and large-scale diffuse emission

Contribution ID: 14

Type: **not specified**

Propagation of PeV particles: Diffuse emission

Wednesday 30 November 2022 17:30 (20 minutes)

Presenter: PERON, Giada (Max Planck Institut für Kernphysik, Heidelberg)

Session Classification: PeV Accelerators: Alternative source types and large-scale diffuse emission

Contribution ID: 15

Type: **not specified**

Mechanisms of acceleration of leptons/hadrons in compact objects

Thursday 1 December 2022 09:20 (20 minutes)

Presenter: AMATO, Elena (INAF - Osservatorio Astrofisico di Arcetri)

Session Classification: Lessons learnt from experimental results and future prospects

Contribution ID: **16**

Type: **not specified**

The Crab Nebula at Very High Energies

Thursday 1 December 2022 09:00 (20 minutes)

Presenter: Dr OLMÍ, Barbara (INAF)

Session Classification: Lessons learnt from experimental results and future prospects

Contribution ID: 17

Type: **not specified**

VHE gamma-ray experimental results: Binary systems at 100 TeV

Thursday 1 December 2022 09:40 (20 minutes)

Presenter: KHANGULYAN, Dmitry (Rikkyo University)

Session Classification: Lessons learnt from experimental results and future prospects

Contribution ID: **18**

Type: **not specified**

Acceleration of leptons/hadrons in compact objects / mechanisms

Session Classification: Lessons learnt from experimental results and future prospects

Contribution ID: **19**

Type: **not specified**

Surveys: Imaging Atmospheric Cherenkov Telescopes

Thursday 1 December 2022 10:30 (20 minutes)

Presenter: DE NAUROIS, Mathieu

Session Classification: Lessons learnt from experimental results and future prospects

Contribution ID: 20

Type: **not specified**

Surveys: Extended Air Shower Arrays

Thursday 1 December 2022 10:50 (20 minutes)

Presenter: CAO, Zhen (Institute of High Energy Physics)

Session Classification: Lessons learnt from experimental results and future prospects

Contribution ID: 21

Type: **not specified**

Surveys: Future

Thursday 1 December 2022 11:10 (20 minutes)

Presenter: HINTON, Jim (MPIK)

Session Classification: Lessons learnt from experimental results and future prospects

Contribution ID: 22

Type: **not specified**

Concluding remarks

Thursday 1 December 2022 11:30 (30 minutes)

Presenters: DE ONA WILHELMI, Emma Maria (Z_GA (Gammaastronomie)); LOPEZ-COTO, Ruben (IAA-CSIC)

Session Classification: Lessons learnt from experimental results and future prospects

Contribution ID: 23

Type: **not specified**

On the localization of the CR PeVatrons inside the extended UHE gamma-ray sources

Wednesday 30 November 2022 16:50 (20 minutes)

Presenter: AHARONIAN, Felix (MPIK)

Session Classification: PeV Accelerators: Alternative source types and large-scale diffuse emission

Contribution ID: 24

Type: **not specified**

A spatially resolved study of hard X-ray emission in Kepler's SNR: indications of different regimes of particle acceleration

Synchrotron X-ray emission in young supernova remnants (SNRs) is a powerful diagnostic tool to study the population of high energy electrons accelerated at the shock front.

We performed a spatially resolved spectral analysis of NuSTAR and XMM-Newton observations of the young Kepler's SNR, aiming to study in detail its non-thermal hard X-rays emission.

We selected a set of regions all around the rim of the shell and extracted the corresponding spectra.

The spectra were analyzed by adopting a model of synchrotron radiation in the loss-limited regime, to constrain the dependence of the cutoff energy of the synchrotron radiation on the shock velocity.

We identify two different regimes of particle acceleration, characterized by different Bohm factors.

In the north, where the shock interacts with a dense circumstellar medium (CSM), we found a more efficient acceleration than in the south, where the shock velocity is higher and there are no signs of shock interaction with dense CSM.

Our results suggest an enhanced efficiency of the acceleration process in regions where the shock-CSM interaction generates an amplified and turbulent magnetic field.

By combining hard X-ray spectra with radio and γ -ray observations of Kepler's SNR, we modelled the spectral energy distribution. In the light of our results we propose that the observed γ -ray emission is mainly hadronic, and originates in the northern part of the shell.

Primary author: SAPIENZA, Vincenzo (University of Palermo)

Co-authors: Dr BAMBÀ, Aya; Dr BOCCHINO, Fabrizio; Prof. PERES, Giovanni; Prof. MICELI, Marco (UNIPA); Dr ORLANDO, Salvatore; Dr KATSUDA, Satoru; Mr NAGAYOSHI, Tsutomu; Dr TERADA, Yukikatsu

Presenter: SAPIENZA, Vincenzo (University of Palermo)

Session Classification: PeV Accelerators: The traditional paradigm - SNRs as PeVatrons

Contribution ID: 25

Type: **not specified**

The supernova remnant SN 1006 as a Galactic particle accelerator

The origin of cosmic-rays is an open issue of high-energy astrophysics. Supernova remnants are expected to be the main source of Galactic cosmic rays up to energies of about 3 PeV, provided that they transfer a significant fraction of their kinetic energy to the particles ($\sim 10\%$). In particular, the loss of such a large fraction of energy is predicted to alter the shock dynamics (shock modification) by enhancing the shock compression ratio above the canonical value of 4. The bilateral supernova remnant SN 1006 is an ideal target to study shock modification because of its evolution in a fairly uniform environment. SN 1006 shows bright synchrotron X-ray emission from ultrarelativistic electrons accelerated at the shock front in its northeastern and southwestern limbs. If efficient hadron acceleration occurs in these regions, we should observe shock modification therein. We performed a spatially resolved spectral analysis of Chandra and XMM-Newton observations of SN 1006 by selecting narrow regions between the shock front and the contact discontinuity and measuring the density of the X-ray emitting plasma. Our results show an increase of the compression ratio from the characteristic value of 4, in thermal limb, up to ~ 7 in nonthermal limbs, i.e. in regions where the ambient magnetic field is almost parallel to the shock velocity. We conclude that an efficient particle acceleration causes shock modification in quasi-parallel shocks in SN 1006. By comparing our results with state-of-the-art models, we find that SN 1006 is transferring a significant fraction of its kinetic energy to hadrons. The inferred values of compression ratios and cosmic ray slopes are consistent with those expected for modified shocks which is affected by the effects of the postcursor.

Primary author: GIUFFRIDA, Roberta (Università degli studi di Palermo - Osservatorio Astronomico di Palermo)

Co-authors: Ms DECOURCHELLE, Anne (Université Paris-Saclay, Université Paris Cité, CEA, CNRS, AIM); Mr CAPRIOLI, Damiano (Department of Astronomy and Astrophysics & Enrico Fermi Institute, The University of Chicago); Mr GRECO, Emanuele (University of Amsterdam); Mr BOCCHINO, Fabrizio (Osservatorio Astronomico di Palermo); Mr PERES, Giovanni (Università degli studi di Palermo - Osservatorio Astronomico di Palermo); Mr VINK, Jacco (Anton Pannekoek Institute, GRAPPA, University of Amsterdam); Mr MICELI, Marco (Università degli studi di Palermo - Osservatorio Astronomico di Palermo); Mr ORLANDO, Salvatore (Osservatorio Astronomico di Palermo)

Presenter: GIUFFRIDA, Roberta (Università degli studi di Palermo - Osservatorio Astronomico di Palermo)

Session Classification: PeV Accelerators: The traditional paradigm - SNRs as PeVatrons

Contribution ID: 26

Type: **not specified**

Origin of the very high energy gamma-ray emission from the Crab nebula

LHAASO has detected gamma-ray emission from the Crab Nebula up to PeV energies. We show here that our recent model for electron acceleration at pulsar wind termination shocks can fit well both the inverse Compton and the synchrotron emission from the Nebula. Integrating individual particle trajectories in a model of the magnetic field and flow pattern near the shock, we find that drift motion on the shock surface maintains either electrons or positrons on Speiser orbits in a ring-shaped region close to the equatorial plane of the pulsar, where they are accelerated up to multi-PeV energies by the first-order Fermi mechanism. We calculate the inverse Compton emission from these electrons, and demonstrate that the observed $> \text{TeV}$ gamma-ray emission from the Crab Nebula can be well reproduced for reasonable parameters of the Crab pulsar wind and turbulence levels in the nebula. Comparing to the LHAASO observations of the Crab Nebula, we can place novel constraints on parameters of the Crab pulsar wind that are still poorly known.

Primary author: GIACINTI, Gwenael (TDLI / SJTU Shanghai)

Presenter: GIACINTI, Gwenael (TDLI / SJTU Shanghai)

Session Classification: PeV Accelerators: Alternative source types and large-scale diffuse emission

Contribution ID: 27

Type: **not specified**

A bright (near) future in the PeVatron era with the ASTRI Mini-Array

Despite the enormous efforts done in very recent years, both theoretically and experimentally, the basic three questions about the Cosmic Ray (CR) origin remain without clear answers: what are their sources, how are they accelerated, how do they propagate?

Gamma-ray astronomy plays a fundamental role in this field. Both relativistic protons and electrons can emit in the gamma-ray band through different processes but only the detection of hadronic gamma-ray emission can probe CR acceleration. The recent results published by the LHAASO collaboration revealed the existence of several PeV sources. Most of them are likely related to pulsars, well known leptonic factories, and/or their nebulae, PWNe (e.g. the Crab Nebula for all), the only sources where we expect to see electrons accelerated up to PeV energies. Consequently, a gamma-ray detection at these energies still cannot be considered the final proof of hadronic acceleration. Furthermore, the limited angular resolution of LHAASO at TeV scale makes associations uncertain and, thus, more detailed and deeper studies are needed.

In this context, the ASTRI Mini-Array can play a fundamental role. The first three telescopes will be operative by fall 2023 and the full array of nine telescopes in 2025. With its unprecedented sensitivity and angular resolution in the multi-TeV band with respect to the existing IACTs, not only this instrument can extend the gamma-ray spectra of candidate PeVatrons but it can also help to distinguish whether the emission is actually connected with pulsars/PWN systems, shedding light on the nature of the highest energies emission.

Primary author: CARDILLO, MARTINA (IAPS-INAF)

Presenter: CARDILLO, MARTINA (IAPS-INAF)

Session Classification: PeV Accelerators: Alternative source types and large-scale diffuse emission

Contribution ID: 28

Type: **not specified**

Cosmic and gamma rays from young supernovae

Supernovae issued from massive star explosion produce collisionless shocks after the breakout. These shocks propagate in the dense circum-stellar wind of the progenitor star. The combination of a high density medium and shock speeds at a fraction of c make these places as potential sites of high energy cosmic ray acceleration because particles can self-drive an efficient magnetic field amplification. Under some favorable circumstances multi PeV cosmic ray energies can be reached within day timescales after the shock breakout. As Cosmic Rays propagate in dense medium they are prone to produce gamma-rays from nuclear interaction. These GeV-TeV gamma-rays if detected would be strong evidences these objects can be the missing Pevatrons. However, the gamma-ray signal detection is hampered by strong gamma-gamma absorption on soft photospheric photons within the first weeks after the explosion. A detailed time-dependent calculation is hence necessary to evaluate their detectability by the Cerenkov Telescope Array. This detectability as well as the multi-wavelength /-messenger signatures of different classes of supernovae is the main object of this talk.

Primary authors: Dr MARCOWITH, Alexandre (Laboratoire Univers Particle Montpellier); CRISTOFARI, Pierre (Observatoire de Paris)

Co-authors: Dr RENAUD, Matthieu (LUPM); Dr DWARKADAS, Vikram (University of Chicago); Dr TATISCHEFF, Vincent (IJClab)

Presenter: Dr MARCOWITH, Alexandre (Laboratoire Univers Particle Montpellier)

Session Classification: PeV Accelerators: Alternative source types and large-scale diffuse emission

Contribution ID: 29

Type: **not specified**

Gamma-ray emission from young supernova remnants in dense environments

Supernova remnants are known to accelerate cosmic rays from the detection of non-thermal emission of radio waves, X-rays, and gamma rays. The presence of cut-offs in the gamma-ray spectra of several young SNRs led to the idea that the highest energies might only be achieved during the very initial stages of a remnant's evolution. Unfortunately, the gamma-ray luminosity is assumed to peak in the first weeks after the Supernova explosion where strong $\gamma\gamma$ -absorption attenuates the observable signal. Here, we investigate to which extend the interaction of SNR-shocks with dense structures in the medium around luminous blue variable (LBV) stars can boost the gamma-ray emission months to years after the explosion.

We use the time-dependent acceleration code RATPaC to study the acceleration of cosmic rays in supernovae expanding into dense environments around massive stars. We performed spherically symmetric 1-D simulations in which we simultaneously solve the transport equations for cosmic rays, magnetic turbulence, and the hydrodynamical flow of the thermal plasma in the test-particle limit.

We investigated typical parameters of the circumstellar medium (CSM) in the freely expanding winds around LBV stars and added dense structures that arise from episodes of highly-enhanced mass-loss of the progenitors. The results are compared to calculation of smooth, unstructured winds.

We find that the interactions with the dense structures happens typically after a few months for LBV progenitors. During the interaction stage, the $\gamma\gamma$ -absorption by photons emitted from the Supernova's photosphere became negligible. The gamma-ray luminosity of the interacting SNRs can surpass the internal/unabsorbed peak-luminosity that arises shortly after the explosion. As a consequence, the observable flux can be considerably higher compared to the signal expected shortly after the explosion where $\gamma\gamma$ -absorption is important and where most gamma-ray observatories search for transient signals from these Supernovae. Further, the change of the shock-speed during the shock-shell interaction boosts the achievable maximum energy beyond a PeV, where early interactions yield higher peak-energies.

Primary author: BROSE, Robert (Dublin Institute for Advanced Studies)

Co-authors: SUSHCH, Iurii (North-West University, South Africa); MACKEY, Jonathan (Dublin Institute for Advanced Studies)

Presenter: BROSE, Robert (Dublin Institute for Advanced Studies)

Session Classification: PeV Accelerators: The traditional paradigm - SNRs as PeVatrons

Contribution ID: 30

Type: **not specified**

Supernova remnant/starformation scenarios: why it is necessary to separate the energy requirement for Galactic cosmic rays from the PeVatron requirement

Supernova remnants (SNRs) have long been considered to be the dominant source of Galactic cosmic rays, which was then implied to mean that they provided most of the energy to power cosmic rays as well as that they are capable to accelerate protons up to the cosmic-ray knee. The lack of evidence for PeV cosmic rays inside SNRs, as well as theoretical considerations, has made this scenario untenable. At the same time the latest LHAASO and IACTs results suggest that PeVatrons lurk inside starforming regions. Here I will talk about why SNRs should still be considered the main scenario for the explanation of the energy in Galactic cosmic rays, but show that cosmic-ray data allows for a second component of energies between 10-1000 TeV. This second component could be a subset of supernovae/SNRs, re-acceleration inside star forming regions, or pulsars.

Primary author: VINK, Jacco (University of Amsterdam)

Presenter: VINK, Jacco (University of Amsterdam)

Session Classification: PeV Accelerators: The traditional paradigm - SNRs as PeVatrons

Contribution ID: 31

Type: **not specified**

Search for the Galactic accelerators of Cosmic-Rays up to the Knee with the Pevatron Test Statistic

The recently introduced PeVatron Test Statistic (PTS) method in the framework of Cherenkov Telescope Array (CTA) PeVatrons study offers a new approach to detect spectral signatures of hadronic PeVatrons. The PTS is, for the first time, applied to PeVatron candidate sources' data from gamma-ray observatories to test for the origin of Cosmic Rays (CRs) at energies around and above the knee of the CR spectrum. Public data from gamma-ray observatories are analyzed jointly within hadronic emission models, assuming conservative systematic errors. It is outruled with a high statistical significance of more than 5σ that the two shell type Supernova remnants (SNRs) RX J1713.7–3946 and Vela Jr., as well as the gamma-ray source HESS J1745–290, which is spatially coincident with the dynamical center of the Galaxy, are PeVatrons. These results were previously derived by other means but the analyses confirm the performance and reliability of the PTS. Assuming that the gamma-ray emission from the region which contains the SNR G106.3+2.7 and the Boomerang nebula has a single component, a PeVatron hypothesis can be outruled with a statistical significance of more than 5σ . No statistically significant conclusion with respect to an association to a PeVatron is drawn for the diffuse gamma-ray emission around the Galactic Center (GC), and the two unidentified gamma-ray sources LHAASO J2108+5157 and HESS J1702–420A. However, it is argued that data from the Northern CTA and the Southern Wide-field Gamma-ray Observatory (SWG0) will respectively allow to decide whether these two sources are associated to a PeVatron or not. Based on the straw-man design sensitivity for SWG0, the experiment is expected to have sufficient point source sensitivity to probe large parts of the relevant parameter space for PeVatrons after 5 years of observations.

Primary authors: ANGÜNER, Ekrem Oguzhan (CPPM); SPENGLER, Gerrit (ZEU-CTA (CTA))

Presenter: ANGÜNER, Ekrem Oguzhan (CPPM)

Session Classification: PeV Accelerators: Alternative source types and large-scale diffuse emission

Contribution ID: 32

Type: **not specified**

Supernova remnant/star formation scenarios: why it is necessary to separate the energy requirement for Galactic cosmic rays from the PeVatron requirement

Tuesday 29 November 2022 13:00 (5 minutes)

Presenter: VINK, Jacco (Anton Pannekoek Institute, GRAPPA, University of Amsterdam)

Session Classification: Contributions

Contribution ID: 33

Type: **not specified**

Gamma-ray emission from young supernova remnants in dense environments

Tuesday 29 November 2022 13:05 (5 minutes)

Presenter: BROSE, Robert (Z_THAT (Theoretische Astroteilchenphysik))

Session Classification: Contributions

Contribution ID: 34

Type: **not specified**

A spatially resolved study of hard X-ray emission in Kepler's SNR: indications of different regimes of particle acceleration

Tuesday 29 November 2022 13:10 (5 minutes)

Presenter: SAPIENZA, Vincenzo (University of Palermo)

Session Classification: Contributions

Contribution ID: 35

Type: **not specified**

The supernova remnant SN 1006 as a Galactic particle accelerator

Tuesday 29 November 2022 13:15 (5 minutes)

Presenter: GIUFFRIDA, Roberta (Università degli studi di Palermo - Osservatorio Astronomico di Palermo)

Session Classification: Contributions

Contribution ID: 36

Type: **not specified**

Cosmic and gamma rays from young supernovae

Wednesday 30 November 2022 19:00 (5 minutes)

Presenter: MARCOWITH, Alexandre (Laboratoire Univers Particle Montpellier)

Session Classification: Contributions

Contribution ID: 37

Type: **not specified**

Origin of the very high energy gamma-ray emission from the Crab nebula

Tuesday 29 November 2022 13:20 (5 minutes)

Presenter: GIACINTI, Gwenaël (MPIK Heidelberg)

Session Classification: Contributions

Contribution ID: **38**

Type: **not specified**

Search for the Galactic accelerators of Cosmic-Rays up to the Knee with the Pevatron Test Statistic

Wednesday 30 November 2022 19:05 (5 minutes)

Presenter: ANGÜNER, Ekrem Oguzhan (CPPM)

Session Classification: Contributions

Contribution ID: 39

Type: **not specified**

A bright (near) future in the PeVatron era with the ASTRI Mini-Array

Wednesday 30 November 2022 19:10 (5 minutes)

Presenter: CARDILLO, MARTINA (IAPS-INAF)

Session Classification: Contributions