

Precise predictions for the trilinear Higgs self-coupling in the Standard Model and beyond

predicting κ_λ in *any* model at the one-loop order

Henning Bahl, Johannes Braathen, **Martin Gabelmann**, Georg Weiglein
DESY Theory Workshop, 28.09.2022



The SM scalar sector

> V_{SM} fixed at tree-level by $m_h \approx 125 \text{ GeV}$ and $v = (\sqrt{2}G_F)^{-1/2} \approx 246 \text{ GeV}$:

$$\begin{aligned} V_{\text{SM}}(h) &= \frac{m_h^2}{2} h^2 + 3 \frac{m_h^2}{v} h^3 + 3 \frac{m_h^2}{v^2} h^4 \\ &= \frac{m_h^2}{2} h^2 + \lambda_{hhh} h^3 + \lambda_{hhhh} h^4 \end{aligned}$$

> However: λ_{hhh} (and λ_{hhhh}) experimentally unknown.

The SM scalar sector

> V_{SM} fixed at tree-level by $m_h \approx 125 \text{ GeV}$ and $v = (\sqrt{2}G_F)^{-1/2} \approx 246 \text{ GeV}$:

$$\begin{aligned} V_{\text{SM}}(h) &= \frac{m_h^2}{2} h^2 + 3 \frac{m_h^2}{v} h^3 + 3 \frac{m_h^2}{v^2} h^4 \\ &= \frac{m_h^2}{2} h^2 + \lambda_{hhh} h^3 + \lambda_{hhhh} h^4 \end{aligned}$$

> However: λ_{hhh} (and λ_{hhhh}) experimentally unknown.


> BSM case: deformation of the scalar potential possible!

$$V_{\text{BSM}}(h, \dots) = \frac{m_h^2}{2} h^2 + 3 \frac{m_h^2}{v} \kappa_{\lambda}^{\text{BSM}} h^3 + 3 \frac{m_h^2}{v^2} \kappa_{2\lambda} h^4 + \dots$$

$\kappa_{\lambda}^{\text{BSM}}$: describes deviation from SM: $\kappa_{\lambda}^{\text{BSM}} = \frac{\lambda_{hhh}^{\text{BSM}}}{\lambda_{hhh}^{\text{SM}}}$ in model "BSM".

> $|\kappa_{\lambda}| \gg 1$ possible even though other (known) couplings behave very SM-like

Higgs pair production (in the alignment limit)




$$pp \rightarrow hh: -0.4 < \kappa_\lambda < 6.3 \quad [\text{ATLAS-CONF-2022-050}]$$

When to apply the κ_λ -constraint to BSM models?

> only κ_λ is *significantly* modified by BSM physics

> all other couplings SM-like

→ in many models given by experimental constraints



Higher-order corrections to $\lambda_{hhh}^{\text{BSM}}$

A few studies of the trilinear Higgs self-couplings (i.e. κ_λ) already exists:

- > SM [Kanemura et al. '04]
- > SM + singlet [Kanemura et al. '16]
- > THDM [Kanemura et al. '04][Basler et al. '17][Braathen et al. '19]
- > THDM + singlet [Basler et al. '19]
- > Triplet extensions [Aoki et al. '18] [Chiang et al. '18]
- > MSSM [Brucherseifer et al. '13]
- > NMSSM [Dao et al. '13] [Dao et al. '15][Dao et al. '22 (tbp)]

of which many can have sizeable deviations from $\kappa_\lambda = 1$ and hence also from $\sigma^{\text{SM}}(hh)$ [Abouabid et al. '21]

Higher-order corrections to $\lambda_{hhh}^{\text{BSM}}$

A few studies of the trilinear Higgs self-couplings (i.e. κ_λ) already exists:

- > SM [Kanemura et al. '04]
- > SM + singlet [Kanemura et al. '16]
- > THDM [Kanemura et al. '04][Basler et al. '17][Braathen et al. '19]
- > THDM + singlet [Basler et al. '19]
- > Triplet extensions [Aoki et al. '18] [Chiang et al. '18]
- > MSSM [Brucherseifer et al. '13]
- > NMSSM [Dao et al. '13] [Dao et al. '15][Dao et al. '22 (tbp)]

of which many can have sizeable deviations from $\kappa_\lambda = 1$ and hence also from $\sigma^{\text{SM}}(hh)$ [Abouabid et al. '21]

However, there many more models to explore:

- > THDM-variants
- > singlet-variants
- > extended gauge sectors
- > extended fermion sectors
- > non-minimal SUSY, e.g. seesaw extensions, dirac gauginos, SplitSUSY, ...
- > (+ combinations)
- > ...

Higher-order corrections to $\lambda_{hhh}^{\text{BSM}}$

A few studies of the trilinear Higgs self-couplings (i.e. κ_λ) already exists:

- > SM [Kanemura et al. '04]
- > SM + singlet [Kanemura et al. '16]
- > THDM [Kanemura et al. '04][Basler et al. '17][Braathen et al. '19]
- > THDM + singlet [Basler et al. '19]
- > Triplet extensions [Aoki et al. '18] [Chiang et al. '18]
- > MSSM [Brucherseifer et al. '13]
- > NMSSM [Dao et al. '13] [Dao et al. '15][Dao et al. '22 (tbp)]

of which many can have sizeable deviations from $\kappa_\lambda = 1$ and hence also from $\sigma^{\text{SM}}(hh)$ [Abouabid et al. '21]

However, there many more models to explore:


- > THDM-variants
- > singlet-variants
- > extended gauge sectors
- > extended fermion sectors
- > non-minimal SUSY, e.g. seesaw extensions, dirac gauginos, SplitSUSY, ...
- > (+ combinations)
- > ...

→ **framework to calculate $\lambda_{hhh}^{\text{BSM}}$ for large class of "BSM's"**

Ingredients


Ingredients

$(\lambda_{hhh}^{\text{BSM}})^{\text{one-loop}}$




Example ingredient: generic fermion triangle

Idea: compute *generic* diagrams i.e. assume most generic



Example ingredient: generic fermion triangle


Idea: compute *generic* diagrams i.e. assume most generic



- > couplings $C_i = P_L C_i^L + P_R C_i^R$, $P_{R/L} = \frac{1 \pm \gamma_5}{2}$
- > as well as loop-masses m_{f_i} and
- > external momenta p_i , $i = 1, 2, 3$.

Example ingredient: generic fermion triangle

Idea: compute *generic* diagrams i.e. assume most generic



> couplings $C_i = P_L C_i^L + P_R C_i^R$, $P_{R/L} = \frac{1 \pm \gamma_5}{2}$


> as well as loop-masses m_{f_i} and

> external momenta p_i , $i = 1, 2, 3$.

$$\begin{aligned}
 &= 2\mathbf{B0}(p_3^2, m_2^2, m_3^2)(C_1^L(C_2^L C_3^R m_{f_1} + C_2^R C_3^L m_{f_2} + C_2^R C_3^R m_{f_3}) + C_1^R(C_2^R C_3^L m_{f_1} + \\
 &C_2^L C_3^L m_{f_2} + C_2^L C_3^R m_{f_3})) + m_{f_1} \mathbf{C0}(p_2^2, p_3^2, p_1^2, m_1^2, m_3^2, m_2^2)((C_1^L C_2^L C_3^R + \\
 &C_1^R C_2^R C_3^L)(p_1^2 + p_2^2 - p_3^2) + 2(C_1^L C_2^L C_3^L + C_1^R C_2^R C_3^R)m_{f_2} m_{f_3} + \\
 &2m_{f_1}(C_1^L(C_2^L C_3^R m_{f_1} + C_2^R C_3^L m_{f_2} + C_2^R C_3^R m_{f_3}) + C_1^R(C_2^R C_3^L m_{f_1} + C_2^L C_3^L m_{f_2} + \\
 &C_2^L C_3^R m_{f_3}))) + \mathbf{C1}(p_2^2, p_3^2, p_1^2, m_1^2, m_3^2, m_2^2)(2p_2^2(C_1^L C_3^R(C_2^L m_{f_1} + C_2^R m_{f_2}) + \\
 &C_1^R C_3^L(C_2^R m_{f_1} + C_2^L m_{f_2})) + (p_1^2 + p_2^2 - p_3^2)((C_1^L C_2^L C_3^R + C_1^R C_2^R C_3^L)m_{f_1} + \\
 &(C_1^L C_2^R C_3^L + C_1^R C_2^L C_3^R)m_{f_3})) + \mathbf{C2}(p_2^2, p_3^2, p_1^2, m_1^2, m_3^2, m_2^2)((p_1^2 + p_2^2 - \\
 &p_3^2)(C_1^L C_3^R(C_2^L m_{f_1} + C_2^R m_{f_2}) + C_1^R C_3^L(C_2^R m_{f_1} + C_2^L m_{f_2})) + 2p_1^2((C_1^L C_2^L C_3^R + \\
 &C_1^R C_2^R C_3^L)m_{f_1} + (C_1^L C_2^R C_3^L + C_1^R C_2^L C_3^R)m_{f_3}))
 \end{aligned}$$

Example ingredient: generic fermion triangle

Idea: compute *generic* diagrams i.e. assume most generic



> couplings $C_i = P_L C_i^L + P_R C_i^R$, $P_{R/L} = \frac{1 \pm \gamma_5}{2}$

> as well as loop-masses m_{f_i} and

> external momenta p_i , $i = 1, 2, 3$.

$$\begin{aligned}
 &= 2\mathbf{B0}(p_3^2, m_2^2, m_3^2)(C_1^L(C_2^L C_3^R m_{f_1} + C_2^R C_3^L m_{f_2} + C_2^R C_3^R m_{f_3}) + C_1^R(C_2^R C_3^L m_{f_1} + \\
 &C_2^L C_3^L m_{f_2} + C_2^L C_3^R m_{f_3})) + m_{f_1} \mathbf{C0}(p_2^2, p_3^2, p_1^2, m_1^2, m_3^2, m_2^2)((C_1^L C_2^L C_3^R + \\
 &C_1^R C_2^R C_3^L)(p_1^2 + p_2^2 - p_3^2) + 2(C_1^L C_2^L C_3^L + C_1^R C_2^R C_3^R)m_{f_2} m_{f_3} + \\
 &2m_{f_1}(C_1^L(C_2^L C_3^R m_{f_1} + C_2^R C_3^L m_{f_2} + C_2^R C_3^R m_{f_3}) + C_1^R(C_2^R C_3^L m_{f_1} + C_2^L C_3^L m_{f_2} + \\
 &C_2^L C_3^R m_{f_3}))) + \mathbf{C1}(p_2^2, p_3^2, p_1^2, m_1^2, m_3^2, m_2^2)(2p_2^2(C_1^L C_3^R(C_2^L m_{f_1} + C_2^R m_{f_2}) + \\
 &C_1^R C_3^L(C_2^R m_{f_1} + C_2^L m_{f_2})) + (p_1^2 + p_2^2 - p_3^2)((C_1^L C_2^L C_3^R + C_1^R C_2^R C_3^L)m_{f_1} + \\
 &(C_1^L C_2^R C_3^L + C_1^R C_2^L C_3^R)m_{f_3})) + \mathbf{C2}(p_2^2, p_3^2, p_1^2, m_1^2, m_3^2, m_2^2)((p_1^2 + p_2^2 - \\
 &p_3^2)(C_1^L C_3^R(C_2^L m_{f_1} + C_2^R m_{f_2}) + C_1^R C_3^L(C_2^R m_{f_1} + C_2^L m_{f_2})) + 2p_1^2((C_1^L C_2^L C_3^R + \\
 &C_1^R C_2^R C_3^L)m_{f_1} + (C_1^L C_2^R C_3^L + C_1^R C_2^L C_3^R)m_{f_3}))
 \end{aligned}$$

> insert concrete BSM model

> evaluate with the help of COLLIER^[Denner et al. '16]

> public code anyH3^[Bahl, Braathen, MG, Weiglein tbp]

Feature list (so far) of anyH3

> import/convert arbitrary UFO models

> definition of renormalisation schemes

```
# schemes.yml
renormalization_schemes:
  OS:
    mass_counterterms:
      h1: OS
      h2: OS
    VEV_counterterm: OS
  MS:
    mass_counterterms:
      h1: MS
      h2: MS
    VEV_counterterm: MS
```

> optional: full p^2 dependence

> numerical / analytical / \LaTeX outputs

> restrict to certain topologies


> restrict to certain particles in the loop

> python-library with command-line- and Mathematica-interface

```
from anyBSM import anyH3
myfancymodel = anyH3(
    'path/to/UFO/model',
    scheme = 'OS')
result = myfancymodel.lambdahhh()
```


> ...

Decoupling in the alignment limit



- > alignment means $\kappa_{\lambda}^{\text{tree-level}} = 1$
- > recover SM result for $M_{\text{BSM}} \rightarrow \infty$
- > many models present and cross-checked
- > easy to implement new models (UFO)

Non-decoupling in the alignment limit



- > mass splitting within the same multiplet
- > induces large couplings for $M_{BSM} \rightarrow \infty$
- > corrections large-enough to exclude otherwise unconstrained parameter space
- > (see also talk by Johannes Braathen tomorrow)

Summary

- > developed computer code anyH3 (anyBSM) for λ_{hhh} in arbitrary ren. QFTs
 - at the full one-loop order
 - with arbitrary choice of renormalization schemes
- > uses UFO input (generate with SARAH, FeynRules or use a custom one)
- > analytical results; fast numerical results; SM: $\mathcal{O}(0.2\text{ s})$, MSSM: $\mathcal{O}(0.5\text{ s})$
- > many models already implemented:
SM, SM+**singlet**, **THDM**, **NTHDM**, various **triplet extensions** and **MSSM**
- > reproduced known results in the SM, SSM, THDM and TSM

Summary

- > developed computer code anyH3 (anyBSM) for λ_{hhh} in arbitrary ren. QFTs
 - at the full one-loop order
 - with arbitrary choice of renormalization schemes
- > uses UFO input (generate with SARAH, FeynRules or use a custom one)
- > analytical results; fast numerical results; SM: $\mathcal{O}(0.2\text{ s})$, MSSM: $\mathcal{O}(0.5\text{ s})$
- > many models already implemented:
SM, SM+**singlet**, **THDM**, **NTHDM**, various **triplet extensions** and **MSSM**
- > reproduced known results in the SM, SSM, THDM and TSM

Future ideas / todos:

- > publish
- > more models / cross-checks
- > go beyond one-loop
- > non-SM self-couplings (e.g. $\kappa_{\lambda_{Hhh}}$)
- > κ_t and κ_{tt}

Backup

Another ingredient: Renormalisation of λ_{hhh}

$$\delta\lambda_{hhh}^{\text{CT}} = \text{---} \bigcirc \text{---} = ?$$

- > one-loop \rightarrow renormalisation of all parameters entering $\lambda_{hhh}^{(0),\text{BSM}}$
- > reminder: $\lambda_{hhh}^{(0),\text{SM}} = 3 \frac{m_h^2}{v^2}$
- > \rightarrow generalization (reminder: quasi-alignment!):

$$\lambda_{hhh}^{(0),\text{BSM}} \equiv \lambda_{hhh}^{(0),\text{BSM}} \left(\underbrace{v^{\text{SM}}, m_h^{\text{SM}}}_{\text{SM Higgs sector}}, m_{H_i}^{\text{BSM}}, v_i^{\text{BSM}}, \dots \right)$$

\swarrow BSM masses \swarrow vevs \swarrow

- > user's choice:
 - SM sector: fully OS **or** $\overline{\text{MS}}/\overline{\text{DR}}$
 - BSM masses: OS **or** $\overline{\text{MS}}/\overline{\text{DR}}$
 - additional couplings/vevs: most likely $\overline{\text{MS}}$ **but also custom ren. conditions possible!**
- couplings etc. (that can't be expressed in terms of masses)

$$\delta\lambda_{hhh}^{\text{CT}} = \sum_x \frac{\partial \lambda_{hhh}^{(0),\text{BSM}}}{\partial x} \delta x, \quad x = (m_h^{\text{SM}})^{\text{OS}/\overline{\text{MS}}}, (v^{\text{SM}})^{\text{OS}/\overline{\text{MS}}}, (m_{H_i}^{\text{BSM}})^{\text{OS}/\overline{\text{MS}}}, (\dots)^{\overline{\text{MS}}/\text{custom}}$$

(Default) Renormalization choice of $(v^{\text{SM}})^{\text{OS}}$ and $(m_i^2)^{\text{OS}}$

- > $v^{\text{OS}} \equiv \frac{2M_W^{\text{OS}}}{e} \sqrt{1 - \frac{M_W^{2\text{OS}}}{M_Z^{2\text{OS}}}}$ with
 - $\delta^{(1)} M_V^{2\text{OS}} = \frac{\Pi_V^{(1),T}}{M_V^{2\text{OS}}}(p^2 = M_V^{2\text{OS}}), V = W, Z$
 - $\delta^{(1)} e^{\text{OS}} = \frac{1}{2} \dot{\Pi}_\gamma(p^2 = 0) + \text{sign}(\sin \theta_W) \frac{\sin \theta_W}{M_Z^2 \cos \theta_W} \Pi_{\gamma Z}(p^2 = 0)$
- > attention (i): $\rho^{\text{tree-level}} \neq 1 \rightarrow$ further CTs needed (depends on the model)
 \rightarrow ability to define *custom* renormalisation conditions
- > scalar masses: $m_i^{\text{OS}} = m_i^{\text{pole}}$
 - $\delta^{\text{OS}} m_i^2 = -\widetilde{\text{Re}} \Sigma_{h_i}^{(1)}|_{p^2=m_i^2}$
 - $\delta^{\text{OS}} Z_i = \widetilde{\text{Re}} \frac{\partial}{\partial p^2} \Sigma_{h_i}^{(1)}|_{p^2=m_i^2}$
- > attention (ii): scalar mixing may also require further CTs/tree-level relations

All bosonic one- & two-point functions and their derivatives for general QFTs are required for flexible OS renormalisation.


Tadpole contributions to λ_{hhh}


- > In the SM: once λ_{hhh} is expressed in terms of *physical* input parameters, its result is independent of the treatment of the tadpoles:

$$\delta^{(1)}\lambda_{hhh} \supset -\frac{3}{v^2}\delta^{(1)}t_{\text{finite}}$$


- > However: UFO models do (often) **not** contain the explicit dependence on the tree-level/one-loop/one-loop-CT tadpoles.
- > Thus: we choose the Fleischer-Jegerlehner treatment $t^{\text{tree-level}} = 0$ and renormalize $\delta^{(1)}t^{\text{CT}}|_{\text{finite}} = 0$ in the $\overline{\text{MS}}$ scheme per default (can also turn-off automatic tadpoles and implement own scheme).

→ only need to take into account tadpole contributions to all two- and three-point functions:







Workflow



Simplest model to consider for cross-check: SM



Leading two-loop $\mathcal{O}(\alpha_t(\alpha_t + \alpha_s))^{\text{OS}}$: $\mathcal{O}(+1.4\%)$ [Senaha '18] [Braathen et al. '19]

W Mass

- > start with HO corrections to muon decay: $M_W^2 \left(1 - \frac{M_W^2}{M_Z^2}\right) = \frac{\pi\alpha_{\text{em}}}{\sqrt{2}G_F} [1 + \Delta r]$
- > and solve for: $M_W^2 = M_Z^2 \left[\frac{1}{2} + \sqrt{\frac{1}{4} - \frac{\pi\alpha_{\text{em}}}{\sqrt{2}G_F M_Z^2} (1 + \Delta r)} \right]$
- > with: $\Delta r^{(1)} = 2\delta^{(1)} e + \frac{\Pi_W^{(1),T}(0) - \delta^{(1)} M_W^2}{M_W^2} - \frac{\delta^{(1)} \sin^2 \theta_W}{\sin^2 \theta_W} + \delta_{\text{vertex+box}}$
- > and: $\frac{\delta^{(1)} \sin^2 \theta_W}{\sin^2 \theta_W} = \frac{\cos^2 \theta_W}{\sin^2 \theta_W} \left(\frac{\Pi_W^{(1),T}(M_W^2)}{M_W^2} - \frac{\Pi_W^{(1),T}(M_Z^2)}{M_Z^2} \right)$

It's all there but:

- > $\delta_{\text{vertex+box}}^{\text{SM}} = -\frac{2 \text{sign}(\sin \theta_W)}{\cos \theta_W \sin \theta_W M_Z^2} \Pi_{Z\gamma}(p^2 = 0) + \frac{\alpha_{\text{QED}}}{4\pi \sin^2 \theta_W} \left(6 + \frac{7-4 \sin^2 \theta_W}{2 \sin^2 \theta_W} \right) \log(\cos^2 \theta_W)$
- > $\delta_{\text{vertex+box}}^{\text{BSM}} = \text{needs to be implemented}$

However:

- > in many models $\Delta r \supset \frac{\delta \sin^2 \theta_W}{\sin \theta_W} \approx \delta \rho$ is the dominant effect!