


Status of Deeply Inelastic Parton Distributions

Johannes Blümlein
DESY



- Introduction and Method
- QCD Analysis of Unpolarized Structure Functions
- Λ_{QCD} and $\alpha_s(M_Z^2)$
- What would we like to know ?

DEEPLY INELASTIC SCATTERING



space – like process : $q^2 = (l-l')^2 = -Q^2 < 0$ $W^2 = (p+q)^2 \geq M_p^2$

$$x = \frac{Q^2}{2p \cdot q}, \quad y = \frac{p \cdot q}{p \cdot l} \quad 0 \leq x, y \leq 1$$

DIS: Microscopy of the Nucleon


- determination of all quark densities and the gluon distribution
- determination of all polarized parton densities

DIS: Fundamental Tests of QCD

- precision measurement of Λ_{QCD} and $\alpha_s(M_Z^2)$
- Thorough verification of the prediction of the light cone expansion: to higher twist
- Test of linear and non-linear resummations

Challenges for Theory: perturbative and non-perturbative

- higher order precision calculations and data analysis
- Lattice gauge theory results for Λ_{QCD} and hadronic ME



Highest order corrections of HO QCD in DIS

- Running α_s : $O(\alpha_s^4)$ Larin, van Ritbergen, Vermaseren 1997
- Unpol. anomalous dimensions and Wilson coefficients: $O(\alpha_s^3)$
Moch, Vermaseren, Vogt 2004/05
- Unpol. NS anomalous dimension 2nd Moment: $O(\alpha_s^4)$ Baikov, Chetyrkin 2006
- Pol. anomalous dimension: $O(\alpha_s^2)$; $\Delta P_{NS}^{qq}, \Delta P_{qG}$: $O(\alpha_s^3)$ Mertig, van Neerven, 1995;
Vogelsang 1995; Moch, Vermaseren, Vogt
- Pol. Wilson coefficients: $O(\alpha_s^2)$; $\Delta C_{NS}^{qq}, \Delta C_{qG}$: van Neerven, Zijlstra 1994 $O(\alpha_s^3)$ to come
- Transversity: $O(\alpha_s^2)$, some moments anom. dim.: $O(\alpha_s^3)$, Hayashigaki, Kanazawa, Koike;
Kumano, Miyama; Vogelsang; 1997; Gracey 2006
- Unpol. Heavy Flavor Wilson Coefficients: $O(\alpha_s^2)$ Laenen, van Neerven, Riemersma, Smith, 1993
Fast Mellin Space code: Blümlein & Alekhin, 2003
- Pol. Heavy Flavor Wilson Coefficients: $O(\alpha_s^1)$, Watson 1982
- $Q^2 \gg m^2$ Pol. Heavy Flavor Wilson Coefficient : $O(\alpha_s^2)$ van Neerven, Smith et al. 1996,
Blümlein & Klein 2007
- $Q^2 \gg m^2$ Unpol. Heavy Flavor Wilson Coefficient F_L : $O(\alpha_s^3)$
Blümlein, De Freitas, van Neerven, S. Klein 2005

DIS Structure Functions @ Twist 2

$$F_j(x, Q^2) = \hat{f}_i(x, \mu^2) \otimes \sigma_j^i \left(\alpha_s, \frac{Q^2}{\mu^2}, x \right)$$

↑ bare pdf ↑ sub – system cross – sect.

$$= \hat{f}_i(x, \mu^2) \otimes \Gamma_k^i \left(\alpha_s(R^2), \frac{M^2}{\mu^2}, \frac{M^2}{R^2} \right)$$

finite pdf $\equiv f_k$

$$\otimes C_j^k \left(\alpha_s(R^2), \frac{Q^2}{\mu^2}, \frac{M^2}{R^2}, x \right)$$

finite Wilson coefficient

Move to Mellin space :

$$F_j(N) = \int_0^1 dx x^{N-1} F_j(x)$$

Diagonalization of the convolutions \otimes into ordinary products.

Evolution Equations

$$\left[M \frac{\partial}{\partial M} + \beta(g) \frac{\partial}{\partial g} - 2\gamma_\psi(g) \right] F_i(N) = 0$$

$$\left[M \frac{\partial}{\partial M} + \beta(g) \frac{\partial}{\partial g} + \gamma_\kappa^N(g) - 2\gamma_\psi(g) \right] f_k(N) = 0$$

$$\left[M \frac{\partial}{\partial M} + \beta(g) \frac{\partial}{\partial g} - \gamma_\kappa^N(g) \right] C_j^k(N) = 0$$

CALLAN–SYMNANZIK equations for mass factorization


\equiv ALTARELLI–PARISI evolution equations

x-space :

$$\frac{d}{d \log(\mu^2)} \begin{pmatrix} q^+(x, Q^2) \\ G(x, Q^2) \end{pmatrix} = \frac{\alpha_s}{2\pi} \mathbf{P}(x, \alpha_s) \otimes \begin{pmatrix} q^+(x, Q^2) \\ G(x, Q^2) \end{pmatrix}$$

$$\mathbf{P}(x, \alpha_s) = \mathbf{P}^{(0)}(x) + \frac{\alpha_s}{2\pi} \mathbf{P}^{(1)}(x) + \left(\frac{\alpha_s}{2\pi} \right)^2 \mathbf{P}^{(2)}(x) + \dots$$

Anomalous Dimensions and Wilson Coefficients



Vermaseren, Moch, Vogt 2004

The Basic Functions of massless QCD to $w=5 \equiv 3$ Loops

Representative : $S_1(N) = \psi(N+1) + \gamma_E$ and its derivatives.

Weight $w=3$:

$$F_1(N) = \mathbf{M} \left[\frac{\ln(1+x)}{1+x} \right] (N)$$

$$F_2(N) = \mathbf{M} \left[\frac{\text{Li}_2(x)}{1+x} \right] (N), \quad F_3(N) = \mathbf{M} \left[\left(\frac{\text{Li}_2(x)}{1-x} \right)_+ \right] (N)$$

Yndurain et al., 1981: $F_2(N)$

Weight $w=4$:

$$F_4(N) = \mathbf{M} \left[\frac{S_{1,2}(x)}{1+x} \right] (N), \quad F_5(N) := \mathbf{M} \left[\left(\frac{S_{1,2}(x)}{1-x} \right)_+ \right] (N)$$

$F_3(N) - F_5(N)$: J.B., S. Moch, 2003; J.B., V. Ravindran ,2004

Weight w=5 :

$$F_{6,7}(N) = \mathbf{M} \left[\left(\frac{\text{Li}_4(x)}{1 \pm x} \right)_{(+)} \right] (N), \quad F_8(N) = \mathbf{M} \left[\frac{S_{1,3}(x)}{1+x} \right] (N),$$

$$F_{9,10}(N) = \mathbf{M} \left[\left(\frac{S_{2,2}(x)}{1 \pm x} \right)_{(+)} \right] (N), \quad F_{11}(N) = \mathbf{M} \left[\frac{\text{Li}_2^2(x)}{1+x} \right] (N),$$

$$F_{12,13}(N) := \mathbf{M} \left[\left(\frac{S_{2,2}(-x) - \text{Li}_2^2(-x)/2}{1 \pm x} \right)_{(+)} \right] (N)$$

$F_6(N) - F_{13}(N)$: J.B., S. Moch, 2004.

Massless QCD to 3 Loops depends on 14 Functions.

⇒ Representation for 3 Loop Wilson Coefficients under way.

Complex Analysis of these Functions


- Construct exact analytic continuations to **complex N**
- The functions are meromorphic
(up to soft corrections, which have a simple structure)
- Asymptotic Representation
- Recursion $z + 1 \rightarrow z$
- Solve the Evolution Equations fully analytically and form an **analytic expression** for the Structure functions in Mellin Space at all Q^2
- Include the **heavy flavor** Wilson coefficients in Mellin Space
- Perform a **single** fast, numerical Mellin inversion
(at high precision)



\Rightarrow Fastest and most Precise Way of Analysis

2. QCD Analysis of Unpolarized Parton Distributions


DIS range
Nucleon structure:

$$10^{-5} < x < 0.9,$$
$$1 < Q^2 < 50.000 \text{ GeV}^2$$







Parton Distributions: Overview





H1

World Data Analysis: Valence Distributions



World data:
NS-analysis

$$W^2 > 12.5 \text{ GeV}^2, Q^2 > 4 \text{ GeV}^2$$



N^3LO :

$$\alpha_s(M_Z^2) = 0.1141^{+0.0020}_{-0.0022}$$

J.B., H. Böttcher,
A. Guffanti,
(hep-ph/0607200)

Why an $O(\alpha_s^4)$ analysis can be performed?

assume an $\pm 100\%$ error on the Pade approximant $\longrightarrow \pm 2 \text{ MeV}$ in Λ_{QCD}





$$\gamma_n^{approx:3} = \frac{\gamma_n^{(2)^2}}{\gamma_n^{(1)}}$$

Baikov & Chetyrkin, April 2006:



$$\begin{aligned} \gamma_2^{3;NS} = & \frac{32}{9}a_s + \frac{9440}{243}a_s^2 + \left[\frac{3936832}{6561} - \frac{10240}{81}\zeta_3 \right] a_s^3 \\ & + \left[\frac{1680283336}{1777147} - \frac{24873952}{6561}\zeta_3 + \frac{5120}{3}\zeta_4 - \frac{56969}{243}\zeta_5 \right] a_s^4 \end{aligned}$$

The results agree better than 20%.



Valence Distributions



Valence Distributions





Valence Distributions: higher twist



- agreement between p and d analysis
- LGT determination of interest

Flavor distributions: light quarks




J. Stirling, 2004

More work needed.

HERMES probably could measure $s(x, Q^2)$ in an independent way.


Flavor distributions: light quarks



CTEQ 2007


DIS Parton Distributions ...

Charm




$F_2^{c\bar{c}}(x, Q^2)$ will be very well measured at HERA.

Slope of F_2 at low x






H1



J.B., A. Guffanti 2005


Very likely, that the $\overline{\text{MS}}$ -gluon is remains positive!

Perturbative or non-perturbative growth?




- H1 QEDC 1997 \oplus E665
- \triangle H1 1997 \star NMC
- \circ H1 SV 1995 \square SLAC
- ZEUS BPT — ALLM97

Gluon Density



H1




MRST 02 vs CTEQ 6

More work needed; $\overline{\text{MS}}$ - vs scheme-invariant evolution.

$F_L(x, Q^2)$ could be decisive.

Gluon Density



Not both distributions can be correct.

$F_L(x, Q^2)$ could be decisive.

MRST06 vs Alekhin: 2006

More work needed ! BFG Analysis in progress.

Moments of PDF's: PT + data

| f | n | This Fit N ³ LO | MRST04 NNLO | A02 NNLO | | Moment | BB, NLO |
|-------------|-----|-------------------------------|----------------|-------------|---------------------------|--------|--------------------|
| u_v | 2 | 0.3006 ± 0.0031 | 0.285 | 0.304 | Δu_v | 0 | 0.926 |
| | 3 | 0.0877 ± 0.0012 | 0.082 | 0.087 | | 1 | 0.163 ± 0.014 |
| | 4 | 0.0335 ± 0.0006 | 0.032 | 0.033 | | 2 | 0.055 ± 0.006 |
| d_v | 2 | 0.1252 ± 0.0027 | 0.115 | 0.120 | Δd_v | 0 | -0.341 |
| | 3 | 0.0318 ± 0.0009 | 0.028 | 0.028 | | 1 | -0.047 ± 0.021 |
| | 4 | 0.0106 ± 0.0004 | 0.009 | 0.010 | | 2 | -0.015 ± 0.009 |
| $u_v - d_v$ | 2 | 0.1754 ± 0.0041 | 0.171 | 0.184 | $\Delta u_v - \Delta d_v$ | 0 | 1.267 |
| | 3 | 0.0559 ± 0.0015 | 0.055 | 0.059 | | 1 | 0.210 ± 0.025 |
| | 4 | 0.0229 ± 0.0007 | 0.022 | 0.024 | | 2 | 0.070 ± 0.011 |

J.B., H. Böttcher, A. Guffanti, 2006

J.B., H. Böttcher, 2002


Lattice Results : developping; different fermion-types studied.
Low values of m_π crucial; values approach 270 MeV now.

3. Λ_{QCD} and $\alpha_s(M_Z^2)$

$$\frac{\delta\alpha_{\text{em}}(0)}{\alpha_{\text{em}}(0)} \sim 3 \cdot 10^{-11}$$

$$\frac{\delta\alpha_{\text{weak}}}{\alpha_{\text{weak}}} \sim 7 \cdot 10^{-4}$$

$$\frac{\delta\alpha_s(M_Z^2)}{\alpha_s(M_Z^2)} > 2 \cdot 10^{-2}$$



P. Zerwas, 2004

Overview of the Analyses

- Various NLO analyses; \Rightarrow Precision requires NNLO analysis and higher!
- Mixed S- and NS-NNLO analyses $e(\mu)N$ world data
- S- and NS-NNLO moment analyses νN world data
- NS-N³LO analysis $e(\mu)N$ world data
- NLO analyses polarized $e(\mu)N$ world data
- Lattice measurements

$$\alpha_s(M_Z^2)$$

| NLO | $\alpha_s(M_Z^2)$ | expt | theory | Ref. |
|----------|-------------------|--------------|----------------------|------|
| CTEQ6 | 0.1165 | ± 0.0065 | | [1] |
| MRST03 | 0.1165 | ± 0.0020 | ± 0.0030 | [2] |
| A02 | 0.1171 | ± 0.0015 | ± 0.0033 | [3] |
| ZEUS | 0.1166 | ± 0.0049 | | [4] |
| H1 | 0.1150 | ± 0.0017 | ± 0.0050 | [5] |
| BCDMS | 0.110 | ± 0.006 | | [6] |
| GRS | 0.112 | | | [10] |
| BBG | 0.1148 | ± 0.0019 | | [9] |
| BB (pol) | 0.113 | ± 0.004 | $+0.009$ -0.006 | [7] |

NLO

| NNLO | $\alpha_s(M_Z^2)$ | expt | theory | Ref. |
|-------------------|-------------------|----------------------|--------------|------|
| MRST03 | 0.1153 | ± 0.0020 | ± 0.0030 | [2] |
| A02 | 0.1143 | ± 0.0014 | ± 0.0009 | [3] |
| SY01(ep) | 0.1166 | ± 0.0013 | | [8] |
| SY01(ν N) | 0.1153 | ± 0.0063 | | [8] |
| GRS | 0.111 | | | [10] |
| A06 | 0.1128 | ± 0.0015 | | [11] |
| BBG | 0.1134 | $+0.0019 / - 0.0021$ | | [9] |
| N ³ LO | $\alpha_s(M_Z^2)$ | expt | theory | Ref. |
| BBG | 0.1141 | $+0.0020 / - 0.0022$ | | [9] |

NNLO and N³LO

BBG: $N_f = 4$: non-singlet data-analysis at $O(\alpha_s^4)$: $\Lambda = 234 \pm 26 \text{ MeV}$


Lattice results :

Alpha Collab: $N_f = 2$ Lattice; non-pert. renormalization $\Lambda = 245 \pm 16 \pm 16 \text{ MeV}$

QCDSF Collab: $N_f = 2$ Lattice, pert. reno. $\Lambda = 261 \pm 17 \pm 26 \text{ MeV}$

Lepage et al.: Larger Values, to be discussed.


$$\alpha_s(M_Z^2)$$



J.B., H. Böttcher, A. Guffanti, 2006

More Global Analyses

- $\alpha_s(M_Z^2)$ for different data sets included are too different !
⇒ applies also to HERA: IS vs FS; and also DIS vs TEVATRON-jet



M. Cooper-Sarkar, 2005




4. The Needs : What would we like to know ?

HERA:

- Collect high luminosity for $F_2(x, Q^2)$, $F_2^{c\bar{c}}(x, Q^2)$, $g_2^{c\bar{c}}(x, Q^2)$, and measure $h_1(x, Q^2)$.
- Measure : $F_L(x, Q^2)$. This is a key-question for HERA.

$$F_L(x, Q^2)$$

M. Klein, 2004: Projection for a possible measurement at HERA
 \Rightarrow of central importance to study the small x behaviour of the gluon distribution



4. Future Avenues : What would we like to know ?

HERA:

- Collect high luminosity for $F_2(x, Q^2)$, $F_2^{c\bar{c}}(x, Q^2)$, $g_2^{c\bar{c}}(x, Q^2)$, and measure $h_1(x, Q^2)$.
- Measure : $F_L(x, Q^2)$. This is a key-question for HERA.

RHIC & LHC:

- Improve constraints on gluon and sea-quarks: polarized and unpolarized. DIS PDF's \Longleftrightarrow Collider PDF's


JLAB:

- High precision measurements in the large x domain at unpolarized and polarized targets; supplements HERA's high precision measurements at small x .

L_q from DVCS

HERA and JLAB : Improve DVCS data

Theory widely developed, cf. rev. Belitsky & Radyushkin, 2005



Expected DVCS asymmetry $A_{UT}^{\sin(\phi-\phi_S)\cos\phi}$ with $b_v = 1, b_s = \infty, J_u = 0.4(0.2, 0.0), J_d = 0.0$ in the Regge (left panel) and factorized (right panel) ansatz, at the average kinematics of the full measurement. $E = 0$ denotes zero effective contribution from the GPD E. The projected statistical error for 8M DIS events is shown. The systematic error is expected to not exceed the statistical one.

F. Ellinghaus et al. 2005

The measurement of L_q off data is model-dependent at the moment.

Lattice calculations at low pion masses are needed to complete the picture

Graph Resummation and Saturation

Further study of proposed mechanisms needed: RHIC, LHC
for nucleus-nucleus collisions.

ep scattering: partly different mechanisms

more studies would be welcome; link to higher twist contributions
in gluon-dynamics


How do the non-perturbative and perturbative parts factorize ?

Conservation laws and interplay between the small x and
medium x range behaviour

New DIS Machines

Where to go ?

- High energies : small x , large Q^2 desirable.
- High luminosities : ELIC: \sqrt{s} between CERN and HERA energies



R. Ent, 2004
high precision physics
polarized and unpolarized

Would be an important extension of the
present programmes in many respects.

Enhancing Precision Further...

- What is the correct value of $\alpha_s(M_z^2)$? $\overline{\text{MS}}$ -analysis vs. scheme-invariant evolution helps. Compare non-singlet and singlet analysis; careful treatment of heavy flavor. (Theory & Experiment)
- Flavor Structure of Sea-Quarks: More studies needed. (All Experiments)
- Revisit polarized data upon arrival of the 3-loop anomalous dimensions; NLO heavy flavor contributions needed. (Theory)
- QCD at Twist 3: $g_2(x, Q^2)$, semi-exclusive Reactions, Transversity, diffraction in polarized scattering (HERMES, High Precision polarized experiments, JLAB, ELIC)
- Comparison with Lattice Results: α_s , Moments of Parton Distributions, Angular Momentum.

Enhancing Precision Further...

- Calculation of more hard scattering reactions at the 3-loop level: LHC
- Further perfection of the mathematical tools:
⇒ Algorithmic simplification of Perturbation theory in higher orders.
- Even higher order corrections needed ?