

The CS Kernel for dynamical z_{max}

March 23, 2023

The CS Kernel

The CSS factorisation formula:

$$\frac{d\sigma}{dQ^2 dy dq_T^2} = \frac{2\pi}{9} \frac{\alpha_{\text{em}}^2(Q)}{sQ^2} |C_V(Q, \mu_Q)|^2 \int_0^\infty db b J_0(bq_T) \\ \times \sum_q e_q^2 f_{q,h_1}(x_1, b; \mu_Q, Q^2) f_{\bar{q},h_2}(x_2, b; \mu_Q, Q^2), \quad (1)$$

The CS Kernel is defined as

$$\frac{df_{q,h}(x, b; \mu, \zeta)}{d \ln \zeta} = -\mathcal{D}(b, \mu) f_{q,h}(x, b; \mu, \zeta)$$

The CS Kernel from ratios

$$\mathcal{D}(b, \mu_0) = \frac{\ln \left(\frac{\Sigma_1}{\Sigma_2} \right) - \ln Z(Q_1, Q_2) - 2\Delta_R(Q_1, Q_2; \mu_0)}{4 \ln(Q_2/Q_1)} - 1$$

where Σ is the Hankel transformation of the cross-section:



$$\Sigma(s, y, Q, b) = \int_0^\infty dq_T q_T J_0(q_T b) \frac{d\sigma}{dQ^2 dy dq_T^2}.$$

and

$$\Delta_R(Q_1, Q_2; \mu_0) = \int_{\mu_{Q_2}}^{\mu_{Q_1}} \frac{d\mu}{\mu} \gamma_F(\mu, Q_1) - 2 \ln \left(\frac{Q_1}{Q_2} \right) \int_{\mu_0}^{\mu_{Q_2}} \frac{d\mu}{\mu} \Gamma_{\text{cusp}}(\mu) \quad Z(Q_1, Q_2) = \frac{\alpha_{\text{em}}^2(Q_1) |C_V(Q_1, \mu_{Q_1})|^2}{\alpha_{\text{em}}^2(Q_2) |C_V(Q_2, \mu_{Q_2})|^2}$$


The DY cross section: PBSet2 vs Dynamical z_{max}

Dynamical z_{max} with $q_0 = 1$ GeV and $q_s = 0.5$ GeV




The CS kernel in CASCADE: PBSet2 vs Dynamical z_{max}

Dynamical z_{max} with $q_0 = 1$ GeV and $q_s = 0.5$ GeV



The CS kernel in CASCADE: Flat behaviour at large b

Is the intrinsic k_T distribution?



The End