

Freeze out Freeze in and the Flavon

One model working at two scales

Dedicated to 2023



Theory Workshop



Tom Tong

arXiv: 2307.14972

Co-directed by Rusa Mandal

50th ANNIVERSARY EDITION

CLINT EASTWOOD



THE GOOD THE BAD and THE UGLY


co-starring
LEE VAN CLEEF

also starring
ELI WALLACH
in the role of TUCO



directed by
SERGIO LEONE

Evidence of new physics

Evidence of new physics






Evidence of new physics





Evidence of new physics

Mystery of flavor




Evidence of new physics

Mystery of flavor






- SM flavor structure
- 13 parameters
- 20 with massive neutrinos



Evidence of new physics

Mystery of flavor



Summary for the impatient




"THAT'S THE END OF MY PRESENTATION. ANY QUESTIONS?"

Summary for the impatient



"THAT'S THE END OF MY PRESENTATION. ANY QUESTIONS?"




Summary for the impatient



"THAT'S THE END OF MY PRESENTATION. ANY QUESTIONS?"

Thanks for watching !




- ✓ • Freeze-out WIMP
- ✓ • Freeze-in FIMP

Hierarchy in the fermion mass

$$\frac{M_e}{M_t} \approx 3 \times 10^{-6}$$



Top Yukawa $y_t \approx 1$



Hierarchy in the fermion mass

$$\frac{m_e}{M_t} \approx 3 \times 10^{-6}$$



Top Yukawa $y_t \approx 1$




Hierarchy in the fermion mass

$$\frac{m_e}{M_t} \approx 3 \times 10^{-6}$$


Top Yukawa $y_t \approx 1$




Global U(1) Froggatt-Nielsen symmetry





$$y \bar{f}_L H f_R$$





Global U(1) Froggatt-Nielsen symmetry



Global U(1) Froggatt-Nielsen symmetry



Global U(1) Froggatt-Nielsen symmetry




$$\left\{ \begin{array}{l} c \sim 0(1) \\ \frac{\phi \rightarrow v_\phi}{M} = \varepsilon \\ q_L - q_R = n \end{array} \right.$$

Froggatt-Nielsen charges

$$\xi = \frac{V_\phi}{M} \approx V_{us} \approx 0.23$$



CKM




$$\left\{ \begin{array}{l} C \sim O(1) \\ \frac{\phi \rightarrow V_\phi}{M} = \xi \\ g_L - g_R = n \end{array} \right.$$


Froggatt-Nielsen charges

$$\xi = \frac{V_\phi}{M} \approx V_{us} \approx 0.23$$

$$n_u^{ij} = \begin{pmatrix} 8 & 4 & 3 \\ 7 & 3 & 2 \\ 5 & 1 & 0 \end{pmatrix}, \quad n_d^{ij} = \begin{pmatrix} 7 & 6 & 6 \\ 6 & 5 & 5 \\ 4 & 3 & 3 \end{pmatrix}, \quad n_e^{ij} = \begin{pmatrix} 9 & 6 & 4 \\ 8 & 5 & 3 \\ 8 & 5 & 3 \end{pmatrix}$$



CKM




$$\left\{ \begin{array}{l} c \sim 0(1) \\ \frac{\phi \rightarrow V_\phi}{M} = \xi \\ g_L - g_R = n \end{array} \right.$$


Froggatt-Nielsen charges

$$\xi = \frac{V_\phi}{M} \approx V_{us} \approx 0.23$$

$$n_u^{ij} = \begin{pmatrix} 8 & 4 & 3 \\ 7 & 3 & 2 \\ 5 & 1 & 0 \end{pmatrix}, \quad n_d^{ij} = \begin{pmatrix} 7 & 6 & 6 \\ 6 & 5 & 5 \\ 4 & 3 & 3 \end{pmatrix}, \quad n_e^{ij} = \begin{pmatrix} 9 & 6 & 4 \\ 8 & 5 & 3 \\ 8 & 5 & 3 \end{pmatrix}$$



CKM



$$q_L = \begin{pmatrix} q_{Q_1} & q_{Q_2} & q_{Q_3} \\ q_u & q_c & q_t \\ q_d & q_s & q_b \end{pmatrix} = \begin{pmatrix} 3 & 2 & 0 \\ -5 & -1 & 0 \\ -4 & -3 & -3 \end{pmatrix}$$


$$q_R = \begin{pmatrix} q_{L_1} & q_{L_2} & q_{L_3} \\ q_e & q_\mu & q_\tau \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ -8 & -5 & -3 \end{pmatrix}$$

$$\left\{ \begin{array}{l} C \sim O(1) \\ \frac{\phi \rightarrow V_\phi}{M} = \xi \\ q_L - q_R = n \end{array} \right.$$


Froggatt-Nielsen charges

$$\Sigma = \frac{V_\phi}{M} \approx V_{us} \approx 0.23$$

$$n_u^{ij} = \begin{pmatrix} 8 & 4 & 3 \\ 7 & 3 & 2 \\ 5 & 1 & 0 \end{pmatrix}, \quad n_d^{ij} = \begin{pmatrix} 7 & 6 & 6 \\ 6 & 5 & 5 \\ 4 & 3 & 3 \end{pmatrix}, \quad n_e^{ij} = \begin{pmatrix} 9 & 6 & 4 \\ 8 & 5 & 3 \\ 8 & 5 & 3 \end{pmatrix}$$




CKM



$$\begin{array}{c}
 \left(\begin{array}{ccc} q_{Q_1} & q_{Q_2} & q_{Q_3} \\ q_u & q_c & q_t \\ q_d & q_s & q_b \end{array} \right) = \left(\begin{array}{ccc} 3 & 2 & 0 \\ -5 & -1 & 0 \\ -4 & -3 & -3 \end{array} \right) \\
 q_L \quad \quad \quad q_R \\
 \left(\begin{array}{ccc} q_{L_1} & q_{L_2} & q_{L_3} \\ q_e & q_\mu & q_\tau \end{array} \right) = \left(\begin{array}{ccc} 1 & 0 & 0 \\ -8 & -5 & -3 \end{array} \right)
 \end{array}$$

$$\left\{ \begin{array}{l}
 C \sim O(1) \\
 \frac{\phi \rightarrow V_\phi}{M} = \Sigma \\
 q_L - q_R = n
 \end{array} \right.$$




Right-handed neutrino and DM

$$\left\{ N_1 \quad N_2 \quad N_3 \right\}$$

Heavy

A hand-drawn diagram showing three elements labeled N_1 , N_2 , and N_3 enclosed in a black brace. Below the brace, a green checkmark is drawn above the word "Heavy".




Right-handed neutrino and DM


$\left\{ N_1 \quad N_2 \quad N_3 \right\}$

Heavy


$C_\nu \left(\frac{\phi}{M} \right)^{n_\nu} \bar{L} \tilde{H} N_{2,3}$

Yukawa: 3×2 matrix





Right-handed neutrino and DM




$$C_\nu \left(\frac{\phi}{M} \right)^{n_\nu} \bar{L} \tilde{H} N_{2,3}$$

Yukawa: 3×2 matrix


X	X
X	X
X	X

DM





Right-handed neutrino and DM



$$C_\nu \left(\frac{\phi}{M} \right)^{n_\nu} \bar{L} \tilde{H} N_{2,3}$$

Yukawa: 3×2 matrix

X	X
X	X
X	X

$$\frac{1}{2} C_N \left(\frac{\phi}{M} \right)^{n_N} M \bar{N} N$$

Majorana: 3×3 matrix

X	O	O
O	X	X
O	X	X

Flavon couplings

$$H \rightarrow V_{EW} + \frac{1}{\sqrt{2}}(h + iG_0)$$

Flavon couplings

$$H \rightarrow V_{EW} + \frac{1}{\sqrt{2}}(h + iG_0)$$

$$\phi \rightarrow V_\phi + \frac{1}{\sqrt{2}}(S + iA)$$

Flavon couplings

$$H \rightarrow V_{EW} + \frac{1}{\sqrt{2}}(h + iG_0)$$

$$\phi \rightarrow V_\phi + \frac{1}{\sqrt{2}}(S + iA)$$


$$\left. \begin{array}{l} M_S \sim V_\phi \\ 10 \text{ GeV} \leq M_A \ll M_S \end{array} \right\}$$

Flavon couplings

$$H \rightarrow V_{EW} + \frac{1}{\sqrt{2}}(h + iG_0)$$

$$\phi \rightarrow V_\phi + \frac{1}{\sqrt{2}}(S + iA)$$

$$\left. \begin{array}{l} M_S \sim V_\phi \\ 10 \text{ GeV} \leq M_A \ll M_S \end{array} \right\}$$




Flavon couplings

$$H \rightarrow V_{EW} + \frac{1}{\sqrt{2}}(h + iG_0)$$

$$\phi \rightarrow V_\phi + \frac{1}{\sqrt{2}}(S + iA)$$

$$\left. \begin{array}{l} M_S \sim V_\phi \\ 10 \text{ GeV} \leq M_A \ll M_S \end{array} \right\}$$




Flavon couplings

$$H \rightarrow V_{EW} + \frac{1}{\sqrt{2}}(h + iG_0)$$

$$\phi \rightarrow V_\phi + \frac{1}{\sqrt{2}}(S + iA)$$

$$\left. \begin{aligned} M_S &\sim V_\phi \\ 10 \text{ GeV} &\leq M_A \leq M_S \end{aligned} \right\}$$






Flavon couplings

$$H \rightarrow V_{EW} + \frac{1}{\sqrt{2}}(h + iG_0)$$

$$\phi \rightarrow V_\phi + \frac{1}{\sqrt{2}}(S + iA)$$

$$\left. \begin{array}{l} M_S \sim V_\phi \\ 10 \text{ GeV} \leq M_A \leq M_S \end{array} \right\}$$






Flavon couplings

$$H \rightarrow V_{EW} + \frac{1}{\sqrt{2}}(h + iG_0)$$


$$\phi \rightarrow V_\phi + \frac{1}{\sqrt{2}}(S + iA)$$

$$\left. \begin{array}{l} M_S \sim V_\phi \\ 10 \text{ GeV} \leq M_A \leq M_S \end{array} \right\}$$




Experimental constraints

Experimental constraints



Experimental constraints


B-meson mixing



$$\left(\frac{M_\alpha}{\text{GeV}}\right) \times \left(\frac{V_\phi}{\text{TeV}}\right) \geq 180$$



Experimental constraints

B -meson mixing




$$\left(\frac{M_\alpha}{\text{GeV}}\right) \times \left(\frac{V_\phi}{\text{TeV}}\right) \geq 180$$

$\mu \rightarrow e \tau$ (MEG)




Experimental constraints

B -meson mixing



$$\left(\frac{M_\alpha}{\text{GeV}}\right) \times \left(\frac{V_\phi}{\text{TeV}}\right) \geq 180$$

$u \rightarrow e \tau$ (MEG)




$$\left(\frac{M_\alpha}{\text{GeV}}\right) \times \left(\frac{V_\phi}{\text{TeV}}\right) \geq 190^*$$

* For $M_\alpha \lesssim 100 \text{ GeV}$


Experimental constraints

B -meson mixing



$$\left(\frac{M_\alpha}{\text{GeV}}\right) \times \left(\frac{V_\phi}{\text{TeV}}\right) \geq 180$$

$u \rightarrow e \tau$ (MEG)



$$\left(\frac{M_\alpha}{\text{GeV}}\right) \times \left(\frac{V_\phi}{\text{TeV}}\right) \geq 190^*$$


*For $M_\alpha \lesssim 100 \text{ GeV}$

Top decay
ATLAS

$$\begin{aligned} &\text{Br}(t \rightarrow c a) \\ &\times \text{Br}(a \rightarrow b \bar{b}) \end{aligned}$$


Experimental constraints

B -meson mixing



$$\left(\frac{M_\alpha}{\text{GeV}}\right) \times \left(\frac{V_\phi}{\text{TeV}}\right) \geq 180$$

$u \rightarrow e \gamma$ (MEG)




$$\left(\frac{M_\alpha}{\text{GeV}}\right) \times \left(\frac{V_\phi}{\text{TeV}}\right) \geq 190^*$$

* For $M_\alpha \lesssim 100 \text{ GeV}$

Top decay
ATLAS

$$\begin{aligned} &\text{Br}(t \rightarrow c a) \\ &\times \text{Br}(a \rightarrow b \bar{b}) \end{aligned}$$



EFT... at what scale? $V_\phi = ?$

EFT... at what scale? $V_\phi = ?$


Freeze out
Freeze in
and the Flavon

One model working at two scales

TeV scale: freeze out

m_a vs V_ϕ

$$\Omega h^2 = 0.12$$




TeV scale: freeze out

$$\left(\frac{m_a}{\text{GeV}}\right) \times \left(\frac{V_\phi}{\text{TeV}}\right) \geq \begin{cases} 180 & B_s \\ 190^* & u \rightarrow e\gamma \end{cases}$$

(A horizontal orange line is drawn at approximately 600 GeV.)

m_a vs V_ϕ



$$\Omega h^2 = 0.12$$




TeV scale: freeze out

m_a vs V_ϕ

$$\Omega h^2 = 0.12$$




TeV scale: freeze out




m_a vs V_ϕ

$$\Omega h^2 = 0.12$$





TeV scale: freeze out

m_{DM} vs V_ϕ





TeV scale: freeze out

m_{DM} vs V_ϕ




TeV scale: freeze out

m_{DM} vs V_ϕ




TeV scale: freeze out with various λ_ϕ




TeV scale: freeze out with various λ_ϕ


$\lambda\phi^4$ term in
flavon potential




TeV scale: freeze out with various λ_ϕ





$\lambda\phi^4$ term in
flavon potential



TeV scale: freeze out with various λ_ϕ




$\lambda\phi^4$ term in
flavon potential



High scale: (IR) freeze in m_{DM} vs V_ϕ

α



High scale: (IR) freeze in

m_{DM} vs V_ϕ



High scale: (IR) freeze in

m_{DM} vs V_ϕ




High scale: (IR) freeze in

m_{DM} vs V_ϕ




"Low" T_{RH}





High scale: (UV) freeze in


m_{DM} vs V_ϕ




High scale: (UV) freeze in m_{DM} vs V_ϕ



Direct detection



Direct detection





Spin-dependent σ

Momentum suppressed

10^{-5} too small !

Direct detection




Spin-dependent σ

Momentum suppressed

10^{-5} too small !



Direct detection




Spin-dependent σ

Momentum suppressed

10^{-5} too small !





Direct detection




Spin-dependent σ

Momentum suppressed




10^{-5} too small !







Spin-independent σ



Summary for the impatient







Summary for the impatient




- ✓ • Freeze-out WIMP
- ✓ • Freeze-in FIMP

Summary for the impatient



Next : Baryogenesis



- ✓ • Freeze-out WIMP
- ✓ • Freeze-in FIMP