

Contribution ID: 26

Type: Poster

Sphaleron Rate from Lattice Gauge Theory

The "Sphaleron Rate" (imaginary linear-in-frequency part of the topological density retarded Green's function) determines the real-time relaxation rate of axial quark number for light quarks in a hot medium, and is relevant in heavy-ion collisions and electroweak baryogenesis. We recently showed how it can be determined in pureglue QCD via standard Euclidean simulations, via a novel saddlepoint method. We extend this work to find the sphaleron rate for (2+1)-flavor QCD with $N_{\tau} = 8 - 16$ and HISQ action at almost physical pion masses in the temperature range 0.2 - 3 GeV or 1.2 - 18 times the crossover temperature T_{pc} .

Similar to the pure gauge case, the QCD result is well described across the range of 1.6 - 8 times T_{pc} as $\Gamma_s \simeq 20(\alpha_s T)^4$, where α_s is the MSbar coupling at $\overline{\mu} = 2\pi T$, determined using the gradient-flow technique.

Primary authors: Prof. MOORE, Guy (TU Darmstadt); WEBER, Johannes Heinrich (Humboldt University of Berlin); Mr BARROSO MANCHA, Marc (TU Darmstadt)

Presenter: WEBER, Johannes Heinrich (Humboldt University of Berlin)