

LCFIPlus variables



Name	Description	Normalization factor	Used by category
trk1d0sig	d0 significance of track with highest d0 significance	1	A, B, C, D
trk2d0sig	d0 significance of track with second highest d0 significance	1	A, B, C, D
trk1z0sig	z0 significance of track with highest d0 significance	1	A, B, C, D
trk2z0sig	z0 significance of track with second highest d0 significance	1	A, B, C, D
trk1pt	transverse momentum of track with highest d0 significance	$1/E_{\text{jet}}$	A, B, C, D
trk2pt	transverse momentum of track with second highest d0 significance	$1/E_{\text{jet}}$	A, B, C, D
jprobr	joint probability in the r-phi plane using all tracks	1	A, B, C, D
jprobr5sigma	joint probability in the r-phi plane using all tracks having impact parameter significance exceeding 5 sigma	1	A, B, C, D
jprobz	joint probability in the z projection using all tracks	1	A, B, C, D
jprobz5sigma	joint probability in the z projection using all tracks having impact parameter significance exceeding 5 sigma	1	A, B, C, D
d0bprob	product of b-quark probabilities of d0 values for all tracks, using b/c/q d0 distributions	1	A, B, C, D
d0cprob	product of c-quark probabilities of d0 values for all tracks, using b/c/q d0 distributions	1	A, B, C, D
d0qprob	product of q-quark probabilities of d0 values for all tracks, using b/c/q d0 distributions	1	A, B, C, D
z0bprob	product of b-quark probabilities of z0 values for all tracks, using b/c/q z0 distributions	1	A, B, C, D
z0cprob	product of c-quark probabilities of z0 values for all tracks, using b/c/q z0 distributions	1	A, B, C, D
z0qprob	product of q-quark probabilities of z0 values for all tracks, using b/c/q z0 distributions	1	A, B, C, D
nmuon	number of identified muons	1	A, B, C, D
nelectron	number of identified electrons	1	A, B, C, D
trkmass	mass of all tracks exceeding 5 sigma significance in d0/z0 values	1	A, B, C, D

used as global variables

LCFIPlus variables

Name	Description	Normalization factor	Used by category
1vtxprob	vertex probability with all tracks associated in vertices combined	1	B, C, D
vtxlen1	decay length of the first vertex in the jet (zero if no vertex is found)	$1/E_{\text{jet}}$	B, C, D
vtxlen2	decay length of the second vertex in the jet (zero if number of vertex is less than two)	$1/E_{\text{jet}}$	D
vtxlen12	distance between the first and second vertex (zero if number of vertex is less than two)	$1/E_{\text{jet}}$	D
vtxsig1	decay length significance of the first vertex in the jet (zero if no vertex is found)	$1/E_{\text{jet}}$	B, C, D
vtxsig2	decay length significance of the second vertex in the jet (zero if number of vertex is less than two)	$1/E_{\text{jet}}$	D
vtxsig12	vtxlen12 divided by its error as computed from the sum of the covariance matrix of the first and second vertices, projected along the line connecting the two vertices	$1/E_{\text{jet}}$	D
vtxdirang1	the angle between the momentum (computed as a vector sum of track momenta) and the displacement of the first vertex	E_{jet}	B, C, D
vtxdirang2	the angle between the momentum (computed as a vector sum of track momenta) and the displacement of the second vertex	E_{jet}	D
vtxmult1	number of tracks included in the first vertex (zero if no vertex is found)	1	B, C, D
vtxmult2	number of tracks included in the second vertex (zero if number of vertex is less than two)	1	D
vtxmult	number of tracks which are used to form secondary vertices (summed for all vertices)	1	D
vtxmom1	magnitude of the vector sum of the momenta of all tracks combined into the first vertex	$1/E_{\text{jet}}$	B, C, D
vtxmom2	magnitude of the vector sum of the momenta of all tracks combined into the second vertex	$1/E_{\text{jet}}$	D
vtxmass1	mass of the first vertex computed from the sum of track four-momenta	1	B, C, D
vtxmass2	mass of the second vertex computed from the sum of track four-momenta	1	D
vtxmass	vertex mass as computed from the sum of four momenta of all tracks forming secondary vertices	1	B, C, D
vtxmasspc	mass of the vertex with minimum pt correction allowed by the error matrices of the primary and secondary vertices	1	B, C, D
vtxprob	vertex probability; for multiple vertices, the probability P is computed as $1-P = (1-P_1)(1-P_2)\dots(1-P_N)$	1	B, C, D



Results: loss & accuracy



- overtraining
- accuracy ~79% in validation data (before ~76%)

Results: confusion matrices



validation data



- better identification efficiencies for c jets and light jets (66% & 85% before)
- improved purity in all classes (before 88% for b jets, 68% for c jets, 74% for light jets)
- especially separation between c jets and light jets improved



ROC curves

validation data



ROC curves - comparison to LCFIPlus

validation data





Next steps

- implement the new variables into my Marlin processor
- use new variables as features of charged jet constituents (per track and not combined for all tracks)?
- add more variables of LCFIPlus?
- start to integrate into iLCSoft, make it usable for others
 - meeting with Frank, Thomas & Uli
- testing over-sampling, other activation functions, ...

Backup

Results: confusion matrices



validation data



- identification efficiencies ~80% for b jets & light jets
- c jet identification quite low (66%)
- especially separation between c jets and light jets should be improved



ROC curves - comparison to LCFIPlus

validation data





- slightly better performance for b jet identification vs. c jet background
- better performance for c jet efficiencies vs. b jet background below ~90% c jet identification efficiency
- worse performance in b jet / c jet identification vs. light jets (especially for c jets)



LCFIPlus variables





LCFIPlus variables





LCFIPlus variables




LCFIPlus variables




LCFIPlus variables




LCFIPlus variables




LCFIPlus variables




LCFIPlus variables




LCFIPlus variables




LCFIPlus variables




LCFIPlus variables




LCFIPlus variables




LCFIPlus variables




LCFIPlus variables




LCFIPlus variables






LCFIPlus variables






hb




Performance LCFIPlus



Performance LCFIPlus



Architecture & data pre-processing





- classify jets into **three classes**: b jets, c jets & light jets
- **ordering of input particles** by (as applied in CMS)
 - impact parameter significance for charged jet constituents
 - shortest angular distance to a secondary vertex (by momentum if there is no secondary vertex) for neutral jet constituents
 - flight distance significance for secondary vertices
- if a value of a features is not available, the value is set to -10
- **normalize input features** to mean 0, std 1

Input features - global variables

- jet momentum
- jet transverse momentum
- number of charged jet constituents
- number of neutral jet constituents
- number of secondary vertices



5 input features



Input features - charged jet constituents

- track momentum / jet momentum
- transverse track momentum relative to jet
- dot product of jet and track momentum w.r.t. jet momentum
- $\Delta R(\text{track}, \text{jet})$,
- d_0 , d_0 significance
- Z_0 , Z_0 significance
- 3D impact parameter, 3D impact parameter significance
- track reconstructed in PV?
- is electron?, is muon?, lepton momentum relative to jet, lepton transverse momentum relative to the jet, lepton momentum / jet momentum
- kaon-ness of charged particles, track momentum fraction weighted with kaon-ness
- χ^2/ndf



19 input features



Input features - neutral jet constituents

- momentum of neutral jet constituent
- fraction of the jet momentum carried by neutral jet constituent
- $\Delta R(\text{jet axis, neutral candidate})$,
- is photon?
- fraction of neutral candidate energy deposited in the hadronic calorimeter



5 input features



Input features - secondary vertices



- SV mass
- number of tracks in SV
- $\Delta R(\text{SV}, \text{jet})$
- SV energy / jet energy
- SV energy
- cosine of the angle between the secondary vertex flight direction and the direction of the secondary vertex momentum
- 3D impact parameter, 3D impact parameter significance
- χ^2 , ndf

10 input features





Training

- activation functions: relu / softmax (last layer)
- cross entropy loss
- optimizer: Adam
- regularization: batch normalization, dropout (0.1)
- batch size: 200
- learning rate: 0.0003
- learning rate is halved if validation loss stagnates for 10 epochs
- number of epochs: 100
- Xavier weight initialization



Results: loss & accuracy



- accuracy ~76% in training & validation data
- epoch 89: learning rate halved
- train longer? decrease dropout rate?