

Analysing New Samples

Leo Rozanov



KDP Lab | University of Chicago | Muon Colliders



Muon Gun Data Selection


- Data from

/collab/project/snowmass21/data/muonc/fmeloni/Data


MuC_MuColl10_v0A/reco/muonGun_pT_0_50,

pt_50_250, and pt_250_1000

- (note: no BIB)
- I matched the reconstructed Muon tracks to truth Muons using only dR matching with $dR < 0.1$
- No requirement on muon pT, nhits, etc.





Distribution Comparison








d0 Bulk vs Tails using 5σ cutoff (pt: 0-50)

- Mean from Fit: -7.573352379753462e-05
- Sigma from Fit: 0.004501198659859312








p_T Bulk vs Tails using 5σ cutoff (pt: 0-50)

- Mean from Fit: 0.28309599682693337
- Sigma from Fit: 0.002726736622490658
- (NB: off-centre resolution, not sure why)



d0 Bulk vs Tails using 5σ cutoff (pt: 50-250)






- Mean from Fit: 0.0003061676728036983
- Sigma from Fit: 0.0029933548097349676



p_T Bulk vs Tails using 5σ cutoff (pt: 50-250)






- Mean from Fit: 0.2854270814443211
- Sigma from Fit: 0.00705020790980997
- (NB: off-centre resolution, not sure why)

p_T resolution vs p_T for Slice 0



d0 Bulk vs Tails using 5σ cutoff (pt: 250-1000)

- Mean from Fit: 0.0003061676728036983
- Sigma from Fit: 0.0029933548097349676



p_T Bulk vs Tails using 5σ cutoff (pt: 250-1000)

- Mean from Fit: 0.2854270814443211
- Sigma from Fit: 0.00705020790980997
- (NB: off-centre resolution, not sure why)

