

Hybrid static potentials from Laplacian eigenmodes

R. Höllwieser, T. Korzec, F. Knechtli,

M. Peardon, L. Struckmeier, J.-A. Urrea-Niño



University of Wuppertal, Trinity College Dublin



FOR 5269



Lattice QCD



- ▶ link variables $U_\mu(x) = \exp(i \int_x^{x+a\hat{\mu}} A_\mu dx^\mu)$
- ▶ Wilson line $U_s(\vec{x}, \vec{y}) = \exp(i \int_{\vec{x}}^{\vec{y}} A_\mu dx^\mu) = \prod U_\mu$
- ▶ path-ordered product of link variables, on-/off-axis
- ▶ plaquette, Wilson loop $W(R, T)$, static $\bar{Q}Q$ pair



Motivation

- ▶ hybrid mesons (valence glue, exotics, XYZ), (hybrid) static-light
- ▶ identify bound states using Born-Oppenheimer approximation
Braaten et al. (2014), Capitani et al. (2019)
- ▶ observation of (hybrid) string breaking in QCD, e.g., Bali et al. (2008), Bulava et al. (2019)
- ▶ calculate the (hybrid) static potential with high resolution
- ⇒ on the lattice we have to work with off-axis separated quarks
- ▶ the spatial part of the Wilson loop has to go over stair-like paths through the lattice → not unique, computationally expensive
- ⇒ alternative operator which ensures gauge invariance of the quark-anti-quark $\bar{Q}(\vec{x})U_s(\vec{x}, \vec{y})Q(\vec{y})$ trial state
- ▶ required gauge transformation behavior:

$$U'_s(\vec{x}, \vec{y}) = G(\vec{x})U_s(\vec{x}, \vec{y})G^\dagger(\vec{y})$$



Laplace trial states


- idea taken from [Neitzel et al. \(2016\)](#) SU(2)
- eigenvectors $v(\vec{x})$ of the 3D covariant lattice Laplace operator
- spatial Wilson line: $U'_s(\vec{x}, \vec{y}) = G(\vec{x})U_s(\vec{x}, \vec{y})G^\dagger(\vec{y})$

$$v'(\vec{x})v'^\dagger(\vec{y}) = G(\vec{x})v(\vec{x})v^\dagger(\vec{y})G^\dagger(\vec{y})$$

- Wilson loop of size $(R = |\vec{r}| = |\vec{x} - \vec{y}|) \times (T = |t_1 - t_0|)$

$$W(R, T) = \langle \text{tr} [U_t(\vec{x}; t_0, t_1) U_s(\vec{x}, \vec{y}; t_1) U_t^\dagger(\vec{y}; t_0, t_1) U_s^\dagger(\vec{x}, \vec{y}; t_0)] \rangle$$



$$\rightarrow \langle \sum_{i,j}^{N_v} \text{tr} [U_t(\vec{x}) \rho_j(t_1) v_j(\vec{x}, t_1) v_j^\dagger(\vec{y}, t_1) U_t^\dagger(\vec{y}) \rho_i(t_0) v_i(\vec{y}, t_0) v_i^\dagger(\vec{x}, t_0)] \rangle$$



Static Perambulator

static perambulator $\tau_{ij}(\vec{x}, t_0, t_1) = v_i^\dagger(\vec{x}, t_0) U_t(\vec{x}; t_0, t_1) v_j(\vec{x}, t_1)$

$$Q(t) = \sum_{\vec{x}, t_0} \langle \sum_{i,j}^{N_v} \tau_{ij}(\vec{x}, t_0, t_0 + t) \rangle$$





expectation value should vanish, no single quarks...



Optimal trial state results

- ▶ Gaussian profile functions: $\rho_i^{(k)}(\lambda_i) = \exp(-\lambda_i^2/2\sigma_k^2)$
- ▶ define correlation matrix W_{kl} using 7 different $\sigma_{k,l}$, SVD ($u_{k,l}$)
- ▶ GEVP: $W(t)\nu^{(n)} = \mu^{(n)}W(t_0)\nu^{(n)}$, $\mu^{(n)}$ give effective energies
- ▶ optimal profiles $\tilde{\rho}_R^{(n)}(\lambda_i) = \nu_k^{(n)} u_{k,l} \exp(-\lambda_i^2/2\sigma_l^2)$
- ▶ $24^3 \times 48$, $\beta = 5.3$, $N_f = 2$, $\kappa = 0.13270$, $a = 0.0658$ fm



Static-hybrid potentials

Hybrid static potentials are characterized by the following quantum numbers Λ_η^ϵ , [Bali et al. \(2005\)](#), [Bicudo et al. \(2015\)](#):

- ▶ $\Lambda = 0, 1, 2, 3, \dots \equiv \Sigma, \Pi, \Delta, \Phi, \dots$, the absolute value of the total angular momentum with respect to the axis of separation of the static quark-antiquark pair
- ▶ $\eta = +, - \equiv g, u$, the eigenvalue corresponding to the operator $\mathcal{P} \circ \mathcal{C}$, i.e. the combination of parity and charge conjugation
- ▶ $\epsilon = +, -$, the eigenvalue corresponding to the operator $\mathcal{P}_x$, which denotes the spatial reflection with respect to a plane including the axis of separation ($\Lambda > 0$ degenerate)

⇒ derived from the continuous group $D_{\infty h}$, which leaves a cylinder along a chosen axis invariant, with irreducible representations (irreps): $A_1^\pm(\Sigma_\pm^+)$, $A_2^\pm(\Sigma_\pm^-)$, $E_1^\pm(\Pi_\pm)$, $E_2^\pm(\Delta_\pm)$, $E_3^\pm(\Phi_\pm), \dots$


⇒ on the lattice we have D_{4h} , with 10 irreps: $A_1^\pm$, $A_2^\pm$, $B_1^\pm$, $B_2^\pm$, $E^\pm$

⇒ subduction relations $A_{1,2}^\pm \rightarrow A_{1,2}^\pm$, $E_1^\pm \rightarrow E^\pm$, $E_2^\pm \rightarrow B_1^\pm \oplus B_2^\pm, \dots$



Wilson loop static-hybrid states

standard construction with handles, e.g. Σ_g^- , [Capitani et al. \(2019\)](#)



Laplacian static-hybrid states

- ▶ we can realize gluonic excitations via covariant derivatives

$$\nabla_{\vec{k}} v(\vec{x}) = \frac{1}{2} [U_k(\vec{x}) v(\vec{x} + \hat{k}) - U_k^\dagger(\vec{x} - \hat{k}) v(\vec{x} - \hat{k})]$$
- ▶ derivative based operators that transform according to the 10 irreps of the cubic group O_h : $\nabla_i(T_1)$, $\mathbb{B}_i = \epsilon_{ijk} \nabla_j \nabla_k(T_1)$, $\mathbb{D}_i = |\epsilon_{ijk}| \nabla_j \nabla_k(T_2)$, $\mathbb{E}_i = \mathbb{Q}_{ijk} \nabla_j \nabla_k(E)$, $\nabla^2(A_1)$
- ▶ since D_{4h} is a subgroup of O_h we have the subduction relations $A_1^\pm \rightarrow A_1^\pm$, $A_2^\pm \rightarrow B_1^\pm$, $E^\pm \rightarrow A_1^\pm \oplus B_1^\pm$, $T_1^\pm \rightarrow A_2^\pm \oplus E^\pm$ and $T_2^\pm \rightarrow B_2^\pm \oplus E^\pm$
- ▶ the three components of ∇_i get separated into one that transforms like A_2 (along the separation axis) and two that transform like E (the two orthogonal to the separation axis)
- ▶ with excited gluons also exotic quantum numbers accessible
 $J^{PC} = 0^{--}, 0^{+-}, 1^{-+}, 2^{+-}, 3^{-+}, 4^{+-}, 5^{-+}, \dots$
- ▶ in the pure quark model $P = (-1)^{L+1}$ and $C = (-1)^{L+S}$, with orbital angular momentum $L \in \{0, 1, 2, \dots\}$ and spin $S \in \{0, 1\}$



Laplacian static-hybrid states

$$\blacktriangleright \Sigma_g^+: v(-\vec{x})v^\dagger(\vec{x}) \quad \bullet \dots \bullet^\dagger \quad \leftrightarrow \dots \bullet^\dagger \mp \bullet \dots \leftrightarrow^\dagger$$

$$\blacktriangleright \Sigma_u^+: \nabla_{\vec{x}} v(-\vec{x})v^\dagger(\vec{x}) + v(-\vec{x})[\nabla_{\vec{x}} v(\vec{x})]^\dagger$$

$$\blacktriangleright \Pi_{\pm}^{\pm}: \uparrow \dots \bullet^\dagger \mp \bullet \dots \downarrow^\dagger \quad \bullet \cdot \uparrow^\dagger \bullet \dots \bullet^\dagger \mp \bullet \dots \bullet^\dagger \downarrow \cdot \bullet$$

$$\nabla_{\vec{k}} v(-\vec{x})v^\dagger(\vec{x}) \mp v(-\vec{x})[\nabla_{\vec{k}} v(\vec{x})]^\dagger \quad (\vec{k} \perp \vec{x})$$

$$v(-\vec{x})[\nabla_{\vec{k}} v(-\delta\vec{x})]^\dagger v(-\delta\vec{x})v^\dagger(\vec{x}) \mp v(-\vec{x})V^\dagger(\delta\vec{x})[\nabla_{\vec{k}} v(\delta\vec{x})]v^\dagger(\vec{x})$$

$$\oplus \dots \bullet^\dagger \pm \bullet \dots \oplus^\dagger \quad | \oplus | \dots \bullet^\dagger \pm \bullet \dots | \oplus |^\dagger$$

$$\mathbb{B}_{\vec{k}} v(-\vec{x})v^\dagger(\vec{x}) \pm v(-\vec{x})[\mathbb{B}_{\vec{k}} v(\vec{x})]^\dagger \quad (\vec{k} \perp \vec{x})$$

$$\mathbb{D}_{\vec{k}} v(-\vec{x})v^\dagger(\vec{x}) \pm v(-\vec{x})[\mathbb{D}_{\vec{k}} v(\vec{x})]^\dagger \quad (\vec{k} \perp \vec{x})$$

$$\blacktriangleright \Delta_{\pm}^{\pm}: \otimes \dots \bullet^\dagger \pm \bullet \dots \otimes^\dagger \quad | \otimes | \dots \bullet^\dagger \pm \bullet \dots | \otimes |^\dagger$$

$$\mathbb{B}_{\vec{x}} v(-\vec{x})v^\dagger(\vec{x}) \pm v(-\vec{x})[\mathbb{B}_{\vec{x}} v(\vec{x})]^\dagger$$

$$\mathbb{D}_{\vec{x}} v(-\vec{x})v^\dagger(\vec{x}) \pm v(-\vec{x})[\mathbb{D}_{\vec{x}} v(\vec{x})]^\dagger$$

$$\blacktriangleright \dots$$



Laplacian static-hybrid states

- $\Sigma_g^+(R, T) = \sum_{\vec{x}, t_0, i, j}$

$$\langle \text{tr} [U_t(\vec{x}; t_0, t_1) \rho(\lambda_j) v_j(\vec{x}, t_1) v_j^\dagger(\vec{y}, t_1) U_t^\dagger(\vec{y}; t_0, t_1) \rho(\lambda_i) v_i(\vec{y}, t_0) v_i^\dagger(\vec{x}, t_0)] \rangle$$
- $\Sigma_u^+(R, T) = \sum_{\vec{x}, t_0, i, j, \vec{k} || \vec{r}}$

$$\langle \text{tr} [U_t(\vec{x}; t_0, t_1) \rho(\lambda_j) \{ [\nabla_{\vec{k}} v_j](\vec{x}, t_1) v_j^\dagger(\vec{y}, t_1) + v_j(\vec{x}, t_1) [\nabla_{\vec{k}} v_j]^\dagger(\vec{y}, t_1) \} U_t^\dagger(\vec{y}; t_0, t_1) \rho(\lambda_i) \{ [\nabla_{\vec{k}} v_i](\vec{y}, t_0) v_i^\dagger(\vec{x}, t_0) + v_i(\vec{y}, t_0) [\nabla_{\vec{k}} v_i]^\dagger(\vec{x}, t_0) \}] \rangle$$
- $\Pi_{u/g}(R, T) = \Pi_{\mp}(R, T) = \sum_{\vec{x}, t_0, i, j, \vec{k} \perp \vec{r}}$

$$\langle \text{tr} [U_t(\vec{x}; t_0, t_1) \rho(\lambda_j) \{ [\nabla_{\vec{k}} v_j](\vec{x}, t_1) v_j^\dagger(\vec{y}, t_1) \pm v_j(\vec{x}, t_1) [\nabla_{\vec{k}} v_j]^\dagger(\vec{y}, t_1) \} U_t^\dagger(\vec{y}; t_0, t_1) \rho(\lambda_i) \{ [\nabla_{\vec{k}} v_i](\vec{y}, t_0) v_i^\dagger(\vec{x}, t_0) \pm v_i(\vec{y}, t_0) [\nabla_{\vec{k}} v_i]^\dagger(\vec{x}, t_0) \}] \rangle$$



Laplacian static-hybrid states

- check the relevant symmetries to characterize hybrid mesons listed above using the following operations, starting with a counter-clockwise 90° rotation $\mathcal{R}$ in the $\mu\nu$ -plane:

$$\mathcal{R}: \quad x'_\mu = -x_\nu \qquad x'_\nu = x_\mu$$

$$U_\mu(x) \rightarrow U_\nu(x') \qquad U_\nu(x) \rightarrow U_\mu^\dagger(x' - \hat{\mu})$$

$$\mathcal{C}: \quad U_k(x) \rightarrow U_k^*(x) \qquad v(x) \rightarrow v^*(x)$$


$$\mathcal{P}: \quad U_k(x_0, \vec{x}) \rightarrow U_k^\dagger(x_0, -\vec{x} - \hat{k}) \qquad v(x_0, \vec{x}) \rightarrow v(x_0, -\vec{x}),$$

$$\mathcal{P}_x: \quad \nabla_{\vec{k} \perp \vec{r}} v \rightarrow -\nabla_{\vec{k} \perp \vec{r}} v \qquad \nabla_{\vec{k} \parallel \vec{r}} v \rightarrow \nabla_{\vec{k} \parallel \vec{r}} v,$$





- for on-axis separations the potential in the continuum $\Pi_\mp$ repr. can be obtained from the $E_1^\mp$ representation of D_{4h}
- for off-axis separations we technically do not have D_{4h}
- we seem to be fine for off-axis separations in a 2D plane (rather than the 3d volume)
- on-axis only for derivatives along the separation axis



Laplacian static-hybrid states



Optimal trial state profiles



Born-Oppenheimer Approximation

- solve the radial Schrödinger equation: $E_{\Lambda_\eta^\epsilon;L,n} \Psi_{\Lambda_\eta^\epsilon;L,n}(r) =$

$$\left(-\frac{1}{2\mu} \frac{d^2}{dr^2} + \frac{L(L+1) - 2\Lambda^2 + J_{\Lambda_\eta^\epsilon}(J_{\Lambda_\eta^\epsilon} + 1)}{2\mu r^2} + V_{\Lambda_\eta^\epsilon}(r) \right) \Psi_{\Lambda_\eta^\epsilon;L,n}(r)$$

$\mu = m_Q m_{\bar{Q}} / (m_Q + m_{\bar{Q}})$, for $m_c = 1628$ MeV or $m_b = 4977$ MeV

- parametrization of potentials $V_{\Lambda_\eta^\epsilon}(r)$ from [Capitani et al. \(2019\)](#)

$$V_{\Sigma_g^+}(r) = V_0 - \alpha/r + \sigma r$$

$$V_{\Pi_u}(r) = A_1/r + A_2 + A_3 r^2$$

- $E_{\Lambda_\eta^\epsilon;L,n}$ contain the self-energies of the static quarks

$$\Rightarrow m_{\Lambda_\eta^\epsilon;L,n} = E_{\Lambda_\eta^\epsilon;L,n} - E_{\Sigma_g^+;L=0,n=1} + \bar{m}$$

with $\bar{m}$ the spin averaged mass [Workman et al. \[PDG\] \(2022\)](#)

$$\bar{m}_c = (m_{\eta_c(1S),\text{exp}} + 3m_{J/\Psi(1S),\text{exp}})/4 = 3069(1) \text{ MeV}$$

$$\bar{m}_b = (m_{\eta_b(1S),\text{exp}} + 3m_{\Upsilon(1S),\text{exp}})/4 = 9445(1) \text{ MeV}$$



Born-Oppenheimer Approximation

$L = 0$, $m_Q = m_c = 1628$ MeV:

Braaten et al. (2014)

state	n=1	n=2
Σ_g^+	$\bar{m}_c$	4333(19)
$m_{J/\Psi}(nS)$	3096.900(6)	3674(1)
Π_u	4692(15)	

$L = 0$, $m_Q = m_b = 4977$ MeV:

state	n=1	n=2	n=3	n=4
Σ_g^+	$\bar{m}_b$	10381(17)	11103(20)	11707(19)
$\Upsilon(nS)$	9460.4(1)	10023.4(5)	10355.1(5)	10579.4(1.2)
Π_u	11417(15)			

$$m_{ps}^* - m_{ps} = 932(20) \Leftrightarrow B_c^\pm(2S) - B_c^\pm(1S) = 2397(1) \text{ MeV}$$




Spatial distribution of Laplace trial states

- trivial plaquette scans color field in Wilson loop
- 'test-charge' $v^\dagger v$ in Laplace trial state



$$\psi^{(n)}(\vec{z}, R) = \left\langle \sum_{\vec{x}, t} \left\| \sum_{ij}^{N_v} \tilde{\rho}_R^{(n)}(\lambda_i, \lambda_j) v_i(\vec{x}, t) v_i^\dagger(\vec{z}, t) v_j(\vec{z}, t) v_j^\dagger(\vec{x} + R, t) \right\|_2 \right\rangle$$

with optimal profiles $\tilde{\rho}_R^{(n)}(\lambda_i, \lambda_j) = \sum_{k,l} \nu_k^{(n)} u_{k,l} e^{-\lambda_i^2/4\sigma_l^2} e^{-\lambda_j^2/4\sigma_l^2}$





Spatial distribution of Laplace trial states ($R = 10a$)

$n = 0 :$



$n = 1 :$



Spatial distribution of hybrid states


$$\psi_{\Sigma_u}^{(n)}(\vec{z}, R) = \left\langle \sum_{\vec{x}, t, \vec{k} \parallel \vec{r}} \left\| \sum_{i,j}^{N_v} \tilde{\rho}_{\Sigma_u, R}^{(n)}(\lambda_i, \lambda_j) \right. \right. \\ \left. \left[\nabla_{\vec{k}} v_i(\vec{x}, t) v_i^\dagger(\vec{z}, t) v_j(\vec{z}, t) v_j^\dagger(\vec{x} + \vec{r}, t) \right. \right. \\ \left. \left. \pm v_i(\vec{x}, t) v_i^\dagger(\vec{z}, t) v_j(\vec{z}, t) [\nabla_{\vec{k}} v_j]^\dagger(\vec{x} + \vec{r}, t) \right] \right\|_2 \right\rangle$$

$$\psi_{\Pi_{u/g}}^{(n)}(\vec{z}, R) = \left\langle \sum_{\vec{x}, t, \vec{k} \perp \vec{r}} \left\| \sum_{i,j}^{N_v} \tilde{\rho}_{\Pi_{u/g}, R}^{(n)}(\lambda_i, \lambda_j) \right. \right. \\ \left. \left[\nabla_{\vec{k}} v_i(\vec{x}, t) v_i^\dagger(\vec{z}, t) v_j(\vec{z}, t) v_j^\dagger(\vec{x} + \vec{r}, t) \right. \right. \\ \left. \left. \pm v_i(\vec{x}, t) v_i^\dagger(\vec{z}, t) v_j(\vec{z}, t) [\nabla_{\vec{k}} v_j]^\dagger(\vec{x} + \vec{r}, t) \right] \right\|_2 \right\rangle$$




Spatial distribution of hybrid states ($R = 10a$)


$\Sigma_u :$ $n = 0$




$n = 1$




$n = 2$




$\Pi_u :$ $n = 0$









$n = 1$



$n = 2$



Spatial distribution of hybrid states ($R = 10a$)

 $\Sigma_u : \quad n = 0$  $n = 1$  $n = 2$  $\Pi_u : \quad n = 0$  $n = 1$  $n = 2$ 

Static-light meson operators

$$\begin{aligned}
 C_{\mathcal{O}}^{sl}(t) &= \frac{1}{N_f} \sum_{\vec{x}, t_0, i} \left\langle \bar{Q}(\vec{x}, t_0 + t) \mathcal{O} q^i(\vec{x}, t_0 + t) \bar{q}^i(\vec{x}, t_0) \gamma_4 \mathcal{O} Q(\vec{x}, t_0) \right\rangle \\
 &= - \sum_{\vec{x}, t_0} \left\langle \text{tr}_{c,d} \left(\underbrace{\mathcal{O} \mathcal{D}(\vec{x}, t_0 + t; \vec{x}, t_0) \gamma_4}_{\text{light propagator}} \underbrace{\mathcal{O} U_t(\vec{x}; t_0, t_0 + t) P_-}_{\text{static propagator}} \right) \right\rangle
 \end{aligned}$$

$\mathcal{O}$	J^P	j^P	O_h	not.
$\gamma_5, \gamma_5 \gamma_j \nabla_j$ $1, \gamma_j \nabla_j$	$0^- [1^-]$ $0^+ [1^+]$	$(1/2)^-$ $(1/2)^+$	A_1	S P_-
$\gamma_1 \nabla_1 - \gamma_2 \nabla_2$ (and cyclic) $\gamma_5 (\gamma_1 \nabla_1 - \gamma_2 \nabla_2)$ (and cyclic)	$2^+ [1^+]$ $2^- [1^-]$	$(3/2)^+$ $(3/2)^-$	E	P_+ $D_{\pm}$
$\gamma_1 \nabla_2 \nabla_3 + \gamma_2 \nabla_3 \nabla_1 + \gamma_3 \nabla_1 \nabla_2$ $\gamma_5 (\gamma_1 \nabla_2 \nabla_3 + \gamma_2 \nabla_3 \nabla_1 + \gamma_3 \nabla_1 \nabla_2)$	$3^- [2^-]$ $3^+ [2^+]$	$(5/2)^-$ $(5/2)^+$	A_2	D_+ $F_{\pm}$



$$C_{S/P_-}^{sl}(t) = - \sum_{\vec{x}, t_0} \left\langle \text{tr}_{c,d} \left(\mathcal{D}(\vec{x}, t_0 + t; \vec{x}, t_0) \gamma_4 P_{\pm} U_t(\vec{x}; t_0, t_0 + t) \right) \right\rangle$$




Static-light meson operators

$$C_{S/P_-}^{sl}(t) = - \sum_{t_0, i, j} \left\langle \rho(\lambda_i) \rho(\lambda_j) \text{Tr}_d \{ [v_i^\dagger D^{-1} \gamma_4 v_j](t_0 + t, t_0) P_\pm \} \right. \\ \left. \sum_{\vec{x}} v_j^\dagger(\vec{x}, t_0) U_t(\vec{x}; t_0, t_0 + t) v_i(\vec{x}, t_0 + t) \right\rangle$$

- ▶ light perambulators $v_i^\dagger(t_1) D_{\alpha\beta}^{-1} \gamma_4 v_j(t_0)$ from distillation framework [Peardon et al. \(2009\)](#), [Knechtli et al \(2022\)](#)
- ▶ static perambulators $v^\dagger(\vec{x}, t_0) U_t(\vec{x}; t_0, t_0 + t) v(\vec{x}, t_0 + t)$
- ▶ Gaussian profiles ρ , SVD, GEVP $\Rightarrow$ optimal profiles...





Static-light S -wave meson states



Static-light meson states

ensemble	$N_s^3 \times N_t$	N_f	m_π	$m_s(P_-) - m_{ps}(S)$ [MeV]
Em1	$24^3 \times 48$	2	2.2GeV	$2393.5 - 1764.9 = 628.6(2.1)$
A0 heavy	$72^3 \times 24$	3+1	1GeV	$1821 - 1350.4 = 470.6(5.8)$
A1 heavy	$96^3 \times 32$	3+1	1GeV	$1815.2 - 1347.2 = 468.0(3.3)$
A1 light	$96^3 \times 32$	3+1	420MeV	$1412 - 1122.7 = 289.3(5.2)$



String breaking and tetra-quark operators

- ▶ combine static and light(charm)-quark perambulators
- ▶ building blocks for observation of string breaking

$$| = v^\dagger(0)U_t v(t) \rightarrow \mathcal{P}, \bullet \text{---} \bullet = v^\dagger(t)D_{\alpha\beta}^{-1}\gamma_4 v(0) \rightarrow \mathcal{D}$$

$$C(t) = \begin{pmatrix} \begin{array}{cc} \bullet & \bullet \\ | & | \\ \bullet & \bullet \end{array} & \sqrt{N_f} \times \begin{array}{cc} \bullet & \bullet \\ \text{---} & \text{---} \\ \bullet & \bullet \end{array} \\ \sqrt{N_f} \times \begin{array}{cc} \bullet & \bullet \\ | & | \\ \bullet & \bullet \\ \text{---} & \text{---} \end{array} & -N_f \times \begin{array}{cc} \bullet & \bullet \\ \text{---} & \text{---} \\ \bullet & \bullet \\ \text{---} & \text{---} \end{array} + \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \end{pmatrix}$$

$$C_{11}(t) \rightarrow \mathcal{P}(\vec{x})\mathcal{P}^\dagger(\vec{y}) \quad \hat{r} = |\vec{y} - \vec{x}|, P_\pm = (1 \pm \gamma_4)/2$$

$$C_{12}(t) \rightarrow \sqrt{N_f} \text{Tr}_{c,d} \mathcal{P}(\vec{x}) P_- \gamma \hat{r} \mathcal{D}(\vec{x}, \vec{y}, t) \mathcal{P}^\dagger(\vec{y})$$

$$C_{21}(t) \rightarrow -\sqrt{N_f} \text{Tr}_{c,d} \mathcal{P}(\vec{x}) \mathcal{P}^\dagger(\vec{y}) P_+ \gamma \hat{r} \mathcal{D}^\dagger(\vec{x}, \vec{y}, 0)$$

$$C_{22}(t) \rightarrow N_f \text{Tr}_{c,d} \mathcal{P}(\vec{x}) P_+ \mathcal{D}(\vec{x}, \vec{y}, t) \mathcal{P}^\dagger(\vec{y}) P_- \mathcal{D}^\dagger(\vec{x}, \vec{y}, 0) \\ - \delta_{ij} \text{Tr}_{c,d} [\mathcal{P}(\vec{x}) P_+ \mathcal{D}_i^\dagger(\vec{x}, 0, t)] \text{Tr}_{c,d} [\mathcal{P}^\dagger(\vec{y}) P_- \mathcal{D}_j(\vec{y}, 0, t)]$$



Conclusions & Outlook

- ✓ alternative operator for a static quark-anti-quark potentials based on Laplacian eigenmodes, replacing Wilson loop
- ✓ improved Laplace trial states (optimal profiles) give earlier effective mass plateaus and better signal
- ✓ computational advantage for high resolution of the potential energy as off-axis distances basically come "for free"
- ✓ hybrid static potentials (hybrid meson masses), instead of "gluonic handles" (excitations) use derivatives of v
- ✓ implementation of static-light (charm) correlator using "perambulators" $v(t_1)D^{-1}v(t_2)$ from distillation framework
- 🔧 putting together building blocks for string breaking in QCD (mixing matrix of static and light quark propagators)
- 🔧 more (hybrid) static(-light) and multi-quark potentials
- ? questions, discussion, many possible applications...



Acknowledgements

THANK YOU



Leibniz-Rechenzentrum
der Bayerischen Akademie der Wissenschaften




NRW-FAIR
Netzwerk


Ministerium für
Kultur und Wissenschaft
des Landes Nordrhein-Westfalen




Charmonium spectrum on Em1



Glueball spectrum on Em1, $m_G = 1955(75)\text{MeV}$



Glueball spectrum on Em1, $m_G = 1919(63)\text{MeV}$



Acknowledgements

THANK YOU



Leibniz-Rechenzentrum
der Bayerischen Akademie der Wissenschaften



NRW-FAIR
Netzwerk

Ministerium für
Kultur und Wissenschaft
des Landes Nordrhein-Westfalen

