

QCD at the Extremes

From DIS to $pp \rightarrow jets$

Jet production in pp

- x-section (i.e. for light and heavy quark ($t\bar{t}$) production)


$$\sigma(pp \rightarrow q\bar{q}X) = \int \frac{dx_1}{x_1} \frac{dx_2}{x_2} x_1 G(x_1, \bar{q}) x_2 G(x_2, \bar{q}) \times \hat{\sigma}(\hat{s}, \bar{q})$$

- with gluon densities

$$xG(x, \bar{q})$$

- hard x-section:


$$\frac{d\sigma}{dt} = \frac{1}{64\hat{s}^2} |M_{ij}|^2$$




Lowest Order Diagrams

Ellis, Stirling, Webber
QCD & collider physics p248


$$qq' \rightarrow qq'$$




$$qq \rightarrow qq$$



$$q\bar{q} \rightarrow gg$$



$$gg \rightarrow gg$$



2 → 2 processes

invariant matrix elements for
2 → 2 processes with
massless partons

Process	$\sum \mathcal{M} ^2 / g^4$	$\theta^* = \pi/2$
$qq' \rightarrow qq'$	$\frac{4}{9} \frac{\hat{s}^2 + \hat{u}^2}{\hat{t}^2}$	2.22
$q\bar{q}' \rightarrow q\bar{q}'$	$\frac{4}{9} \frac{\hat{s}^2 + \hat{u}^2}{\hat{t}^2}$	2.22
$qq \rightarrow qq$	$\frac{4}{9} \left(\frac{\hat{s}^2 + \hat{u}^2}{\hat{t}^2} + \frac{\hat{s}^2 + \hat{t}^2}{\hat{u}^2} \right) - \frac{8}{27} \frac{\hat{s}^2}{\hat{u}\hat{t}}$	3.26
$q\bar{q} \rightarrow q'\bar{q}'$	$\frac{4}{9} \frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2}$	0.22
$q\bar{q} \rightarrow q\bar{q}$	$\frac{4}{9} \left(\frac{\hat{s}^2 + \hat{u}^2}{\hat{t}^2} + \frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2} \right) - \frac{8}{27} \frac{\hat{u}^2}{\hat{s}\hat{t}}$	2.59
$q\bar{q} \rightarrow gg$	$\frac{32}{27} \frac{\hat{t}^2 + \hat{u}^2}{\hat{t}\hat{u}} - \frac{8}{3} \frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2}$	1.04
$gg \rightarrow q\bar{q}$	$\frac{1}{6} \frac{\hat{t}^2 + \hat{u}^2}{\hat{t}\hat{u}} - \frac{3}{8} \frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2}$	0.15
$gq \rightarrow gq$	$-\frac{4}{9} \frac{\hat{s}^2 + \hat{u}^2}{\hat{s}\hat{u}} + \frac{\hat{u}^2 + \hat{s}^2}{\hat{t}^2}$	6.11
$gg \rightarrow gg$	$\frac{9}{2} \left(3 - \frac{\hat{t}\hat{u}}{\hat{s}^2} - \frac{\hat{s}\hat{u}}{\hat{t}^2} - \frac{\hat{s}\hat{t}}{\hat{u}^2} \right)$	30.4

Remember: W+jet and dijets

- check on propagator:

$$qq \rightarrow qq$$

$$|M|^2 \sim \frac{s^2 + u^2}{t^2}$$


$$\sim \frac{1}{t^2} \sim \frac{1}{(1 - \cos\theta^*)^2}$$

$$qq \rightarrow W + g :$$

$$|M|^2 \sim \frac{1}{\cancel{t}} \sim \frac{1}{(1 - \cos\theta^*)}$$



CDF Collaboration (F. Abe et al.). Phys. Rev. Lett. 73:2296-2300, 1994.

Ellis, Stirling, Webber
QCD & collider physics p248



Color Flow in pp

- quarks carry color
- anti-quarks carry anti-color
- gluons carry color – anti-color
 - connect to color singlet systems
 - watch out pp or $p\bar{p}$




Color Flow in pp

The Lund Monte Carlo For High P(T) Physics [H.U. Bengtsson](#)
Comput.Phys.Commun.31:323,1984.

process: $gg \rightarrow q_i \bar{q}_i$


grams:



plitudes:

$$g^2 f^{abc} T_{\alpha\beta}^c \bar{u}_i^\beta(q_4) \frac{\epsilon_1^\kappa \epsilon_2^\lambda \gamma^\mu}{\hat{s}} C_{\kappa\lambda\mu}(q_1, q_2, -q_1 - q_2) v_i^\alpha(q_3) \\ - i g^2 T_{\alpha\gamma}^b T_{\gamma\beta}^a \bar{u}_i^\beta(q_4) \epsilon_1^\kappa \frac{q_1 - q_4}{\hat{t}} \epsilon_2^\lambda v_i^\alpha(q_3) \\ - i g^2 T_{\alpha\gamma}^a T_{\gamma\beta}^b \bar{u}_i^\beta(q_4) \epsilon_2^\lambda \frac{q_1 - q_3}{\hat{u}} \epsilon_1^\kappa v_i^\alpha(q_3)$$

Colour flows:



String configurations:



Colour factors: A: $T_{\alpha\gamma}^b T_{\gamma\beta}^a$; B: $T_{\alpha\gamma}^a T_{\gamma\beta}^b$

Amplitudes:

$$A: -i g^2 \bar{u}_i^\beta(q_4) \left[\epsilon_1^\kappa \frac{q_1 - q_4}{\hat{t}} \epsilon_2^\lambda + \frac{\epsilon_1^\kappa \epsilon_2^\lambda \gamma^\mu}{\hat{s}} C_{\kappa\lambda\mu}(q_1, q_2, -q_1 - q_2) \right] v_i^\alpha(q_3) \\ B: -i g^2 \bar{u}_i^\beta(q_4) \left[\epsilon_2^\lambda \frac{q_1 - q_3}{\hat{u}} \epsilon_1^\kappa + \frac{\epsilon_1^\kappa \epsilon_2^\lambda \gamma^\mu}{\hat{s}} C_{\kappa\lambda\mu}(q_1, q_2, -q_1 - q_2) \right] v_i^\alpha(q_3)$$

Cross-sections:

$$A: \frac{\pi\alpha_s^2}{\hat{s}^2} \frac{1}{6} \left(\frac{\hat{u}}{\hat{t}} - 2 \frac{\hat{u}^2 + \hat{t}^2}{\hat{s}^2} \right); \quad B: \frac{\pi\alpha_s^2}{\hat{s}^2} \frac{1}{6} \left(\frac{\hat{t}}{\hat{u}} - 2 \frac{\hat{t}^2}{\hat{s}^2} \right)$$





$$A + B = \dots \left(\frac{\hat{u}}{\hat{t}} - 2 \frac{\hat{u}^2 + \hat{t}^2}{\hat{s}^2} + \frac{\hat{t}}{\hat{u}} \right) = \frac{\hat{u}^2 + \hat{t}^2}{\hat{t}\hat{u}} - 2 \frac{\hat{u}^2 + \hat{t}^2}{\hat{s}^2}$$

Color Flow in pp

Process: $gg \rightarrow gg$

The Lund Monte Carlo For High P(T) Physics [H.U. Bengtsson](#)
Comput.Phys.Commun.31:323,1984.

Diagrams:



Amplitudes:







$$s: -ig^2 \frac{1}{s} f^{abe} f^{cde} \epsilon_1^\kappa \epsilon_2^\lambda \epsilon_3^\mu \epsilon_4^\nu C_{\kappa\lambda\tau}(-q_1, -q_2, q_1 + q_2) C_{\mu\nu\tau}(q_3, q_4, -q_3 - q_4)$$

$$t: -ig^2 \frac{1}{t} f^{dae} f^{bce} \epsilon_1^\kappa \epsilon_2^\lambda \epsilon_3^\mu \epsilon_4^\nu C_{\nu\kappa\tau}(q_4, -q_1, q_1 - q_4) C_{\lambda\mu\tau}(-q_2, q_3, q_2 - q_3)$$

$$u: -ig^2 \frac{1}{u} f^{cae} f^{bde} \epsilon_1^\kappa \epsilon_2^\lambda \epsilon_3^\mu \epsilon_4^\nu C_{\mu\kappa\tau}(q_3, -q_1, q_1 - q_3) C_{\lambda\nu\tau}(-q_2, q_4, q_2 - q_4)$$


$$4: -ig^2 f^{abe} f^{cde} (g_{\kappa\mu} g_{\lambda\nu} - g_{\kappa\nu} g_{\lambda\mu}) - ig^2 f^{dae} f^{bce} \\ \times (g_{\kappa\mu} g_{\lambda\nu} - g_{\kappa\lambda} g_{\nu\mu}) + ig^2 f^{cae} f^{bde} (g_{\kappa\lambda} g_{\mu\nu} - g_{\kappa\nu} g_{\lambda\mu})$$

Colour flows:



The general pp case

- basic elements are:
 - **Matrix Elements:**
 - on shell/off shell
 - **PDFs**
 - TMD PDFs
 - **Parton Shower**
 - angular ordering
- Proton remnant and hadronization



$$\sigma(pp \rightarrow q\bar{q} + X) = \int \frac{dx_{g1}}{x_{g1}} \frac{dx_{g2}}{x_{g2}} \int d^2 k_{t1} d^2 k_{t2} \hat{\sigma}(\hat{s}, k_t, \mu) \times x_{g1} \mathcal{A}(x_{g1}, k_{t1}, \mu) x_{g2} \mathcal{A}(x_{g2}, k_{t2}, \mu)$$

Parton shower


- simulate explicitly parton radiation with evolution of parton densities
- advantage to include properly energy momentum conservation in each step
- perform resummation numerically
- Parton shower follows backward evolution:

$$\Pi = \exp \left[- \int_{\mu_l^2}^{\mu_h^2} \frac{d\mu'^2}{\mu'^2} \int^{z_{\text{dyn}}} \frac{dz}{z} \hat{P}(z) \frac{f(x/z, \mu^2)}{f(x, \mu^2)} \right]$$

- Emitted partons should have *resolvable* energy (or p_T) with: $p_T > q_{t \text{ cut}} \sim 1\text{GeV}$

$$z_{\text{dyn}} = 1 - \frac{q_{t \text{ cut}}}{\mu'}$$


- With $z_{\text{dyn}} \ll 1$ soft gluons with $p_T < 1\text{GeV}$ are neglected.



Parton shower


- Parton shower with collinear parton densities adds new information on transverse momentum of partons, by explicitly undoing the evolution.
 - not necessarily same parameters as in evolution:
 - $z_{\text{dyn}} \ll 1$ versus $z \rightarrow 1$
 - order of splitting functions
 - α_s heavy flavors
 - ordering conditions (angular ordering ?)
 - etc
 - inconsistencies arise from energy momentum conservation
 - special treatment needs to be done to conserve energy momentum at the end
 - boost, rotation of hard process

$$\alpha_s^{PS}(p_\perp)$$






TMD parton shower

- TMD parton shower uses information from TMDs (obtained from fits to inclusive data)
 - uses exactly the same parameters as in evolution
 - cut on z values
 - order of splitting functions
 - α_s , heavy flavors
 - ordering conditions (angular ordering...)
 - inconsistencies arise from energy momentum conservation stay, when using collinear ME
 - special treatment needs to be done to conserve energy momentum at the end
 - boost, rotation of hard process



Parton shower and TMDs

- Check that including initial state PS and TMDs does not change the x-section of the hard process.



How to calculate x-sections then ?

MCatNLO method

- NLO calculation of dijets:

$$d\sigma = \left[B(\Phi_B) + \hat{V}(\Phi_B) \right] + R(\Phi_R) d\Phi_r$$

- diverging spectrum of gluon radiation
- cancelled by virtual corrections
- spectrum at low p_T of gluon (or p_T of dijet system) unphysical
- introduce auxiliary counter term $C(\Phi_R)$

$$d\sigma = \left[B(\Phi_B) + V(\Phi_B) + \int d\Phi C \right] + R(\Phi_R) d\Phi_r - C d\Phi$$

- subtract and add collinear, soft divergent term
 - calculated matrix element with subtraction
 - physical quantity obtained only if $C(\Phi_R)$ is added back by TMD or parton shower
- method used in MCatNLO with CASCASDE3 for TMD or with HERWIG/PYTHIA for parton shower





MCatNLO method

- **MCatNLO** method (Frixione, S. and Webber, B. R. *Matching NLO QCD computations and parton shower simulations*, *JHEP*, 06(2002), 029)):



- subtract collinear and soft part
- add it back by using TMD

$p_{T,12} > 100 \text{ GeV}$





Jet production at the LHC

Tumasyan, A. and others Measurement and QCD analysis of double-differential inclusive jet cross sections in proton-proton collisions at $\sqrt{s} = 13$ TeV, JHEP, 022022(), 142




Jet production to constrain PDFs

Tumasyan, A. and others Measurement and QCD analysis of double-differential inclusive jet cross sections in proton-proton collisions at $\sqrt{s} = 13$ TeV, JHEP, 022022(), 142







Jet Mass – Sudakov peak

- Jets in final state: time-like evolution of parton radiation
 - follows similar evolution equation as for parton densities
 - evolution from large time-like mass to small scale where hadronization happens
- Evolution according to Sudakov form factors (similar to initial state)
- Jet mass comes from parton radiation:
 - low mass
 - at low transverse momenta, suppression from Sudakov
 - large mass
 - perturbative radiation pattern



Jet Mass – Sudakov peak


Sirunyan, A. M. and others Measurements of the differential jet cross section as a function of the jet mass in dijet events from proton-proton collisions at $\sqrt{s}=13$ TeV, JHEP, 112018(), 113



k_t - factorization

- use high energy (kt -) factorization:

(Catani, Ciafaloni, Hautmann NPB 366 (1991) 135,
Gribov, Levin, Ryskin, Phys. Rep. 100, (1983), 1,
Collins, Ellis, NPB 360, (1991), 3)




- with $\sigma(ep \rightarrow e' q \bar{q}) = \int \frac{dy}{y} d^2 Q \frac{dx_g}{x_g} \int d^2 k_t \hat{\sigma}(\hat{s}, k_t,) x_g \mathcal{A}(x_g, k_t,)$

$$\int^{Q^2} d^2 k_t x_g \mathcal{A}(x_g, k_t, \bar{q}) \simeq x_g G(x_g, Q^2)$$

- t -channel gluon with virtuality $k^2 = -k_t^2$ dominates the process in the high energy limit $s \gg \hat{s}$
- collinear limit obtained by: $\hat{\sigma}(\hat{s}, 0, Q) \cdot \Theta(Q - k_T)$

- **BUT k_t -factorization is proven only for small x **



Off-shell matrix element calculation

A.van Hameren Katie, <https://arxiv.org/abs/1611.00680>

- Automated tool exists for LO calculation: KaTie
 - however k_t factorization only proven for small x
- NLO calculation much more difficult, since virtual corrections are also k_t dependent
- First steps toward automated NLO calculation are done
- Where are k_t -dependent matrix elements relevant ?

Why off-shell matrix elements ?

Baranov, S. and others CASCADE3 A Monte Carlo event generator based on TMDs,
Eur. Phys. J. C, 81(2021), 425

- Behavior of ME as function of k_t :

- for small k_t converges to collinear result


- for large k_t has suppression
- suppression appears at “standard factorization scale”: $Q^2 + 4m^2$

- collinear factorization:

$$\mu^2 \sim Q^2 + 4m^2$$

$$\int_0^{\mu^2} d\hat{\sigma}(, \dots)$$



$$\mu^2 \sim 4m^2_{\text{TMD/PS}} \perp i < \mu^2$$



Predictions with TMDs and off-shell ME

- Calculate jet production with KaTie and TMDs
 - no special kinematic reshuffling needed
 - even without explicit parton shower, $\Delta\phi$ well described



<https://arxiv.org/abs/1712.05932>



Predictions with TMDs and off-shell ME



- Calculate jet production with KaTie and TMDs
 - no special kinematic reshuffling needed
 - even without explicit parton shower, $\Delta\phi$ well described
 - for small $\Delta\phi$ parton shower is needed – more jets

<https://arxiv.org/abs/1712.05932>





MCatNLO method

- **MCatNLO** method (*Frixione, S. and Webber, B. R. Matching NLO QCD computations and parton shower simulations, JHEP, 06(2002), 029*):
 - subtract collinear and soft part
 - add it back by using TMD
- check $\Delta\phi$ distribution




Measurements at large p_T . $\Delta\phi$






Jets at large p_T

- At large p_T , jets become smaller, pencil-like
- much better experimental resolution
 - enables studies in new environment






Measurements at large p_T . $\Delta\phi$ back-to-back



- In back-to-back region

- for $p_T^{leading} > 200 \text{ GeV}$ sensitive to “soft gluon” k_T kick of $\sim 10 \text{ GeV}$




Measurements at large p_T . $\Delta\phi$ back-to-back




- In back-to-back region

- for $p_T^{leading} > 900 \text{ GeV}$ sensitive to “soft gluon” k_T kick of $\sim 100 \text{ GeV}$

Measurements at large p_T . $\Delta\phi$ back-to-back





Measurements at large p_T . $\Delta\phi$ back-to-back



- Sensitivity in back-to-back region on differences in “small kt” region

Measurements at large p_T



- differences to standard parton shower approaches are observed
- details matter

But even this is not the full story...

- factorization breaking in $pp \rightarrow j_1 j_2 X$

J. Collins, J.W. Qiu [hep-ph 0705.2141](#)





FIG. 8 (color online). The exchange of two extra gluons, as in this graph, will tend to give nonfactorization in unpolarized cross sections.

- factorization breaking also in $t\bar{t}$ -production at large p_t^{top}

S. Catani, M. Grazzini, and A. Torre. Transverse-momentum resummation for heavy-quark hadroproduction. [arXiv 1408.4564](#)





Testing possible factorization breaking

- No real calculations are available
- Compare with experimental measurements
 - and test with scenario, where factorization effects are absent or at least much smaller:
 - compare jj with Zj (in Zj color connections to of initial to final state are reduced, since Z is colorless).





<https://arxiv.org/abs/2204.01528>






Dijet and Z+j comparisons I

<https://arxiv.org/abs/2204.01528>




Dijet and Z+j comparisons II

<https://arxiv.org/abs/2204.01528>



Dijet and Z+j comparisons III

<https://arxiv.org/abs/2204.01528>



The role of soft gluons

Role of soft gluons in inclusive distributions

- Does evolution equation work with cut z_M ?

$$\int_x^1 dz P(z) \rightarrow \int_x^{z_M} dz P(z)$$

$$\mu^2 \frac{df_a}{d\mu^2} = \int \frac{dz}{z} \frac{d_s}{2\pi} P_{ab}(z) \int_b \left(\frac{x}{z}, z^2 \right)$$

$$P_{ab} = \underbrace{R_{ab}} + \frac{\bar{K}_{ab}}{(1-z)_+} + \underbrace{D_{ab} \delta(1-z)}$$

$$PCR1 = R_{ab} + \frac{K_{ab}}{(1-z)}$$

$$\mu^2 \frac{df}{d\mu^2} = \int \frac{dz}{z} \frac{d_s}{2\pi} P \cdot f\left(\frac{x}{z}\right) = \int \frac{dz}{z} \frac{d_s}{2\pi} \left(R_{ab} + \frac{K_{ab}}{(1-z)_+} + D_{ab} \delta(1-z) \right) f$$

$$= \int_x^1 \frac{dz}{z} \frac{d_s}{2\pi} \left(R_{ab} f + \frac{f(x/z) - f(x)}{1-z} \right) + D_{ab} f(x)$$

$$= \int_x^{z_M} \frac{dz}{z} \left(R_{ab} f + \frac{1}{1-z} \cdot f \right)$$


$$+ \int_{z_M}^1 \frac{dz}{z} \left(\cancel{R_{ab} f} + \frac{\cancel{f(x/z)}}{1-z} - \frac{\cancel{f(x)}}{1-z} \right) + D_{ab} f(x)$$

$$\boxed{z_M \rightarrow 1}$$

Role of soft gluons in inclusive distributions

<https://arxiv.org/abs/2312.08655>

- Perform evolution with PB method with and without cut $z_{\text{dyn}} = 1 - \frac{q_t \text{ cut}}{\mu'}$




$$q_t > 1$$

$$z < 1 - \frac{q_t}{\mu'}$$

Role of soft gluons in TMD distributions

<https://arxiv.org/abs/2312.08655>

- Perform evolution with PB method with and without cut $z_{\text{dyn}} = 1 - \frac{q_{t \text{ cut}}}{\mu'}$
down, $x = 0.01$, $\mu = 100 \text{ GeV}$



Cutoff and connection to cross section

- For simplicity consider again DIS cross section (at NLO)

<https://arxiv.org/abs/2309.11802>

$$F_2(x, Q^2) = x \sum_{q, \bar{q}} e_q^2 \int_x^1 \frac{d\xi}{\xi} q(\xi, Q^2) \left[\delta \left(1 - \frac{x}{\xi} \right) + \frac{\alpha_s}{2\pi} C_{\overline{\text{MS}}} \left(\frac{x}{\xi} \right) + \dots \right]$$

- with:



$$C_q^{\overline{\text{MS}}}(z) = C_F \left[2 \left(\frac{\log(1-z)}{1-z} \right)_+ - \frac{3}{2} \left(\frac{1}{1-z} \right)_+ - (1+z) \log(1-z) \right. \\ \left. - \frac{1+z^2}{1-z} \log z + 3 + 2z - \left(\frac{\pi^2}{3} + \frac{9}{2} \right) \delta(1-z) \right]$$

- to ensure proper cancellation of terms in parton density as well as in cross section, $z_M \rightarrow 1$
- see similar study in Frixione, Webber <https://arxiv.org/abs/2309.15587>

Soft gluons in Parton Shower

<https://arxiv.org/abs/2309.11802>

- With PB-TMD parton shower study effect of $z_{\text{dyn}} = 1 - \frac{q_{t \text{ cut}}}{\mu'}$




- Huge effect on soft partons
 - they are important for inclusive distributions: pdfs and TMDs

Soft gluons in Parton Shower

<https://arxiv.org/abs/2309.11802>

- With PB-TMD parton shower study effect of $z_{\text{dyn}} = 1 - \frac{q_t \text{ cut}}{\mu'}$





- Huge effect on soft partons
 - they are important for inclusive distributions: pdfs and TMDs
 - see DY q_T spectra
- in Lund string fragmentation these soft partons do not change hadron spectra
 - no issue for hadrons or jets from parton shower !

Expectations at high luminosity

Expectations at highest luminosity: jets I



<https://arxiv.org/abs/1902.04070>



- At highest luminosity (13 TeV, $L=3\text{ab}$)
 - jets with p_T up to 4 TeV can be observed
 - dijet masses up to 6 TeV ($\sim 1/2$ of $\sqrt{s}$)

Expectations at highest luminosity: jets II



<https://arxiv.org/abs/1902.04070>



- At highest luminosity (13 TeV, L=3ab)
 - p_T distribution for different flavors (even for boosted W and top) are similar
 - masses play no role
 - $\Delta\phi$ shows differences at smaller angles, due to different production mechanism

Expectations at highest luminosity: jets III

<https://arxiv.org/abs/1902.04070>



- At highest luminosity (13 TeV, $L=3\text{ab}$)
 - b-jets come mainly from $g \rightarrow b\bar{b}$
 - look for pairs of B-mesons inside jet:
 - at high p_T fraction of pairs of B-mesons is large $> 70\%$
 - b-jet comes mainly from gluon splitting