

Contribution submission to the conference Karlsruhe 2024

Distinguishing Axion-Like Particles from Extended Higgs Sector Models in $t\bar{t}$ production at the LHC — ANKE

BIEKÖTTER¹, THOMAS BIEKÖTTER², ALEXANDER GROHSJEAN³, SVEN HEINEMEYER⁴, •LAURIDS JEPPE⁵, CHRISTIAN SCHWANENBERGER^{5,3}, and GEORG WEIGLEIN^{5,3} — ¹PRISMA+ Cluster of Excellence & Institute of Physics (THEP) & Mainz Institute for Theoretical Physics, Johannes Gutenberg University, Mainz, Germany — ²Institute for Theoretical Physics, Karlsruhe Institute of Technology, Karlsruhe, Germany — ³Universität Hamburg, Hamburg, Germany — ⁴Instituto de Física Teórica UAM-CSIC, Madrid, Spain — ⁵Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany

We present an analysis of the sensitivity of LHC searches for new spin-0 particles produced via gluon-fusion and decaying into top quark-antiquark ($t\bar{t}$) final states to generic axion-like particles (ALPs) coupled to top-quarks and gluons. We derive new limits on the effective ALP Lagrangian in the linear representation in terms of the Wilson coefficients c_t and c_G based on the existing CMS search using 35 fb^{-1} collected at $\sqrt{s} = 13\text{ TeV}$. We further investigate possible distinctions between ALPs and pseudoscalar Higgs bosons as predicted by the two-Higgs-doublet model (2HDM), and find that distinction is possible with data anticipated to be collected during the high-luminosity of the LHC for a significant range of the effective ALP-gluon coupling.

Part: T
Type: Vortrag;Talk
Topic: 2.18 Suche nach neuen Teilchen; 2.18
Search for New Particles
Keywords: BSM; ALP; Heavy Higgs; 2HDM; Top
Email: laurids.jeppe@desy.de