

Measuring kinematic anisotropies with pulsar timing arrays

Ameek Malhotra

Whispers from the Dark Universe (2024),
DESY-Hamburg

*Based on Phys. Rev. D 110, 063526
and [arxiv:2406.04957](https://arxiv.org/abs/2406.04957) with Marisol Cruz,
Gianmassimo Tasinato and Ivonne Zavala*



Swansea
University
Prifysgol
Abertawe

Outline


- ▶ PTA SGWB detection
 - SGWB Anisotropy
 - Kinematic dipole
 - Current Limits and forecasts
- ▶ Summary

**What is the origin of the PTA
signal?**

Origin of PTA signal

Supermassive BH mergers expected to produce amplitude $h_c \sim 10^{-15}$ and spectral index $\gamma = 13/3$ [Phinney (2001), Sesana et al. (2008)+]


The likely one



IPTA joint analysis

Origin of PTA signal

Or is it from the early universe?



NANOGrav 15 year analysis: Search for signals from new physics

Inflation

Large density perturbations \leftrightarrow PBH

Phase Transitions

Additional possibilities studied in
[arXiv: 2306.16219, 2306.16227 + many more!]

Origin of PTA signal


Too early to tell...

- Additional SGWB properties important to figure out origin(s)
- Can SGWB anisotropies help? — currently data is consistent with isotropy

SMBHB Anisotropies


Estimates vary, but **SMBHB** anisotropies are expected to be large

[Mingarelli et al. 2013; Taylor & Gair 2013; Mingarelli et al. 2017), Sato-Polito & Kamionkowski (2023) + more]




NANOGrav 15-year Anisotropic Gravitational-Wave Background

Cosmological SGWB anisotropies



CMB observations indicate large scale inhomogeneity at the 10^{-5} level




In general, cosmological SGWB anisotropies are also expected to be small

See [review by LISA CosWG \(2022\)](#)

Kinematic dipole anisotropy

Largest anisotropy in the CMB is the kinematic dipole

Velocity $\beta = v/c = 1.23 \times 10^{-3}$ towards
 $(l, b) = (264^\circ, 48^\circ)$ in galactic coordinates




COBE dipole detection (1994)

SGWB Kinematic dipole

If SGWB is of early universe origin, then we can expect a kinematic dipole mirroring the CMB dipole

$$I(f, \hat{n}) = \bar{I}(f) \left[1 + (1 - n_I) \beta (\hat{n} \cdot \hat{v}) + \mathcal{O}(\beta^2) \right]$$

$$n_I \equiv \frac{d \ln \bar{I}}{d \ln f}$$



[Cusin and Tasinato (2022)]

PTA response to kinematic dipole



Cross-correlation of timing residuals

$$\langle \delta t_p \delta t_q \rangle \propto \Gamma_{pq}^{(\text{HD})} I + \Gamma_{pq}^{(\text{dipole})} \beta (1 - n_I) I$$

$$\Gamma_{pq}^{\text{dipole}} = \left(\frac{1}{12} + \frac{y_{pq}}{2} + \frac{y_{pq} \ln y_{pq}}{2(1 - y_{pq})} \right) [\hat{v} \cdot \hat{p} + \hat{v} \cdot \hat{q}]$$

$$y_{pq} \equiv \frac{1 - \hat{p} \cdot \hat{q}}{2}$$


[Anholm et al. (2009), Mingarelli et al. (2013), Tasinato (2023)]




Dipole tension

CMB and LSS estimates of β appear to be in tension, $\beta_{\text{LSS}} \approx 2\beta_{\text{CMB}}$

[See Peebles (2022) for a review]



COBE dipole




16.6 source deg^{-2} 17.2

Secrest et al. (2020)

Analysis of NANOGrav data

We implement the kinematic dipole ORF in Enterprise¹ with $\hat{v} = \hat{v}_{\text{CMB}}$


Note: this assumes SGWB of cosmological origin!



Parameter means and 95% C.L

1. <https://github.com/nanograv/enterprise>

What comes next?




May be enough to detect SMBHB anisotropy

But not enough for the kinematic dipole...

Forecasts: SKA era

Idealised scenario with $N \gg 100$ identical pulsars distributed uniformly


We make several simplifying assumptions \rightarrow most optimistic estimate



[Keane et al. (2015), Janssen et al. (2015)]

Forecasts: SKA era


Weak signal results



~30° degree localisation of dipole direction


Challenging even with ~4000 pulsars

$$\beta \times 10^3 = 1.23 \pm 0.61$$



Forecasts: SKA era

Strong signal regime



Detection will be challenging even for futuristic experiments

See also [Depta et al. \(2024\)](#) for strong signal results

Circular polarisation

Cosmological sources e.g. GW from axion-gauge fields [Unal et al. 2023 + more]


PTA blind to circular polarisation monopole — planar detector

$$\Gamma_{ab}^V = \beta (n_V - 1) G_{ab}^{(1)} V$$

PTA response begins at dipole

$$G_{ab}^{(1)} = - \left(\frac{1}{3} + \frac{y_{ab} \ln y_{ab}}{4(1 - y_{ab})} \right) [\hat{v} \cdot (\hat{x}_a \times \hat{x}_b)]$$

Circular polarisation



Degree of circular polarisation

$$\epsilon_V = \frac{V}{I}$$


Unconstrained by current data
(again for cosmo SGWB)

Near maximal polarisation may be detected with SKA ($N_{\text{psr}} \gtrsim 10^3$)


AGWB circular polarisation

Astrophysical estimate $C_\ell^V \simeq C_\ell^I$

[Dall'Armi et al. (2023), Sato-Polito and Kamionkowski (2023)]



$\epsilon_V = 1$ with $N_{\text{psr}} = 150$



Summary

Anisotropy could help distinguish **SMBHB** vs **cosmological scenarios**

Kinematic dipole the largest anisotropy for cosmological SGWB + dependence on pulsar direction w.r.t dipole — challenging even with SKA

Additional information in circular polarisation for both cosmological and astrophysical SGWB

Thank you!

Contact: ameek.malhotra@swansea.ac.uk