

# LHC Physics - Higgs




Image:  
ATLAS/CERN

**Lydia Beresford**

DESY Summer Student Lectures

01.08.24





# Physics Goals of the LHC



**Search for the  
Higgs Boson**



# The Standard Model (SM)

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + i\bar{\psi}\not{D}\psi + h.c.$$

$$+ \bar{\psi}_i y_{ij} \psi_j \phi + h.c.$$

Higgs-fermion interactions

$$+ |D_\mu \phi|^2 - V(\phi)$$

Higgs-self interactions

Higgs-gauge boson (W,Z) interactions

See Markus Diehl's lectures for more on QCD

# The Brout-Englert-Higgs mechanism in the SM

Introduction of the presence of a scalar field into the SM leads to


- **Particles acquire mass**
  - **Bosons:** 3 out of 4 through electroweak symmetry breaking
  - **Fermions:**  $\propto$  Yukawa  $y_{ij}$  couplings
- **Prediction of the existence of a particle**
  - **Higgs Boson**
  - **Higgs Boson interacts with itself**





# The situation before the LHC

Status 2009: SM Higgs mass above 114 GeV and NOT in range 160-170 GeV



Prediction from EW fits 2012: SM Higgs mass  $95^{+30}_{-23} \text{ GeV}$




# Interlude: how to read limit plots

Limit plots are used if we don't see a significant signal

y-axis = 95% CL upper limit on "signal strength"

$$\mu = \sigma_{\text{meas}} / \sigma_{\text{SM}}$$






# Interlude: how to read limit plots

y-axis = 95% CL  
upper limit on  
“signal strength”

$$\mu = \sigma_{\text{meas}} / \sigma_{\text{SM}}$$

Expected  $\pm 1\sigma$   
uncertainty band

Expected  $\pm 2\sigma$   
uncertainty band



With 95% confidence level, we can say that the real signal strength is smaller than the indicated value (calculate for each mass point separately)




# Interlude: how to read limit plots

y-axis = 95% CL  
upper limit on  
“signal strength”

$$\mu = \sigma_{\text{meas}} / \sigma_{\text{SM}}$$

Expected  $\pm 1\sigma$   
uncertainty band

Expected  $\pm 2\sigma$   
uncertainty band




Excluded Tevatron range since observed limit goes below 1



# Interlude: how to read limit plots

The limit for a given model will improve by adding more data

However if a signal is there the observed limit does not improve anymore



If the observed limit is better than the expected  $\rightarrow$  you have a deficit



If the observed limit is worse than the expected  $\rightarrow$  you have an excess

“better” = lower on the y-axis here



# Higgs production modes at the LHC



## Higgs production cross section as a function of $M_H$





# Higgs production modes at the LHC



## Higgs production cross section as a function of $M_H$





# Higgs production modes at the LHC


## Higgs production cross section as a function of $M_H$






# Higgs production modes at the LHC

## Higgs production cross section as a function of $M_H$




## Higgs production in association with $t\bar{t}$ (ttH)






# Higgs decay modes: a little bit of everything





# Higgs decay modes: a little bit of everything




Quiz: can you name the Higgs “discovery channels”?




# Higgs to $\gamma\gamma$

**Fairly clean signature: 2 photons & reconstruct  $m_{\gamma\gamma}$**

- Very good mass resolution  $\rightarrow$  Excellent channel to measure  $m_{\gamma\gamma}$
- Large but smoothly falling di-photon background




**Higgs discovery**



Phys. Lett. B 716 (2012) 1-29

**Full Run 2**



JHEP 07 (2023) 088






# Higgs to ZZ

**Very clean signature: 4 leptons** (electrons & muons, 2 same flavour opposite sign pairs)

**Channel with high S/B ratio**

- Low rate due to branching fraction of ZZ and  $Z \rightarrow$  leptons







# Higgs to ZZ

**Very clean signature: 4 leptons** (electrons & muons, 2 same flavour opposite sign pairs)

**Channel with high S/B ratio**

- Low rate due to branching fraction of ZZ and  $Z \rightarrow$  leptons
- Typically one Z is on-mass shell & the trailing lepton is at low  $p_T$






# Higgs to WW


## Final states including two leptons & two neutrinos

- Higgs mass diluted by presence of neutrinos →  $m_T$  variable is used


**Large event rate but also large bkg from SM WW and top pair production**



### Higgs discovery



### Full Run 2








# Aside: control regions

Large event rate but also large bkg from SM WW and top production

→ Control regions in data needed to estimate backgrounds

## Control Region (CR):

- Little or no signal expected
- Orthogonal to the Signal Region (SR)






# Discovery!


- Strongly motivated
- Significance  $\uparrow$  with luminosity
- Two experiments: ATLAS & CMS
- Several channels

p-value reflects consistency of observed data with the absence of signal





# Higgs production modes at the LHC




Production mode	Cross section (pb)
$ggH$	$48.31 \pm 2.44$
$VBF$	$3.771 \pm 0.807$
$WH$	$1.359 \pm 0.028$
$ZH$	$0.877 \pm 0.036$
$ttH$	$0.503 \pm 0.035$
$bbH$	$0.482 \pm 0.097$
$tH$	$0.092 \pm 0.008$

## At $m_H = 125$ GeV

- $gg \rightarrow H$ : main production mode
- Followed by  $VBF$  then  $WH$

# Higgs decay modes: a little bit of everything




Decay channel	Branching fraction (%)	
$bb$	57.63	$\pm 0.70$
$WW$	22.00	$\pm 0.33$
$gg$	8.15	$\pm 0.42$
$\tau\tau$	6.21	$\pm 0.09$
$cc$	2.86	$\pm 0.09$
$ZZ$	2.71	$\pm 0.04$
$\gamma\gamma$	0.227	$\pm 0.005$
$Z\gamma$	0.157	$\pm 0.009$
$ss$	0.025	$\pm 0.001$
$\mu\mu$	0.0216	$\pm 0.0004$

## At $m_H = 125$ GeV

- $H \rightarrow bb$ : main decay mode but large bkg
- $H \rightarrow \gamma\gamma$ ,  $H \rightarrow ZZ$  and  $H \rightarrow WW$  are the “discovery channels”




# The Higgs turned 10!



### A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery

Ten years after the discovery of the Higgs boson, the ATLAS experiment at CERN probes its kinematic properties with a significantly larger dataset from 2015–2018 and provides further insights on its interaction with other known particles.


The ATLAS Collaboration



### A portrait of the Higgs boson by the CMS experiment ten years after the discovery

The most up-to-date combination of results on the properties of the Higgs boson is reported, which indicate that its properties are consistent with the standard model predictions, within the precision achieved to date.

The CMS Collaboration





# Higgs to $b\bar{b}$

## Highest branching ratio of Higgs decays to two b-quarks

- Large SM backgrounds
- Statistical combination of various “channels” or “regions”
- Often machine learning techniques used



## 3 main channels targeting WH & ZH:

0 leptons ( $Z \rightarrow \nu\nu$ ),

1 lepton ( $W \rightarrow \mu\nu, e\nu$ ),

2 leptons ( $Z \rightarrow \mu\mu, ee$ )


**See lectures by Matthias Komm for  
introduction to Machine Learning**






# Higgs production modes at the LHC


**Gluon fusion (ggF)**




**Vector-boson fusion (VBF)**



**Higgs-strahlung (VH)**




**Higgs production in association with  $t\bar{t}$  (ttH)**



Observation of the **ttH** process provides direct access to the top Yukawa coupling of the Higgs

# ttH: direct probe of top Yukawa coupling

- **Very small production cross section:**  
**one of latest discoveries**
- **Large number of complex final states:**  
Mixture of b-jets, leptons, taus and photons
- **Many different channels:** many different bkg and different systematic uncertainties  
→ Excellent way to cross check each other





# What does the SM predict for the Higgs boson?

**Width**  
~4 MeV for 125 GeV



**Spin and CP**  
Scalar: spin 0, CP even

**Couplings**  
Higgs-Fermion couplings  $\propto$  fermion mass  
Higgs-Boson couplings  $\propto$  boson mass<sup>2</sup>

→ **SM Higgs sector is overall very predictive:**

Knowing the fermion masses, only free parameter is  $m_H$

# Let's test these predictions



- **Measure all properties:**  
Mass, spin, CP, couplings
- **Deviations could point to physics beyond the SM**
- **Higgs can also play an important role in searches for new physics**






# Higgs mass measurements

- Not predicted by SM
- Mass measurements in “golden channels”:  $H \rightarrow \gamma\gamma$  and  $H \rightarrow ZZ$
- Optimised analyses in categories with best mass resolution (photon, electron and muon)
- Reached 0.09% precision



# Standard Model fits after the Higgs discovery: 2022




Knowing the Higgs boson mass has large impact on global fits  
 (compare grey vs blue)



# Higgs width

- **What is the “width” of a particle?**
- Heisenberg uncertainty principle implies energy (i.e. also mass) of all unstable particles must have uncertainty  
Width is inversely proportional to lifetime
- **Larger the width smaller the lifetime**
- **Higgs width predicted to be ~4 MeV**

## Breit-wigner line shape



# Higgs width

## Two ways to access Higgs width:


- **Direct mass measurement:** Limited by experimental resolution to  $\sim 1\text{-}2$  GeV
- **Indirect methods e.g. using off-shell signal strength (away from peak):**

On-shell cross section depends on width, off-shell does not

→ **ratio is sensitive to width!**

Latest CMS result ( $H \rightarrow ZZ$ ):

$$\Gamma_H = 3.2^{+2.4}_{-1.7} \text{ MeV}$$





# Higgs width


## Two ways to access Higgs width:

- **Direct mass measurement:** Limited by experimental resolution to  $\sim 1\text{-}2$  GeV
- **Indirect methods e.g. using off-shell signal strength (away from peak):**

On-shell cross section depends on width, off-shell does not

→ **ratio is sensitive to width!**

**Quiz: why would a Higgs width  $>$  SM prediction be exciting?**



# Higgs spin and CP

## Spin (SM = 0)

- **Spin 1 excluded** using ZZ, WW decays (and by  $H \rightarrow \gamma\gamma$ )
  - **Spin 2 excluded** for number of different tensor structures (~99.9%)
- **Very likely spin 0 as predicted for the SM Higgs**


## Parity (SM: even)

- **Parity odd excluded** at  $> 99.9\%$  (ATLAS, CMS)
- **Admixtures** (CP even and CP odd couplings) **still possible**



# Higgs couplings


So far all measured couplings consistent with SM



Higgs-Fermion couplings  
 $\propto$  fermion mass

Higgs-Boson couplings  
 $\propto$  boson mass<sup>2</sup>

# Higgs couplings summary



No evidence yet  
Probably needs  
future colliders

**Established**


No evidence yet  
No clear route  
to conclusively  
establishing  
Standard Model  
couplings

First evidence  
To be conclusively  
established at the LHC  
within 5–10 years



# Undiscovered decays

## Example $H \rightarrow \mu\mu$



Nature 607 52–59 (2022)



p-value reflects consistency of observed data with the absence of signal

## Challenging: small coupling, large Drell-Yan bkg


→ Categorise events by production mode

→ Use of sophisticated machine learning techniques


## Achieved evidence ( $> 3\sigma$ )



# Di-Higgs production



## Holy grail: di-Higgs production



[arXiv:2406.09971](https://arxiv.org/abs/2406.09971)

- **Extremely interesting but very rare ~1000x rarer than H**
- **Enables us to test the Higgs self-coupling**
- **Deviations from SM expected in many BSM models**

**Higgs-self coupling will be a key focus at the HL-LHC**





# BSM Higgs searches

We indirectly search for BSM physics via precisely measuring the Higgs

We also perform direct searches e.g.

- Flavour violating searches e.g.  $H \rightarrow e\mu$
- Invisible decays of Higgs bosons
- Additional Higgs bosons: lighter, heavier, charged?





# Next lecture




**Search for  
New Physics**





# The Standard Model - fundamental particles







# The Higgs boson - very brief theory reminder

3

Higgs mechanism:

- makes use of one Higgs doublet of complex scalar fields
- to spontaneously break the  $SU(2)_L \times U(1)_Y$  symmetry
- to generate in a gauge invariant way
- the masses of the  $W^\pm$ ,  $Z$  gauge bosons and the fermions

Basically:

In order to give the gauge bosons mass and keep gauge invariance

- > introduce a Higgs field, with a scalar potential
- > find ground state -> express in terms of ground state

-> Higgs boson

-> Gauge bosons with mass

-> Higgs-particle couplings terms —>

Higgs couplings  $\sim$  boson mass<sup>2</sup>  
Higgs couplings  $\sim$  fermion mass



# Higgs Parity

If coupling of Higgs boson to other particles does not change under CP


→ CP-even (scalar)

If all coordinates are flipped, like left and right are flipped in a normal mirror

→ Coupling is CP-odd (pseudoscalar)



# Spin and CP

Effect of spin on  $|\cos\theta^*|$  of the two photons



$$|\cos\theta^*| = \frac{|\sinh(\Delta\eta^{\gamma\gamma})|}{\sqrt{1 + (p_T^{\gamma\gamma}/m_{\gamma\gamma})^2}} \frac{2p_T^{\gamma_1} p_T^{\gamma_2}}{m_{\gamma\gamma}^2}$$

Variables can be defined in the  $H \rightarrow ZZ$  decay that are sensitive to spin and parity






# Higgs production modes at the LHC


Run 2 production rates shown (13 TeV,  $\sim 150 \text{ fb}^{-1}$ )

**Gluon fusion (ggF)**




**$\sim 8 \text{ M}$  events**

**Vector-boson fusion (VBF)**




**$\sim 600 \text{ k}$  events**

**Higgs-strahlung (VH)**



**$\sim 400 \text{ k}$  events**

**Higgs production  
in association with  $t\bar{t}$  (ttH)**





**$\sim 80 \text{ k}$  events**

# Higgs to $\tau\tau$

## $\tau$ leptons are complicated to reconstruct

- Various decay modes including neutrinos
- Analysis through statistical combination of variety of channels

The only lepton heavy enough to decay to hadrons




### 3 prong

$$\tau^\pm \rightarrow \pi^\pm \pi^\mp \pi^\pm \nu_\tau$$

+  $\geq 0$  neutral  $\pi$ 's

### Leptonic

$$\tau^\pm \rightarrow l^\pm \nu_l \nu_\tau$$



### 1 prong




$$\tau^\pm \rightarrow \pi^\pm \nu_\tau$$

+  $\geq 0$  neutral  $\pi$ 's




# Higgs to $\tau\tau$

Large backgrounds from  $Z \rightarrow \tau\tau + \text{jets}$



# Differential Higgs measurement


Enough Higgs candidates to perform **differential measurements** for variety of observables





# Di-Higgs production

Very recent ATLAS Run 2 combination:



Higgs-self coupling will be a key focus at the HL-LHC


# Putting it all together

## Almost all production modes established

**CMS**

Nature 607 (2022) 60-68



138 fb<sup>-1</sup> (13 TeV)





# Higgs mass measurements

- **Not predicted by SM**
- **Mass measurements in “golden channels”:  $H \rightarrow \gamma\gamma$  and  $H \rightarrow ZZ$**
- **Optimised analyses in categories with best mass resolution (photon, electron and muon energy response)**
- **Reached 0.09% precision**



# H → WW control region

Owing to spin correlations in the  $WW^{(*)}$  system arising from the spin-0 nature of the SM Higgs boson and the V-A structure of the  $W$  boson decay vertex, the charged leptons tend to emerge from the primary vertex pointing in the same direction [107]. This kinematic feature is exploited for all jet multiplicities by requiring that  $|\Delta\phi_{\ell\ell}| < 1.8$ , and the dilepton invariant mass,  $m_{\ell\ell}$ , be less than 50 GeV for the 0-jet and 1-jet channels. For the 2-jet channel, the  $m_{\ell\ell}$  upper bound is increased to 80 GeV.

## 6.2.3. $WW$ control sample

The MC predictions of the  $WW$  background in the 0-jet and 1-jet analyses, summed over lepton flavours, are normalised using control regions defined with the same selections as for the signal region except that the  $\Delta\phi_{\ell\ell}$  requirement is removed and the upper bound on  $m_{\ell\ell}$  is replaced with a lower bound:  $m_{\ell\ell} > 80$  GeV.