

Towards background rejection (Particle Identification)

M. Almanza Soto, S. Huang, A. Irles, J.P. Marquez
26/09/2024 LUXE ECAL Workshop - Krakow



UNIVERSITAT
DE
VALÈNCIA

IFIC
INSTITUT DE FÍSICA
CORPUSCULAR

CSIC
CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS



GENERALITAT
VALENCIANA
Conselleria de Educació,
Universitats y Empleo

GVA NEXT
Fondos Next Generation en la Comunitat Valenciana



Financiado por la
Unión Europea
NextGenerationEU

Plan de Recuperación,
Transformación
y Resiliencia

AITANA

Goals

Particle identification for background rejection in NPOD's ECAL

Particles to study:

- Pions
- Neutrons
- Photons


These neutral particles can mimic the expected signal photons in the ECAL, we have to be able to reject them.

The beam dump absorbs most background particles that could come from IP.


Basic principles of PID in High granularity CAL

- Different particles have different behaviors when interacting with the detector material
- EM vs Hadron showers in ECAL have different “shapes” and deposit energy differently


MUON






PION



PHOTON



NEUTRON




Examples of useful event parameters for PID


Making use of the high granularity

- Total number of hits (fired cells)
- Total energy
- Hits per layer
- Energy per layer
- Moliere radius
- Shower start layer (if there is a shower)
- Shower end layer
- Shower length
- Max layer reached
- Transversal distribution per layer and total


Previous work by J.P. Marquez




Previous work by Jesus + my work to use with CALORIMETERHIT




Previous work by Jesus + my work to use with CALORIMETERHIT




Output root file: neutron




drawing branch sume from ntp





Calculation of parameters from simulation: Photon example







Neutron vs photon: 10 GeV




Neutron 10 GeV vs photon 5 GeV



Moliere radius: Neutron vs photon 10 GeV


NEUTRON

drawing branch mol from ntp



PHOTON

drawing branch mol from ntp



Work to do

- Immediate
 - Produce samples for the three particle types for many energies to train ML algorithm
- Long run but not too long
 - Actually use J.P.'s machine learning program and get results/numbers for background rejection