

Parton evolution with α_s at small k_T

DGLAP evolution – solution with parton branching method

$$\bullet f(x, \mu^2) = f(x, \mu_0^2) \Delta_s(\mu^2) + \int^{z_M} \frac{dz}{z} \int \frac{d\mu'^2}{\mu'^2} \cdot \frac{\Delta_s(\mu^2)}{\Delta_s(\mu'^2)} P^{(R)}(z) f\left(\frac{x}{z}, \mu'^2\right)$$

$$f_0(x, \mu^2) = f(x, \mu_0^2) \Delta(\mu^2)$$

DGLAP evolution – solution with parton branching method

$$\bullet f(x, \mu^2) = f(x, \mu_0^2) \Delta_s(\mu^2) + \int^{z_M} \frac{dz}{z} \int \frac{d\mu'^2}{\mu'^2} \cdot \frac{\Delta_s(\mu^2)}{\Delta_s(\mu'^2)} P^{(R)}(z) f\left(\frac{x}{z}, \mu'^2\right)$$


$$f_0(x, \mu^2) = f(x, \mu_0^2) \Delta(\mu^2)$$

from μ' to μ
w/o branching



branching at μ'

from μ to μ'
w/o branching

$$f_1(x, \mu^2) = f(x, \mu_0^2) \Delta(\mu^2) + \int_{\mu_0^2}^{\mu^2} \frac{d\mu'^2}{\mu'^2} \frac{\Delta(\mu^2)}{\Delta(\mu'^2)} \int^{z_M} \frac{dz}{z} P^{(R)}(z) f(x/z, \mu_0^2) \Delta(\mu'^2)$$





Validation of method with QCDnum at **NLO**



- Very good agreement with **NLO** - QCDnum over all x and μ^2
- the same approach works also at NNLO !

Validation of method at **NLO**: z_M - dependence



- No dependence on z_M if z_M is large enough:
 - approximation is of
- Very good agreement with **NLO** - QCDnum

PDFs from Parton Branching method: fit to HERA data

- Convolution of kernel with starting distribution

$$\begin{aligned} x f_a(x, \mu^2) &= x \int dx' \int dx'' \mathcal{A}_{0,b}(x') \tilde{\mathcal{A}}_a^b(x'', \mu^2) \delta(x'x'' - x) \\ &= \int dx' \mathcal{A}_{0,b}(x') \cdot \frac{x}{x'} \tilde{\mathcal{A}}_a^b\left(\frac{x}{x'}, \mu^2\right) \end{aligned}$$

- Fit performed using xFitter frame (with collinear Coefficient functions at NLO)
 - using full HERA I+II inclusive DIS (neutral current, charged current) data
 - in total 1145 data points







$$3.5 \leq Q^2 \leq 50000 \text{ GeV}^2$$

$$4 \cdot 10^{-5} < x < 0.65$$





- using starting distribution as in HERAPDF2.0
 - $\chi^2/ndf = 1.2$

→ Can be easily extended to include any other measurement for fit !

Fits to DIS x-section at NLO: F_2 and F_2^C




Collinear parton distributions after fit



- fit 1 with $\alpha_s(\mu)$
 - $\mu = q$
 - as good as HERAPDF2.0
 $\chi^2/ndf = 1.2$
- fit 2 with $\alpha_s(\mu)$
 - $\mu = \max(1, q(1 - z))$
 - $\chi^2/ndf = 1.21$
- very different gluon distribution obtained at small Q^2





Analytic continuation of α_s into the non-pert region



- α_s with extension to small k_T region
- with taming parameter:





$$q_{eff}^2 = \frac{q^2}{q^2 + M^2}$$
- with analytic continuation ala A. Kotikov et al

Collinear parton distributions after fit with new α_s



- fit with tamed α_s :
 $\chi^2/ndf = 1.21$
- fit with analytic α_s :
 $\chi^2/ndf = 1.75$
 - not really good
 - Especially shape of gluon at starting scale is different !

Fits to DIS x-section at NLO: F_2



Conclusion

- Treatment of small k_T region in QCD fits very important, if scale in α_s is k_T dependent
 - with taming (instead of fixing) α_s already very good fit obtained
 - with analytically continued α_s - difficulty to fit F2 with good chi2
 - shape of gluon at small scales very different

Appendix