

Follow up on Searches for SUSY with 2 leptons and missing transverse energy with the ATLAS experiment




Joao Firmino da Costa
LHC-DESY discussions



Dilepton searches Results from ATLAS

Final state leptons can be produced in different ways...




a) $\tilde{\chi}_i^0 \rightarrow l^\pm \nu \tilde{\chi}_j^\mp$

b) $\tilde{\chi}_i^\pm \rightarrow l^\pm \nu \tilde{\chi}_j^0$

c) $\tilde{\chi}_i^0 \rightarrow l^\pm l^\mp \tilde{\chi}_j^0$

d) $\tilde{\chi}_i^\pm \rightarrow l^\pm l^\mp \tilde{\chi}_j^\pm$




(c) and (d) on a single-leg (opposite-sign)
(a) or (b) on both-legs (opposite-sign or same-sign)

Dilepton final states (opposite sign or same sign)

latest results with 1.04 fb⁻¹ (public)
(for 2010 results : EPJC(2011) 1682)

Opposite-Sign Flavour subtraction search

Use the selected opposite-sign di-leptons and search for excess of same-flavour events over opposite-flavour




Offers a great to measure SUSY mass parameters
Via the measurement of mass edges

Most important bkg (ttbar) are combinatorics so they cancel with the Flavour Subtraction

Quantify the excess of Same Flavour (SF) through the usage of the variable S :

$$S = \underbrace{\frac{N(e^\pm e^\mp)}{\beta(1 - (1 - \tau_e)^2)}}_{\text{SF}} - \underbrace{\frac{N(e^\pm \mu^\mp)}{1 - (1 - \tau_e)(1 - \tau_\mu)}}_{\text{DF}} + \underbrace{\frac{\beta N(\mu^\pm \mu^\mp)}{(1 - (1 - \tau_\mu)^2)}}_{\text{SF}}$$



measured with in Z CR and variation due to process from MC

Opposite-Sign Flavour subtraction search - Signal Regions

3 signal regions :

- 1) $E_{\text{miss}} > 80 \text{ GeV}$ and Z peak veto (80-100)
- 2) $E_{\text{miss}} > 80 \text{ GeV}$, jets $p_T > 20, 20 \text{ GeV}/c$
- 3) $E_{\text{miss}} > 250 \text{ GeV}$


Same sources of backgrounds as OS analysis:

$t\bar{t}$ bar estimation from m_{CT} (for SR3)
or from MC (SR1, SR2)

Z CR : Z peak, low m_{ll} , extrapolate
back to SR using MC

fake leptons : ABCD matrix method

Wt and dibosons : MC estimations



Opposite-Sign Flavour subtraction search - Results

- Determine $\mathcal{S}$ ($\mathcal{S}_b$) done with MC toy experiments:
 - ▶ toy experiment for each SM process in each channel
 - ▶ based on data driven expectations nominal values and uncertainties
- observed $\mathcal{S}_b$ in agreement with SM pred. within 1σ (FS-SR2 within 1.5σ)


	$\mathcal{S}_{obs}$	$\bar{\mathcal{S}}_b$	RMS
FS-SR1	$131.6 \pm 0.6(\text{sys})$	$126.5 \pm 23.5 \pm 17.2$	49.9
FS-SR2	$142.2 \pm 0.6(\text{sys})$	$70.0 \pm 23.2 \pm 16.8$	49.1
FS-SR3	$-3.1 \pm 0.0(03)(\text{sys})$	$0.4 \pm 1.2 \pm 1.2$	4.6

errors on $\bar{\mathcal{S}}_b$ are stat. and syst. errors

Limits:

- Probability of observing a value $\mathcal{S} > \mathcal{S}_{obs}$ evaluated
- Limit set on mean contribution of potential signal ($\bar{\mathcal{S}}_s$)
- assumptions for signal events which are added to the toys as signal:
 - ▶ same BR for ee and $\mu\mu$ in SUSY events and NO contribution of $e\mu$

	$\mathcal{S} > \mathcal{S}_{obs}$ (%)	Limit $\bar{\mathcal{S}}_s$ (95% C.L.)
FS-SR1	46	88
FS-SR2	7	156
FS-SR3	77	4.9



Discussions

Discussion – Experiment vs Theory

CMS provides also limits on the excess of same flavor events dependent on the mass edge position






Figure 6: CL_s 95% confidence level upper limit on cross-section times acceptance as a function of the endpoint in the invariant mass spectrum assuming a triangular shaped signal.

How useful can these nice results be to theorists ?

Discussion - Experiment vs Theory

2 signal regions :

- 1) $E_{\text{miss}} > 80 \text{ GeV}$ and Z peak veto (80-100)
- 2) $E_{\text{miss}} > 80 \text{ GeV}$, jets $p_T > 20, 20 \text{ GeV}/c$



What do you need as information for your fits/models ?
Above + efficiency maps or so ?

Discussion - Experiment vs Theory

Questions :

1 - For fits using shapes (fS), what is the most general template one can use ?

We were thinking of using our models to get a triangular shape, (similar to CMS) but maybe you have another recipe ?

2 - Besides mSUGRA, what other models can we look at ? Suggestions ?

3 - What else with 2 lepton analyses ? What are we not covering ?

Back-up

Same-Sign dilepton search - Results

	Background	Obs.	95% C.L.
SS-SR1	$32.6 \pm 4.4 \pm 4.4$	25	10.2 fb
SS-SR2	$24.9 \pm 4.1 \pm 6.6$	28	20.3 fb

Same Sign [SS-SR1]	$e^\pm e^\pm$	$e^\pm \mu^\pm$	$\mu^\pm \mu^\pm$
Fake	$3.5 \pm 0.7 \pm 2.0$	$14.4 \pm 3.2 \pm 0.6$	$9.2 \pm 4.0 \pm 1.8$
Charge flip	$0.73 \pm 0.05 \pm 0.06$	$1.10 \pm 0.07 \pm 0.08$	<i>neg.</i>
Dibosons	$0.79 \pm 0.27 \pm 0.05$	$1.7 \pm 0.4 \pm 0.3$	$1.1 \pm 0.2 \pm 0.1$
Standard Model	$5.0 \pm 0.8 \pm 2.0$	$17.2 \pm 3.2 \pm 0.6$	$10.3 \pm 3.0 \pm 1.8$
Cosmics	$< 10^{-3}$	$< 10^{-3}$	$< 10^{-3}$
Observed	6	14	5

Same Sign [SS-SR2]	$e^\pm e^\pm$	$e^\pm \mu^\pm$	$\mu^\pm \mu^\pm$
Fake	$1.5 \pm 0.5 \pm 2.0$	$13.4 \pm 3.2 \pm 2.7$	$6.7 \pm 2.5 \pm 1.8$
Charge flip	$0.59 \pm 0.04 \pm 0.04$	$1.36 \pm 0.05 \pm 0.06$	<i>neg.</i>
Dibosons	$0.25 \pm 0.06 \pm 0.13$	$0.9 \pm 0.2 \pm 0.2$	$0.64 \pm 0.05 \pm 0.02$
Standard Model	$2.4 \pm 0.5 \pm 2.0$	$15.6 \pm 3.2 \pm 2.7$	$6.9 \pm 2.5 \pm 1.8$
Cosmics	$< 10^{-3}$	$< 10^{-3}$	$< 10^{-3}$
Observed	4	14	10

fakes are dominant contribution


Opposite-Sign dilepton search - Results

	Background	Obs.	95% C.L.
OS-SR1	$15.5 \pm 1.2 \pm 4.4$	13	9.5 fb
OS-SR2	$13.0 \pm 1.8 \pm 4.1$	17	15.2 fb
OS-SR3	$5.7 \pm 1.1 \pm 3.5$	2	5.0 fb

Opposite Sign [OS-SR1]	$e^{\pm}e^{\mp}$	$e^{\pm}\mu^{\mp}$	$\mu^{\pm}\mu^{\mp}$
$t\bar{t}$	$1.84 \pm 0.07 \pm 0.49$	$5.09 \pm 0.19 \pm 1.35$	$3.34 \pm 0.13 \pm 0.88$
$Z/\gamma^* + \text{jets}$	$0.013 \pm 0.008 \pm 0.67$	$1.03 \pm 0.42 \pm 0.02$	$0.81 \pm 0.26 \pm 0.06$
Fakes	$0.17 \pm 0.19 \pm 0.36$	$0.92 \pm 0.96 \pm 1.15$	$-0.08 \pm 0.03 \pm 0.003$
Dibosons	$0.54 \pm 0.29 \pm 0.07$	$0.04 \pm 0.03 \pm 0.03$	$0.67 \pm 0.25 \pm 0.31$
Single-top	$0.11 \pm 0.11 \pm 0.05$	$0.47 \pm 0.16 \pm 0.16$	$0.48 \pm 0.17 \pm 0.09$
Standard Model	$2.67 \pm 0.37 \pm 1.29$	$7.55 \pm 1.07 \pm 2.13$	$5.30 \pm 0.42 \pm 1.29$
Cosmics	$< 10^{-3}$	$< 10^{-3}$	$< 10^{-3}$
Observed	2	8	3
Opposite Sign [OS-SR2]	$e^{\pm}e^{\mp}$	$e^{\pm}\mu^{\mp}$	$\mu^{\pm}\mu^{\mp}$
$t\bar{t}$	$1.41 \pm 0.11 \pm 0.33$	$3.90 \pm 0.30 \pm 0.95$	$2.56 \pm 0.20 \pm 0.62$
$Z/\gamma^* + \text{jets}$	$0.45 \pm 0.23 \pm 0.44$	$0.84 \pm 0.59 \pm 0.32$	$0.27 \pm 0.14 \pm 0.27$
Fakes	$0.01 \pm 0.14 \pm 0.19$	$2.77 \pm 1.64 \pm 2.11$	$-0.13 \pm 0.04 \pm 0.01$
Dibosons	<i>neg.</i>	$0.03 \pm 0.03 \pm 0.03$	$0.24 \pm 0.21 \pm 0.02$
Single-top	$0.05 \pm 0.10 \pm 0.02$	$0.39 \pm 0.16 \pm 0.25$	$0.09 \pm 0.15 \pm 0.08$
Standard Model	$1.92 \pm 0.31 \pm 0.77$	$7.93 \pm 1.77 \pm 1.41$	$3.16 \pm 0.36 \pm 0.94$
Cosmics	$< 10^{-3}$	$< 10^{-3}$	$< 10^{-3}$
Observed	3	9	5
Opposite Sign [OS-SR3]	$e^{\pm}e^{\mp}$	$e^{\pm}\mu^{\mp}$	$\mu^{\pm}\mu^{\mp}$
$t\bar{t}$	$0.77 \pm 0.14 \pm 0.51$	$2.14 \pm 0.38 \pm 1.41$	$1.40 \pm 0.25 \pm 0.92$
$Z/\gamma^* + \text{jets}$	$0.01 \pm 0.01 \pm 0.17$	<i>neg.</i>	$0.27 \pm 0.20 \pm 0.47$
Fakes	$0.13 \pm 0.13 \pm 0.03$	$0.91 \pm 0.94 \pm 0.04$	$-0.03 \pm 0.02 \pm 0.001$
Single-top	<i>neg.</i>	$0.0 \pm 0.0 \pm 0.02$	$0.10 \pm 0.10 \pm 0.05$
Standard Model	$0.91 \pm 0.19 \pm 0.67$	$3.05 \pm 1.01 \pm 1.43$	$1.77 \pm 0.34 \pm 1.41$
Cosmics	$< 10^{-3}$	$< 10^{-3}$	$< 10^{-3}$
Observed	0	1	1

$t\bar{t}$ is dominant contribution

$t\bar{t}$ Background (OS): Contranverse Mass Tagger



- In the decay of a two pair-produced heavy states δ which decay via $\delta \rightarrow \alpha\chi_i$:

$$m_{CT}^2(\chi_1, \chi_2) = [E_T(\chi_1) + E_T(\chi_2)]^2 - [\mathbf{p}_T(\chi_1) - \mathbf{p}_T(\chi_2)]^2$$

- m_{CT} distributions have endpoints defined by $m(\delta)$, $m(\alpha)$ and the vector sum of transverse momenta of the visible particles upstream of the system for which the contranverse mass is calculated (p_b)
- For the $t\bar{t}$ system $m_{CT}(\ell\ell)$, $m_{CT}(jj)$, $m_{CT}(j\ell, j\ell)$ can be constructed

Contranverse mass tagger

1. Event with least 2 jets with $p_T > 20$ GeV
2. Consider all 2 jet permutations j_1, j_2 , such that the two jets have $p_T > 20$ GeV and $p_T(j_1) + p_T(j_2) + p_T(\ell_1) + p_T(\ell_2) > 100$ GeV
3. $m_{CT}(\ell_1, \ell_2)$ in the allowed area of the $(m_{CT}(\ell_1, \ell_2), p_b(\ell\ell))$ plane
4. Build all pairs $((j_i\ell_1)(j_j\ell_2))$ such that $m(j_i\ell_1) < 155$ GeV and $m(j_j\ell_2) < 155$ GeV
5. One combination with $m_{CT}(jj)$ in the allowed area of the $m_{CT}(jj), p_b(jj)$ plane
6. $m_{CT}(j\ell, j\ell)$ should be compatible with $t\bar{t}$

- m_{CT} tagger has an efficiency of 85% in signal region
- Control region for $t\bar{t}$ estimation:
 - m_{CT} -tagged events
 - $60 < E_T^{\text{miss}} < 80$ GeV