

Flavor Violating Higgs Decays

Joachim Kopp

DESY Theory Workshop 2012
September 27, 2012



MAX-PLANCK-GESELLSCHAFT

Based on work done in collaboration with
Roni Harnik and Jure Zupan
[arXiv:1209.1397](https://arxiv.org/abs/1209.1397)



MAX-PLANCK-INSTITUT
FÜR KERNPHYSIK

Outline

- 1 Flavor mixing in the Higgs sector
- 2 Couplings to leptons
- 3 Couplings to quarks
- 4 Flavor-violating Higgs decays at the LHC
- 5 Summary



Flavor Mixing in the Higgs Sector

Motivation

Scenario 1: Several sources of EW symmetry breaking

... can lead to misalignment of mass matrices and light (125 GeV) Higgs Yukawa couplings

Example: Type III 2-Higgs Doublet model

Scenario 2: Extra Higgs couplings

Assume existence of heavy new particles, which induce effective operators of the form

$$\Delta \mathcal{L}_Y = -\frac{\lambda'_{ij}}{\Lambda^2} (\bar{f}_L^i f_R^j) H (H^\dagger H) + h.c. + \dots ,$$

→ after EWSB, new (but misaligned) contributions to mass matrices and Yukawa couplings

Effective Yukawa Lagrangian

Effective Yukawa Lagrangian

$$\mathcal{L}_Y = -m_i \bar{f}_L^i f_R^i - Y_{ij}^a (\bar{f}_L^i f_R^j) h^a + h.c. + \dots$$

Previously studied by many authors:

- | | |
|--|---|
| Bjorken Weinberg, PRL 38 (1977) 622 | McWilliams Li, Nucl. Phys. B 179 (1981) 62 |
| Shanker, Nucl. Phys. B 206 (1982) 253 | Barr Zee, PRL 65 (1990) 21 |
| Babu Nandi, hep-ph/9907213 | Diaz-Cruz Toscano, hep-ph/9910233 |
| Han Marfatia, hep-ph/0008141 | Kanemura Ota Tsumura, hep-ph/0505191 |
| Blanke Buras Duling Gori Weiler, arXiv:0809.1073 | |
| Casagrande Goertz Haisch Neubert Pfoh, arXiv:0807.4937 | |
| Giudice Lebedev, arXiv:0804.1753 | Aguilar-Saavedra, arXiv:0904.2387 |
| Albrecht Blanke Buras Duling Gemmeler, arXiv:0903.2415 | |
| Buras Duling Gori, arXiv:0905.2318 | Azatov Toharia Zhu, arXiv:0906.1990 |
| Agashe Contino, arXiv:0906.1542 | Davidson Greiner, arXiv:1001.0434 |
| Goudelis Lebedev Park, arXiv:1111.1715 | Blankenburg Ellis Isidori, arXiv:1202.5704 |
| Arribi Cheng Kong, arXiv:1208.4669 | McKeen Pospelov Ritz, arXiv:1208.4597 |
| ... | |

Effective Yukawa Lagrangian

Effective Yukawa Lagrangian

$$\mathcal{L}_Y = -m_i \bar{f}_L^i f_R^i - Y_{ij}^a (\bar{f}_L^i f_R^j) h^a + h.c. + \dots$$


New in this talk:

- Comprehensive list of up-to-date constraints (including subdominant ones)
- First LHC limits
- Strategy for future LHC searches




Couplings to Leptons


Low-energy constraints on LFV in the Higgs sector




$M-\bar{M}$ oscillations






μ - e conversion



$g - 2$, EDMs




$\tau \rightarrow 3\mu, \mu\mu e, \text{etc.}$





$\tau \rightarrow \mu\gamma$

Constraints on $h \rightarrow \mu e$



Constraints on $h \rightarrow \tau\mu$ and $h \rightarrow \tau e$



Substantial flavor violation ($\text{BR}(h \rightarrow \tau\mu, \tau e) \sim 0.01$) perfectly viable.



Couplings to Quarks

Constraints on Higgs couplings to light quarks

- Tight constraints from neutral meson oscillations
- Work in Effective Field Theory:

$$H_{\text{eff}} = C_2^{db} (\bar{b}_R d_L)^2 + \tilde{C}_2^{db} (\bar{b}_L d_R)^2 + C_4^{db} (\bar{b}_L d_R)(\bar{b}_R d_L)$$

- Wilson coefficients constrained in Bona et al., arXiv:0707.0636

Technique	Coupling	Constraint
D^0 oscillations	$ Y_{uc} ^2, Y_{cu} ^2$	$< 5.0 \times 10^{-9}$
	$ Y_{uc} Y_{cu} $	$< 7.5 \times 10^{-10}$
B_d^0 oscillations	$ Y_{db} ^2, Y_{bd} ^2$	$< 2.3 \times 10^{-8}$
	$ Y_{db} Y_{bd} $	$< 3.3 \times 10^{-9}$
B_s^0 oscillations	$ Y_{sb} ^2, Y_{bs} ^2$	$< 1.8 \times 10^{-6}$
	$ Y_{sb} Y_{bs} $	$< 2.5 \times 10^{-7}$
K^0 oscillations	$\Re(Y_{ds}^2), \Re(Y_{sd}^2)$	$[-5.9 \dots 5.6] \times 10^{-10}$
	$\Im(Y_{ds}^2), \Im(Y_{sd}^2)$	$[-2.9 \dots 1.6] \times 10^{-12}$
	$\Re(Y_{ds}^* Y_{sd})$	$[-5.6 \dots 5.6] \times 10^{-11}$
	$\Im(Y_{ds}^* Y_{sd})$	$[-1.4 \dots 2.8] \times 10^{-13}$

Couplings involving top quarks

Loose constraints ($|Y_{qt}|, |Y_{tq}| < \text{few} \times 10^{-1}$) from

- Single top production

CDF arXiv:0812.3400

DØ arXiv:1006.3575

ATLAS arXiv:1203.0529

- $t \rightarrow hq$

Craig et al. arXiv:1207.6794

based on CMS multilepton search arXiv:1204.5341

Not sensitive

- $t \rightarrow Zq$

CMS arXiv:1208.0957



Flavor-Violating Higgs Decays at the Large Hadron Collider

$h \rightarrow \tau\mu$ and $h \rightarrow \tau e$ at the LHC

Basic idea:

- $h \rightarrow \tau\ell$ has the same final state as $h \rightarrow \tau\tau_\ell$
(but is enhanced by $1/\text{BR}(\tau \rightarrow \ell)$)
- Recast $h \rightarrow \tau\tau$ search
here: ATLAS, arXiv:1206.5971
- Use VBF cuts
(much lower BG than gg fusion)

$h \rightarrow \tau\mu$ and $h \rightarrow \tau e$ at the LHC

Basic idea:

- $h \rightarrow \tau\ell$ has the same final state as $h \rightarrow \tau\tau_\ell$
(but is enhanced by $1/\text{BR}(\tau \rightarrow \ell)$)
- Recast $h \rightarrow \tau\tau$ search
here: ATLAS, arXiv:1206.5971
- Use VBF cuts
(much lower BG than gg fusion)

Technicalities:

- Consider only 2-lepton final states here
- Use MadGraph 5, v1.4.6,
Pythia 64, PGS
- Use only 120–160 GeV bin
- Derive one-sided 95% CL limit


$h \rightarrow \tau\mu$ and $h \rightarrow \tau e$ at the LHC

Basic idea:

- $h \rightarrow \tau\ell$ has the same final state as $h \rightarrow \tau\tau_\ell$
(but is enhanced by $1/\text{BR}(\tau \rightarrow \ell)$)
- Recast $h \rightarrow \tau\tau$ search
here: ATLAS, arXiv:1206.5971
- Use VBF cuts
(much lower BG than gg fusion)

Technicalities:

- Consider only 2-lepton final states here
- Use MadGraph 5, v1.4.6,
Pythia 64, PGS
- Use only 120–160 GeV bin
- Derive one-sided 95% CL limit




$h \rightarrow \tau\mu$ and $h \rightarrow \tau e$ at the LHC

Basic idea:

- $h \rightarrow \tau\ell$ has the same final state as $h \rightarrow \tau\tau_\ell$
(but is enhanced by $1/\text{BR}(\tau \rightarrow \ell)$)
- Recast $h \rightarrow \tau\tau$ search
here: ATLAS, arXiv:1206.5971
- Use VBF cuts
(much lower BG than gg fusion)

Technicalities:



- Consider only 2-lepton final states here
- Use MadGraph 5, v1.4.6,
Pythia 64, PGS
- Use only 120–160 GeV bin
- Derive one-sided 95% CL limit





Result:

$$\begin{aligned}\text{BR}(h \rightarrow \tau\mu) &< 0.13 \\ \text{BR}(h \rightarrow \tau e) &< 0.13\end{aligned}$$

Constraints on $h \rightarrow \tau\mu$ and $h \rightarrow \tau e$



Constraints on $h \rightarrow \tau\mu$ and $h \rightarrow \tau e$



Strategy for a dedicated $h \rightarrow \tau\mu$ and $h \rightarrow \tau e$ search


Possible improvements:

- Different invariant mass formula
(assuming 1 neutrino rather than 3)
 - ▶ Avoids smearing of signal
 - ▶ Shifts $Z \rightarrow \tau\tau$ peak to lower invariant mass
- Consider hadronic τ 's
(especially for CMS)
- Modified cuts
 - ▶ CMS $h \rightarrow \tau_{\text{had}}\tau\ell$ search requires $m_T(\ell, \not{p}_T) < 40$ GeV to suppress $W + \text{jets}$
 - ▶ In $h \rightarrow \tau_{\text{had}}\mu$, neutrino and muon typically not collinear
→ large $m_T(\ell, \not{p}_T)$



Strategy for a dedicated $h \rightarrow \tau\mu$ and $h \rightarrow \tau e$ search

Possible improvements:

- Different invariant mass formula (assuming 1 neutrino rather than 3)
 - ▶ Avoids smearing of signal
 - ▶ Shifts $Z \rightarrow \tau\tau$ peak to lower invariant mass
- Consider hadronic τ 's (especially for CMS)
- Modified cuts
 - ▶ CMS $h \rightarrow \tau_{\text{had}}\tau e$ search requires $m_T(\ell, \not{p}_T) < 40$ GeV to suppress $W + \text{jets}$
 - ▶ In $h \rightarrow \tau_{\text{had}}\mu$, neutrino and muon typically not collinear
→ large $m_T(\ell, \not{p}_T)$



Strategy for a dedicated $h \rightarrow \tau\mu$ and $h \rightarrow \tau e$ search



For $Y_{\mu\tau}$, $Y_{\tau\mu}$ close to the current upper limits, **spectacular signals** possible.

Summary

- Flavor-violating Higgs couplings arise in
 - ▶ Models with several sources of electroweak symmetry breaking
 - ▶ Models with heavy fields coupled to the Higgs
- In the lepton sector:
 - ▶ Constraints from $\ell_1 \rightarrow \ell_2 + \gamma$, $\ell_1 \rightarrow \ell_2 + X$, $\mu\text{--}e$ conversion in nuclei, $g - 2$, EDMs, $M - \bar{M}$ oscillations
 - ▶ Strong constraints in the $\mu\text{--}e$ sector
 - ▶ Very weak constraints in the $\tau\text{--}e$ and $\tau\text{--}\mu$ sectors
- In the quark sector:
 - ▶ Strong constraints on couplings to light quarks
 - ▶ Very weak constraints on couplings to top quarks
- At the LHC
 - ▶ Constraints on anomalous top–Higgs couplings from single top production
 - ▶ A recast ATLAS $h \rightarrow \tau_\ell \tau_\ell$ search already provides strongest limits on $h \rightarrow \tau \mu$ and $h \rightarrow \tau e$
 - ▶ A dedicated search would be much more sensitive



Thank you!