

J/ψ photoproduction in Pb-Pb and p-Pb ultra-peripheral collisions with ALICE at LHC

D. De Gruttola* for ALICE Collaboration

*Centro Fermi Roma and Salerno INFN - Italy




Summary

- ✓ LHC as γ Pb and γ p collider (Ultra-peripheral collisions)
- ✓ Physics motivation (gluon distribution in nuclei and nucleons)
- ✓ ALICE and UPCs (detector and trigger description)
- ✓ J/ψ cross section (forward and mid-rapidity)
- ✓ results and comparison with models (gluon shadowing)
- ✓ $\gamma\gamma$ cross section (constraint on QED processes)
- ✓ first results in pA (proton as a target)
- ✓ conclusions (achieved results and on going analyses)

LHC as γ Pb and γ p collider

- ✓ at the LHC heavy ions are accelerated towards each other at ultra relativistic energies
- ✓ being charged particles, they are accompanied by an **electromagnetic field**
- ✓ the EM field can be viewed as a flux of **quasi-real photons**
- ✓ intensity of the photon beam proportional to Z^2
- ✓ photon flux well described in **Fermi-Weizsäcker-Williams** approximation
- ✓ hadronic processes strongly suppressed
- ✓ high σ for γ -induced reactions e.g. **vector meson photoproduction**



- ✓ **virtuality** of the photon dependent on the radius of the emitting particle:

$$Q^2 \approx \left(\frac{\hbar c}{R} \right)^2$$

$$\gamma \text{ from p} \rightarrow Q^2 \approx (250 \text{ MeV})^2$$

$$\gamma \text{ from Pb} \rightarrow Q^2 \approx (30 \text{ MeV})^2$$


Physics motivation

- ✓ possibility to study non linear effects at **low x** in the gluon distribution of the target
- ✓ quarkonia **photo-production** allows to study the gluon density $G(x, Q^2)$ in Pb

$$\left. \frac{d\sigma(\gamma N \rightarrow VN)}{dt} \right|_{t=0} \approx \frac{\alpha_s \Gamma_{ee}}{3\alpha_e M_V^5} 16\pi^3 \left(xG(x, Q^2) \right)^2$$

- ✓ Bjorken-x accessible at LHC $x = (M_V/\sqrt{s_{NN}})\exp(\pm y) \sim 10^{-2} - 10^{-5}$
- ✓ vector meson photo-production as tool to measure **nuclear gluon shadowing** and **saturation**

$$R_g^A(x, Q^2) = \frac{G_A(x, Q^2)}{G_p(x, Q^2)}$$




C A Salgado et al 2012 J. Phys. G.: Nucl. Part. Phys. **39** 015010

Physics motivation

- ✓ some representative values for γA cms energy $W_{\gamma A}$:

- ✧ $y = 0 \rightarrow W_{\gamma A} \sim 90$ GeV
- ✧ $y = -3 \rightarrow W_{\gamma A} \sim 400$ or 20 GeV
(the flux of photons at 400 GeV is $\sim 4\%$ of the flux at 20 GeV)

LHeC Study group ArXiv: 1211.4831




- ✓ **coherent** vector meson production:


- ✧ photon couples coherently to all nucleons
- ✧ $\langle p_T \rangle \sim 1/R_{Pb} \sim 60$ MeV/c
- ✧ no neutron emission in $\sim 80\%$ of cases

- ✓ **incoherent** vector meson production:

- ✧ photon couples to a single nucleon
- ✧ $\langle p_T \rangle \sim 1/R_p \sim 500$ MeV/c
- ✧ target nucleus normally breaks up





- ✓ an interesting physics case is also $\gamma\gamma$ interactions to provide informations on QED processes when $\sqrt{\alpha}$ is replaced by $Z\sqrt{\alpha}$



ALICE layout

Central Barrel
2 π tracking & PID
 $|\eta| < 1$

muon spectrometer
 $-2.5 > \eta > -4$




ACORDE (cosmics)
VZERO scint. (centrality)
 $-1.7 < \eta < -3.7, 2.8 < \eta < 5.1$
T0 (timing)
ZDC (centrality)
FMD (N_{ch} $-3.4 < \eta < 5$)
PMD (N_{γ}, N_{ch})

Detector:
Length: 26 meters
Height: 16 meters
Weight: 10,000 tons

ALICE and UPCs ($J/\psi \rightarrow \mu^+\mu^-$)

UPC forward trigger:

- ✧ single **muon trigger** with $p_T > 1$ GeV/c ($-4 < \eta < -2.5$)
- ✧ hit in **VZERO-C** ($-3.7 < \eta < -1.7$)
- ✧ no hits in **VZERO-A** ($2.8 < \eta < 5.1$)




integrated luminosity $\sim 55 \mu\text{b}^{-1}$

✓ offline event selection:

- ✧ beam gas rejection with VZERO
- ✧ hadronic rejection with ZDC and SPD

✓ track selection:


- ✧ muon tracks: $-3.7 < \eta < -2.5$
- ✧ matching with the trigger
- ✧ radial position for muons at the end of absorber: $17.5 < R_{\text{abs}} < 89.5$ cm
- ✧ p_T dependent DCA cut
- ✧ opposite sign dimuon: $-3.6 < y < -2.6$



ALICE and UPCs ($J/\psi \rightarrow \mu^+\mu^-$ and $J/\psi \rightarrow e^+e^-$)

UPC mid-rapidity trigger:


- ✧ ≥ 2 hits in **SPD**
- ✧ $2 \leq \text{TOF hits} \leq 6$ and back-to-back topology
- ✧ veto on **VZERO-C** and **VZERO-A**




integrated luminosity $\sim 23 \mu\text{b}^{-1}$

✓ offline event selection:


- ✧ rejection with VZERO and FMD
- ✧ primary vertex
- ✧ $\max(p_{T1}, p_{T2}) > 1 \text{ GeV}/c$
- ✧ dE/dx consistent with e/μ
- ✧ opposite sign tracks
- ✧ ZDC cut on number of neutrons emitted in coherent events




ALICE and UPCs



central Pb-Pb collision



UP Pb-Pb collision at mid-rapidity



UP Pb-Pb collision at forward rapidity



two tracks in an otherwise empty detector

detailed studies done to understand the noise and the emptiness of the detector

J/ψ measurements (coherent at forward rapidity)

first measurement of J/ψ photo-production done at LHC

Phys. Lett. B718 (2013) 1273 -1283



p_T distribution fitted using MC samples representing several components:

- ✧ coherent and incoherent J/ψ
- ✧ ψ' feed down
- ✧ $\gamma\gamma \rightarrow \mu^+\mu^-$

distribution peaked at low momentum as expected from coherent production




J/ψ photo-production probes the gluon distribution in Pb at $x \sim 10^{-2}$

J/ψ measurements (coherent at mid-rapidity)

dimuon channel

arXiv:1305.1467 [nucl-ex] submitted to EPJ-C

electrons



muons

p_T distribution fitted using MC samples representing several components:

- ✧ coherent and incoherent J/ψ
- ✧ (coherent and incoherent) ψ' feed down
- ✧ $\gamma\gamma \rightarrow \mu^+\mu^-$
- ✧ hadronic

$p_T < 200$ MeV/c and < 6 neutrons emitted by nuclei

distribution peaked at low momentum as expected from coherent production




J/ψ photo-production probes the gluon distribution in Pb at $x \sim 10^{-3}$

J/ψ measurements (coherent at mid-rapidity)

dielectron channel

arXiv:1305.1467 [nucl-ex]

electrons



muons

p_T distribution fitted using MC samples representing several components:


- ✧ coherent and incoherent J/ψ
- ✧ (coherent and incoherent) ψ' feed down
- ✧ $\gamma\gamma \rightarrow e^+e^-$
- ✧ hadronic

$p_T < 300$ MeV/c and < 6 neutrons emitted by nuclei

distribution peaked at low momentum as expected from coherent production

J/ψ photo-production probes the gluon distribution in Pb at $x \sim 10^{-3}$

Results and comparison with models



arXiv:1305.1467 [nucl-ex]

$$|y| < 0.9 \rightarrow d\sigma_{J/\psi}^{coh} / dy = 2.38_{-0.24}^{+0.34} (stat + syst) \text{ mb}$$

$$-3.6 < y < -2.6 \rightarrow d\sigma_{J/\psi}^{coh} / dy = 1.00 \pm 0.18 (stat)^{+0.24}_{-0.26} (syst) \text{ mb}$$

measured cross section in good agreement with the calculation using the **EPS09** nuclear gluon prediction

✓ **AB:** Adeluyi and Bertulani, PRC85 (2012) 044904
these models use LO pQCD scaled by an effective constant to correct for missing contributions
MSTW08 assumes no nuclear effects, EPS08/09 incorporate nuclear effects according to different parametrizations

✓ **CSS:** Cisek, Szczurek, Sch.fer PRC86 (2012) 014905
color dipole model based on unintegrated gluon distribution of the proton

✓ **STARLIGHT:** Klein, Nystrand PRC60 (1999) 01493
GVDM coupled to a Glauber approach and using HERA data to fix the γp cross section

✓ **GM:** Goncalves, Machado, PRC84 (2011) 011902
color dipole model, where the dipole nucleon cross section is from the IIM saturation model

✓ **RSZ:** Rebyakova, Strikman, Zhalov, PLB 710 (2012) 252
based on LO pQCD amplitude for two gluon exchange where the gluon density incorporates shadowing computed in leading twist approximation




data are closer to models incorporating **nuclear gluon shadowing**

J/ψ measurements (incoherent at mid-rapidity)

dimuon channel

arXiv:1305.1467 [nucl-ex]

electrons



muons

p_T distribution fitted using MC samples representing several components:

- ✧ coherent and incoherent J/ψ
- ✧ (coherent and incoherent) ψ' feed down
- ✧ $\gamma\gamma \rightarrow \mu^+\mu^-$
- ✧ hadronic




$p_T > 200 \text{ MeV/c}$

the ratio $\sigma_{\text{inc}}/\sigma_{\text{coh}}$ provides further constraints on the treatment of the nuclear modifications implemented in the different models

J/ψ measurements (incoherent at mid-rapidity)

dielectron channel

arXiv:1305.1467 [nucl-ex]



p_T distribution fitted using MC samples representing several components:

- ✧ coherent and incoherent J/ψ
- ✧ (coherent and incoherent) ψ' feed down
- ✧ $\gamma\gamma \rightarrow e^+e^-$
- ✧ hadronic

$p_T > 300 \text{ MeV/c}$

the ratio $\sigma_{\text{inc}}/\sigma_{\text{coh}}$ provides further constraints on the treatment of the nuclear modifications implemented in the different models

Results and comparison with models


arXiv:1305.1467 [nucl-ex]

$$|y| < 0.9 \rightarrow d\sigma_{J/\psi}^{inc} / dy = 0.98^{+0.19}_{-0.17} (stat + syst) \text{ mb}$$

✧ none of the three existing models predicts the incoherent cross section correctly

✧ **STARLIGHT** predicts a correct incoherent-to-coherent ratio (0.41)

✧ ALICE measurement $0.41^{+0.10}_{-0.08} (stat + syst)$



✓ **STARLIGHT**: Klein, Nystrand PRC60 (1999) 01493
GVDM coupled to a Glauber approach and using HERA data to fix the γp cross section


✓ **RSZ**: Rebyakova, Strikman, Zhalov, PLB 710 (2012) 252
based on LO pQCD amplitude for two gluon exchange where the gluon density incorporates shadowing computed in leading twist approximation

✓ **LM**: Lappi, Mantysaari, PRC87 (2013) 032201
color dipole model based with Glauber approach and a saturation prescription

the ratio $\sigma_{inc}/\sigma_{coh}$ provides further constraints on the treatment of the nuclear modifications implemented in the different models

$\gamma\gamma$ cross section


arXiv:1305.1467 [nucl-ex]



- ✓ the $\gamma\gamma$ cross section measurement provides important constraints on QED calculations when the vertex $\sqrt{\alpha}$ has to be replaced by $Z\sqrt{\alpha}$
- ✓ due to the large Pb charge, giving $Z\sqrt{\alpha} \sim 0.6$, the inclusion of higher order terms is not straightforward \rightarrow the models including higher order terms predict a reduction of the cross section up to 30%
 - ✧ $[2.2, 2.6] \text{ GeV}/c^2 \rightarrow d\sigma_{\gamma\gamma}^{e^+e^-} = 154 \pm 11(\text{stat})_{-10.8}^{+16.6}(\text{syst}) \mu\text{b}$ precision 12%
 - ✧ $[3.7, 10] \text{ GeV}/c^2 \rightarrow d\sigma_{\gamma\gamma}^{e^+e^-} = 91 \pm 10(\text{stat})_{-8.0}^{+10.9}(\text{syst}) \mu\text{b}$ precision 16%
- ✓ the measured values for the $\gamma\gamma$ cross sections are 20% above but fully compatible within 1.0σ and 1.5σ with the STARLIGHT (LO) prediction for the low and high invariant mass intervals ($128 \mu\text{b}$ and $77 \mu\text{b}$)


\rightarrow the models predicting a strong contribution of higher-order terms (not included in STARLIGHT) are not favored

UPCs in pA




forward rapidity ($J/\psi \rightarrow \mu^+\mu^-$)

semi-forward rapidity ($J/\psi \rightarrow \mu^+\mu^-$)




mid-rapidity ($J/\psi \rightarrow \mu^+\mu^-$
and $J/\psi \rightarrow e^+e^-$)



trigger logic:

- ✓ similar to Pb+Pb case for forward and mid-rapidities, but improved purity
- ✓ semi-forward
 - ✧ veto on V0A and V0C (≥ 5 cells)
 - ✧ veto on SPD multiplicity (≥ 7 outer chips)
 - ✧ single muon with $p_T > 0.5 \text{ GeV}/c$
 - ✧ SPD (≥ 1 chips)

UPCs in pA




J/ψ photoproduction dominated by $\gamma+p$ process

- ✓ first results at forward rapidity
- ✓ analysis ongoing for central and semi-forward samples

accessible kinematics regions:

p+Pb forward	$21 < W_{\gamma p} < 45$ GeV
p+Pb semi-forward	$45 < W_{\gamma p} < 82$ GeV
mid-rapidity	$100 < W_{\gamma p} < 250$ GeV
Pb+p semi-forward	$300 < W_{\gamma p} < 550$ GeV
Pb+p forward	$550 < W_{\gamma p} < 1160$ GeV



possibility to study gluon PDFs in proton up to $x \sim 10^{-5}$


<http://indico.cern.ch/materialDisplay.py?contribId=129&sessionId=15&materialId=slides&confId=27458>

Conclusions


- ✓ LHC as γ Pb and γ p collider to study $\gamma\gamma$, photo-nuclear and γ p processes
- ✓ measurement of **exclusive vector meson** (J/ψ) cross sections to investigate the **gluon distribution in the nuclei**
- ✓ results seem to favor models including **gluon shadowing**
- ✓ $\gamma\gamma$ cross section to set limits on higher order terms in **QED processes**
- ✓ two ALICE papers:
 - ✧ Phys. Lett. B718 (2013) 1273-1283
 - ✧ arXiv:1305.1467 [nucl-ex]
- ✓ on going analyses:
 - ✧ J/ψ cross section in **p+Pb** and **Pb+p** collisions for three different topologies (central, forward and semi-forward) \rightarrow this allows J/ψ photoproduction measurement in γ p in a wide range of center of mass energy ([20,1000] GeV)
 - ✧ ρ^0 cross section in Pb+Pb collisions

back up

p_T distributions (linear scale)



dimuon channel



dielectron channel

p_T distribution fitted using MC samples representing several components:

- ✧ coherent and incoherent J/ ψ
- ✧ (coherent and incoherent) ψ' feed down
- ✧ $\gamma\gamma \rightarrow \mu^+\mu^-$
- ✧ hadronic

distribution peaked at low momentum as expected from coherent production

Feed down ($\psi' \rightarrow J/\Psi + \text{anything}$)

- ✓ fraction f_D of J/Ψ coming from the decay of $\psi' \rightarrow J/\Psi + \text{anything}$ estimated by simulating a sample of coherently produced ψ' with STARLIGHT, using PYTHIA to simulate their decay into J/Ψ
- ✓ contribution from incoherent ψ' expected to be negligible for the enriched coherent J/Ψ samples \rightarrow not considered
- ✓ ψ' polarization can be shared between J/Ψ and the other daughters $\rightarrow \psi'$ decay simulated assuming no polarization, full transverse and full longitudinal polarization for the J/Ψ

for a given polarization P:

$$f_D^P = \frac{\sigma_{\psi'} \cdot BR(\psi' \rightarrow J/\psi + \text{anything}) \cdot (\text{Acc} \times \epsilon)_{\psi' \rightarrow J/\psi}^P}{\sigma_{J/\psi} \cdot (\text{Acc} \times \epsilon)_{J/\psi}}$$

see table in the next slide for the results

Feed down ($\psi' \rightarrow J/\psi + \text{anything}$)

alternatively the ratio ψ' over J/ψ , used to compute the feed-down f_D , can be extracted from the data

due to the limited statistics the two decay channels were combined:


$$N_{\psi'} = 17 \pm 10 \text{ and } N_{J/\psi} = 505 \pm 48$$

$$f_D^P = \frac{N_{\psi'} \cdot BR(J/\psi \rightarrow l^+l^-) \cdot BR(\psi' \rightarrow J/\psi + \text{anything}) \cdot (\text{Acc} \times \epsilon)_{\psi' \rightarrow J/\psi}^P}{N_{J/\psi} \cdot BR(\psi' \rightarrow l^+l^-) (\text{Acc} \times \epsilon)_{\psi' \rightarrow l^+l^-}^P}$$

→ f_D ranges from $11.0 \pm 6.5\%$ for transverse ψ' polarization to $15 \pm 9\%$ for longitudinal ψ' polarization

the average of these estimates is $f_D = 0.10^{+0.05}_{-0.06}$

invariant mass distribution for combined dimuon and dielectron channels



Fit procedure

- ✓ exponential for underlying continuum
(systematics evaluated using polynomial)
- ✓ Crystall Ball (exp+gauss) to extract the J/Ψ signal
- ✓ tail CB parameters (α and n) left free for the coherent sample (systematics evaluated fixing the parameters) and fixed to MC values for the incoherent one
- ✓ incoherent dimuons fitted also using a polynomial to take into account the combinatorial background, as constrained to the LS pair spectrum
- ✓ fit also constrained to a MC cocktail ($J/\Psi + \gamma\gamma$)